



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:16 AM UTC

PDB ID : 4ZDS / pdb_00004zds
Title : Crystal Structure of core DNA binding domain of Arabidopsis Thaliana Transcription Factor Ethylene-Insensitive 3
Authors : Song, J.; Zhu, C.; Zhang, X.; Wen, X.; Liu, L.; Peng, J.; Guo, H.; Yi, C.
Deposited on : 2015-04-18
Resolution : 1.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1365 (1.78-1.78)
Clashscore	190562	1395 (1.78-1.78)
Ramachandran outliers	187476	1382 (1.78-1.78)
Sidechain outliers	187428	1382 (1.78-1.78)
RSRZ outliers	180081	1365 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	134	
1	B	134	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2250 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein ETHYLENE INSENSITIVE 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	132	Total	C	N	O	S	0	1	0
			1071	692	186	188	5			
1	B	129	Total	C	N	O	S	0	0	0
			1038	668	183	182	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	173	SER	-	expression tag	UNP O24606
B	173	SER	-	expression tag	UNP O24606



- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	74	Total	O	0	0
			74	74		
2	B	67	Total	O	0	0
			67	67		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protein ETHYLENE INSENSITIVE 3



- Molecule 1: Protein ETHYLENE INSENSITIVE 3



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	35.02Å 45.17Å 47.57Å 89.46° 77.07° 71.38°	Depositor
Resolution (Å)	46.26 – 1.78 46.26 – 1.78	Depositor EDS
% Data completeness (in resolution range)	96.6 (46.26-1.78) 96.6 (46.26-1.78)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.92 (at 1.78Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.174 , 0.218 (Not available) , 0.237	Depositor DCC
R_{free} test set	1270 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	26.6	Xtriage
Anisotropy	0.152	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 33.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2250	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.27	2/1108 (0.2%)	1.16	3/1508 (0.2%)
1	B	1.24	1/1070 (0.1%)	1.38	13/1455 (0.9%)
All	All	1.26	3/2178 (0.1%)	1.27	16/2963 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	4
All	All	0	8

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	240	PRO	C-O	5.54	1.31	1.24
1	A	200	PRO	C-O	5.52	1.30	1.23
1	A	174	THR	C-O	-5.47	1.19	1.24

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	198	ASP	CA-C-N	14.53	135.35	120.38
1	B	198	ASP	C-N-CA	14.53	135.35	120.38
1	B	260	PHE	CA-C-N	11.41	134.10	119.84
1	B	260	PHE	C-N-CA	11.41	134.10	119.84
1	B	299	ALA	N-CA-C	-8.45	99.37	110.53
1	A	260	PHE	CA-C-N	7.79	129.58	119.84
1	A	260	PHE	C-N-CA	7.79	129.58	119.84
1	B	199	PRO	N-CA-C	-6.31	103.00	110.70
1	B	191	SER	N-CA-C	5.86	117.67	111.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	270	VAL	N-CA-C	5.83	116.01	110.53
1	A	198	ASP	C-N-CD	-5.73	107.98	120.60
1	B	198	ASP	C-N-CD	-5.59	102.08	125.00
1	B	261	PRO	N-CA-C	-5.47	101.20	112.47
1	B	250	GLY	N-CA-C	5.38	118.74	112.50
1	B	300	ARG	N-CA-C	-5.27	99.58	110.80
1	B	260	PHE	C-N-CD	-5.23	103.58	125.00

There are no chirality outliers.

All (8) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	198	ASP	Mainchain,Peptide
1	A	260	PHE	Mainchain,Peptide
1	B	198	ASP	Mainchain,Peptide
1	B	260	PHE	Mainchain,Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1071	0	1090	11	0
1	B	1038	0	1057	13	0
2	A	74	0	0	0	2
2	B	67	0	0	1	2
All	All	2250	0	2147	23	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:299:ALA:O	1:B:300:ARG:HB2	1.73	0.88
1:B:299:ALA:O	1:B:300:ARG:CB	2.39	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:207[B]:GLU:H	1:A:207[B]:GLU:CD	1.99	0.70
1:B:195:GLN:HE22	1:B:201:GLN:HE21	1.41	0.69
1:A:298:LEU:HA	1:A:301:GLU:OE1	1.94	0.68
1:A:195:GLN:OE1	1:A:202:ARG:HD3	1.95	0.67
1:B:207:GLU:H	1:B:207:GLU:CD	2.08	0.60
1:A:280:MET:CE	1:A:284:GLU:HB3	2.39	0.53
1:B:232:GLN:HG2	2:B:404:HOH:O	2.07	0.52
1:B:244:LYS:HG2	1:B:247:TRP:CE3	2.44	0.51
1:A:280:MET:HE2	1:A:284:GLU:C	2.37	0.49
1:B:296:GLU:O	1:B:299:ALA:O	2.32	0.48
1:B:221:ASP:OD1	1:B:221:ASP:N	2.34	0.48
1:A:280:MET:HE2	1:A:284:GLU:HB3	1.96	0.46
1:A:292:ILE:HD13	1:A:292:ILE:HA	1.80	0.45
1:B:195:GLN:HE22	1:B:201:GLN:NE2	2.12	0.44
1:A:274:LYS:HD2	1:A:274:LYS:HA	1.77	0.44
1:A:232:GLN:HG3	1:B:275:CYS:SG	2.58	0.43
1:A:244:LYS:HG2	1:A:247:TRP:CE3	2.54	0.43
1:B:223:TRP:N	1:B:224:PRO:CD	2.82	0.42
1:B:176:HIS:CE1	1:B:225:GLN:HB3	2.56	0.40
1:B:203:ARG:O	1:B:205:PRO:HD3	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:450:HOH:O	2:B:405:HOH:O[1_556]	1.03	1.17
2:A:465:HOH:O	2:B:463:HOH:O[1_556]	1.60	0.60

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	131/134 (98%)	129 (98%)	1 (1%)	1 (1%)	16	6
1	B	127/134 (95%)	124 (98%)	0	3 (2%)	4	0
All	All	258/268 (96%)	253 (98%)	1 (0%)	4 (2%)	7	1

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	300	ARG
1	B	261	PRO
1	B	199	PRO
1	A	261	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	118/119 (99%)	110 (93%)	8 (7%)	14	2
1	B	114/119 (96%)	109 (96%)	5 (4%)	25	7
All	All	232/238 (98%)	219 (94%)	13 (6%)	21	4

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	203	ARG
1	A	207[A]	GLU
1	A	207[B]	GLU
1	A	272	GLN
1	A	292	ILE
1	A	296	GLU
1	A	298	LEU
1	A	302	LEU
1	B	207	GLU
1	B	221	ASP
1	B	272	GLN
1	B	298	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	301	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	196	HIS
1	B	201	GLN
1	B	272	GLN
1	B	294	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	132/134 (98%)	0.09	4 (3%) 52 60	17, 29, 49, 63	1 (0%)
1	B	129/134 (96%)	0.15	2 (1%) 70 77	18, 33, 50, 61	0
All	All	261/268 (97%)	0.12	6 (2%) 61 69	17, 31, 49, 63	1 (0%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	304	PRO	3.6
1	B	275	CYS	2.6
1	A	176	HIS	2.5
1	A	302	LEU	2.2
1	A	301	GLU	2.1
1	B	298	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.