



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 07:37 PM UTC

PDB ID : 4ZE5 / pdb_00004ze5
Title : Structure of Gan1D-E170Q, a catalytic mutant of a putative 6-phospho-beta-galactosidase from *Geobacillus stearothermophilus*
Authors : Lansky, S.; Zehavi, A.; Dvir, H.; Shoham, Y.; Shoham, G.
Deposited on : 2015-04-20
Resolution : 1.48 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6779 (1.50-1.46)
Clashscore	190562	7025 (1.50-1.46)
Ramachandran outliers	187476	6917 (1.50-1.46)
Sidechain outliers	187428	6914 (1.50-1.46)
RSRZ outliers	180081	6781 (1.50-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	485	2% 77% 16% • 5%
1	B	485	% 80% 14% • 5%
1	C	485	2% 78% 15% • 5%
1	D	485	2% 74% 18% • 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	IMD	A	505	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 18228 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative 6-phospho-beta-galactobiosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	459	Total	C	N	O	S	0	21	0
			3902	2518	666	703	15			
1	B	459	Total	C	N	O	S	0	20	0
			3900	2515	662	709	14			
1	C	460	Total	C	N	O	S	0	23	0
			3923	2530	674	705	14			
1	D	460	Total	C	N	O	S	0	16	0
			3872	2497	664	699	12			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	MET	-	initiating methionine	UNP W8QF82
A	-5	ILE	-	expression tag	UNP W8QF82
A	-4	HIS	-	expression tag	UNP W8QF82
A	-3	HIS	-	expression tag	UNP W8QF82
A	-2	HIS	-	expression tag	UNP W8QF82
A	-1	HIS	-	expression tag	UNP W8QF82
A	0	HIS	-	expression tag	UNP W8QF82
A	1	HIS	-	expression tag	UNP W8QF82
A	170	GLN	GLU	engineered mutation	UNP W8QF82
B	-6	MET	-	initiating methionine	UNP W8QF82
B	-5	ILE	-	expression tag	UNP W8QF82
B	-4	HIS	-	expression tag	UNP W8QF82
B	-3	HIS	-	expression tag	UNP W8QF82
B	-2	HIS	-	expression tag	UNP W8QF82
B	-1	HIS	-	expression tag	UNP W8QF82
B	0	HIS	-	expression tag	UNP W8QF82
B	1	HIS	-	expression tag	UNP W8QF82
B	170	GLN	GLU	engineered mutation	UNP W8QF82
C	-6	MET	-	initiating methionine	UNP W8QF82
C	-5	ILE	-	expression tag	UNP W8QF82
C	-4	HIS	-	expression tag	UNP W8QF82

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-3	HIS	-	expression tag	UNP W8QF82
C	-2	HIS	-	expression tag	UNP W8QF82
C	-1	HIS	-	expression tag	UNP W8QF82
C	0	HIS	-	expression tag	UNP W8QF82
C	1	HIS	-	expression tag	UNP W8QF82
C	170	GLN	GLU	engineered mutation	UNP W8QF82
D	-6	MET	-	initiating methionine	UNP W8QF82
D	-5	ILE	-	expression tag	UNP W8QF82
D	-4	HIS	-	expression tag	UNP W8QF82
D	-3	HIS	-	expression tag	UNP W8QF82
D	-2	HIS	-	expression tag	UNP W8QF82
D	-1	HIS	-	expression tag	UNP W8QF82
D	0	HIS	-	expression tag	UNP W8QF82
D	1	HIS	-	expression tag	UNP W8QF82
D	170	GLN	GLU	engineered mutation	UNP W8QF82

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is IMIDAZOLE (CCD ID: IMD) (formula: $C_3H_5N_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			5	3	2		
3	C	1	Total	C	N	0	0
			5	3	2		
3	D	1	Total	C	N	0	0
			5	3	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	616	Total 616	O 616	0	0
4	B	640	Total 640	O 640	0	0
4	C	690	Total 690	O 690	0	0
4	D	610	Total 610	O 610	0	0



i

- Molecule 1: Putative 6-phospho-beta-galactobiosidase





● Molecule 1: Putative 6-phospho-beta-galactobiosidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	101.55Å 97.44Å 105.24Å 90.00° 97.82° 90.00°	Depositor
Resolution (Å)	27.64 – 1.48 27.64 – 1.48	Depositor EDS
% Data completeness (in resolution range)	99.7 (27.64-1.48) 99.7 (27.64-1.48)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.50 (at 1.48Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.132 , 0.159 0.132 , 0.159	Depositor DCC
R_{free} test set	16968 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	10.9	Xtriage
Anisotropy	0.482	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.000 for l,-k,h	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	18228	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 71.27 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.7039e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, IMD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.70	50/4091 (1.2%)	1.47	30/5559 (0.5%)
1	B	1.65	31/4076 (0.8%)	1.48	30/5543 (0.5%)
1	C	1.63	35/4117 (0.9%)	1.44	25/5597 (0.4%)
1	D	1.70	42/4041 (1.0%)	1.56	42/5496 (0.8%)
All	All	1.67	158/16325 (1.0%)	1.49	127/22195 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	2
All	All	0	6

The worst 5 of 158 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	430	LEU	C-O	-9.11	1.12	1.23
1	C	476	ALA	C-O	8.42	1.34	1.24
1	D	440	ARG	CA-C	-8.02	1.42	1.52
1	B	133	ASP	CG-OD1	-7.99	1.10	1.25
1	C	309	ASN	C-O	-7.96	1.17	1.24

The worst 5 of 127 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	93	ASN	CA-C-N	-8.66	105.81	121.32
1	B	93	ASN	C-N-CA	-8.66	105.81	121.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	477	GLU	N-CA-CB	8.41	124.34	111.56
1	A	468	GLN	O-C-N	8.37	130.99	122.12
1	C	352	TRP	CA-CB-CG	7.94	128.69	113.60

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	157[A]	ARG	Sidechain
1	A	157[B]	ARG	Sidechain
1	A	476	ALA	Peptide
1	A	477	GLU	Peptide
1	B	349[A]	ASN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3902	0	3768	33	0
1	B	3900	0	3731	25	0
1	C	3923	0	3783	40	0
1	D	3872	0	3716	50	0
2	A	24	0	32	3	0
2	B	18	0	24	0	0
2	C	12	0	16	0	0
2	D	6	0	8	0	0
3	A	5	0	5	6	0
3	C	5	0	5	0	0
3	D	5	0	5	1	0
4	A	616	0	0	11	0
4	B	640	0	0	9	0
4	C	690	0	0	22	0
4	D	610	0	0	18	0
All	All	18228	0	15093	147	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 147 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:132[A]:MET:HE3	4:D:1075:HOH:O	1.54	1.08
1:A:453:GLU:OE1	3:A:505:IMD:H2	1.59	1.01
1:C:132[A]:MET:HE3	4:C:1110:HOH:O	1.60	1.01
1:B:349[B]:ASN:O	1:B:350[B]:TRP:CD1	2.15	0.99
1:C:149:ARG:HD3	4:C:1004:HOH:O	1.64	0.95

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	478/485 (99%)	462 (97%)	16 (3%)	0	100	100
1	B	476/485 (98%)	461 (97%)	15 (3%)	0	100	100
1	C	480/485 (99%)	462 (96%)	17 (4%)	1 (0%)	43	22
1	D	472/485 (97%)	456 (97%)	14 (3%)	2 (0%)	30	11
All	All	1906/1940 (98%)	1841 (97%)	62 (3%)	3 (0%)	43	22

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	310	PRO
1	C	310	PRO
1	D	167	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	413/412 (100%)	410 (99%)	3 (1%)	76	56
1	B	411/412 (100%)	408 (99%)	3 (1%)	76	56
1	C	415/412 (101%)	411 (99%)	4 (1%)	68	43
1	D	407/412 (99%)	400 (98%)	7 (2%)	53	24
All	All	1646/1648 (100%)	1629 (99%)	17 (1%)	68	43

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	112[B]	ARG
1	D	430	LEU
1	C	312	ASP
1	C	329	THR
1	C	430	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 56 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	170	GLN
1	D	410	GLN
1	C	269	ASN
1	D	349	ASN
1	D	251	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	B	501	-	5,5,5	0.49	0	5,5,5	1.72	2 (40%)
2	GOL	D	501	-	5,5,5	0.44	0	5,5,5	0.94	0
2	GOL	B	502	-	5,5,5	0.68	0	5,5,5	0.92	0
2	GOL	A	501	-	5,5,5	0.77	0	5,5,5	1.31	0
3	IMD	C	503	-	5,5,5	0.86	0	5,5,5	0.65	0
2	GOL	C	501	-	5,5,5	0.68	0	5,5,5	1.20	1 (20%)
2	GOL	A	503	-	5,5,5	0.41	0	5,5,5	1.27	0
2	GOL	C	502	-	5,5,5	0.49	0	5,5,5	1.01	0
3	IMD	D	502	-	5,5,5	0.93	0	5,5,5	0.73	0
3	IMD	A	505	-	5,5,5	0.74	0	5,5,5	1.48	1 (20%)
2	GOL	A	502	-	5,5,5	0.36	0	5,5,5	1.03	1 (20%)
2	GOL	A	504	-	5,5,5	0.65	0	5,5,5	1.01	0
2	GOL	B	503	-	5,5,5	0.50	0	5,5,5	1.00	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	501	-	-	0/4/4/4	-
2	GOL	D	501	-	-	0/4/4/4	-
2	GOL	B	502	-	-	0/4/4/4	-
2	GOL	A	501	-	-	0/4/4/4	-
3	IMD	C	503	-	-	-	0/1/1/1
2	GOL	C	501	-	-	2/4/4/4	-
2	GOL	A	503	-	-	0/4/4/4	-
2	GOL	C	502	-	-	2/4/4/4	-
3	IMD	D	502	-	-	-	0/1/1/1
3	IMD	A	505	-	-	-	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	502	-	-	0/4/4/4	-
2	GOL	A	504	-	-	3/4/4/4	-
2	GOL	B	503	-	-	0/4/4/4	-

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	GOL	C3-C2-C1	-2.93	101.05	111.80
2	C	501	GOL	C3-C2-C1	-2.50	102.62	111.80
2	B	501	GOL	O1-C1-C2	2.36	120.99	110.38
2	A	502	GOL	O3-C3-C2	2.22	120.38	110.38
3	A	505	IMD	C5-N1-C2	-2.05	102.77	107.19

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	501	GOL	O1-C1-C2-C3
2	C	501	GOL	O1-C1-C2-O2
2	A	504	GOL	C1-C2-C3-O3
2	A	504	GOL	O2-C2-C3-O3
2	A	504	GOL	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	502	IMD	1	0
3	A	505	IMD	6	0
2	A	504	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	459/485 (94%)	-0.33	12 (2%) 57 61	5, 11, 28, 67	21 (4%)
1	B	459/485 (94%)	-0.41	7 (1%) 72 76	4, 10, 23, 61	20 (4%)
1	C	460/485 (94%)	-0.49	8 (1%) 69 73	4, 9, 21, 80	23 (5%)
1	D	460/485 (94%)	-0.32	12 (2%) 57 61	5, 11, 25, 90	16 (3%)
All	All	1838/1940 (94%)	-0.38	39 (2%) 63 67	4, 10, 25, 90	80 (4%)

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	350	TRP	6.6
1	D	352	TRP	5.6
1	B	352	TRP	5.2
1	A	352	TRP	5.1
1	C	313	GLY	5.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	IMD	D	502	5/5	0.77	0.16	19,25,31,32	5
2	GOL	B	503	6/6	0.80	0.13	29,34,35,35	0
3	IMD	A	505	5/5	0.81	0.13	33,38,42,45	0
2	GOL	A	503	6/6	0.81	0.14	31,32,35,35	0
2	GOL	D	501	6/6	0.82	0.14	34,35,37,40	0
2	GOL	C	502	6/6	0.86	0.10	30,31,32,33	0
3	IMD	C	503	5/5	0.87	0.13	22,26,31,35	0
2	GOL	A	504	6/6	0.87	0.12	38,40,43,47	0
2	GOL	C	501	6/6	0.93	0.10	17,26,33,44	0
2	GOL	B	502	6/6	0.97	0.07	13,15,21,23	0
2	GOL	A	501	6/6	0.97	0.07	13,15,20,22	0
2	GOL	A	502	6/6	0.97	0.07	13,17,20,22	0
2	GOL	B	501	6/6	0.97	0.08	13,17,22,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.