



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 06:37 PM UTC

PDB ID : 4ZEB / pdb_00004zeb
Title : PBP AccA from *A. tumefaciens* C58 in complex with agrocinopine A
Authors : El Sahili, A.; Morera, S.
Deposited on : 2015-04-20
Resolution : 1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	499	% 91% 7% ..
1	B	499	% 91% 6% ..
2	C	2	100%
2	D	2	50% 50%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8520 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ABC transporter, substrate binding protein (Agrocinopines A and B).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	490	Total	C	N	O	S	0	4	0
			3959	2555	662	723	19			
1	B	489	Total	C	N	O	S	0	2	0
			3950	2549	661	720	20			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	29	MET	-	initiating methionine	UNP Q7D2F4
A	522	HIS	-	expression tag	UNP Q7D2F4
A	523	HIS	-	expression tag	UNP Q7D2F4
A	524	HIS	-	expression tag	UNP Q7D2F4
A	525	HIS	-	expression tag	UNP Q7D2F4
A	526	HIS	-	expression tag	UNP Q7D2F4
A	527	HIS	-	expression tag	UNP Q7D2F4
B	29	MET	-	initiating methionine	UNP Q7D2F4
B	522	HIS	-	expression tag	UNP Q7D2F4
B	523	HIS	-	expression tag	UNP Q7D2F4
B	524	HIS	-	expression tag	UNP Q7D2F4
B	525	HIS	-	expression tag	UNP Q7D2F4
B	526	HIS	-	expression tag	UNP Q7D2F4
B	527	HIS	-	expression tag	UNP Q7D2F4

- Molecule 2 is an oligosaccharide called alpha-D-gulopyranose-(1-2)-4-O-phosphono-beta-D-f ructofuranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	O	P	0	0	0
			26	12	13	1			
2	D	2	Total	C	O	P	0	0	0
			26	12	13	1			

- Molecule 3 is alpha-L-arabinopyranose (CCD ID: ARA) (formula: C₅H₁₀O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	5	5		
3	B	1	Total	C	O	0	0
			10	5	5		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			7	4	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	275	Total 275	O 275	0	0
6	B	253	Total 253	O 253	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ABC transporter, substrate binding protein (Agrocinopines A and B)



- Molecule 1: ABC transporter, substrate binding protein (Agrocinopines A and B)



- Molecule 2: alpha-D-gulopyranose-(1-2)-4-O-phosphono-beta-D-fructofuranose



- Molecule 2: alpha-D-gulopyranose-(1-2)-4-O-phosphono-beta-D-fructofuranose



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	177.21Å 51.24Å 119.95Å 90.00° 114.33° 90.00°	Depositor
Resolution (Å)	37.80 – 1.89 37.80 – 1.89	Depositor EDS
% Data completeness (in resolution range)	98.7 (37.80-1.89) 99.2 (37.80-1.89)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 1.90Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.172 , 0.200 (Not available) , 0.200	Depositor DCC
R_{free} test set	3903 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	30.7	Xtriage
Anisotropy	0.363	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 41.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8520	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, PEG, ARA, TT7, 4GL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	1/4080 (0.0%)	1.15	10/5542 (0.2%)
1	B	0.83	3/4064 (0.1%)	1.15	12/5518 (0.2%)
All	All	0.83	4/8144 (0.0%)	1.15	22/11060 (0.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	484	ALA	CA-C	8.24	1.59	1.53
1	A	372	MET	SD-CE	-6.66	1.62	1.79
1	B	372	MET	SD-CE	-5.58	1.65	1.79
1	B	404	PRO	C-O	-5.04	1.17	1.24

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	156	ASP	CA-CB-CG	7.74	120.34	112.60
1	B	423	TRP	N-CA-C	7.42	122.18	113.20
1	A	423	TRP	N-CA-C	7.14	121.84	113.20
1	B	180	SER	N-CA-C	6.99	121.12	112.59
1	A	180	SER	N-CA-C	6.92	121.04	112.59
1	B	136	ILE	N-CA-C	6.38	117.90	110.62
1	A	130	LEU	N-CA-C	6.30	120.77	112.13
1	B	156	ASP	CA-CB-CG	6.00	118.61	112.60
1	A	211	PHE	CA-CB-CG	5.70	119.50	113.80
1	A	266	ILE	N-CA-C	5.61	116.06	110.23
1	B	149	PHE	N-CA-C	5.60	117.97	110.35
1	A	149	PHE	N-CA-C	5.50	117.83	110.35
1	A	420	ASN	CA-CB-CG	5.47	118.07	112.60
1	B	266	ILE	N-CA-C	5.43	115.88	110.23
1	B	211	PHE	CA-CB-CG	5.41	119.21	113.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	110	PHE	N-CA-C	-5.31	103.38	110.55
1	B	130	LEU	N-CA-C	5.26	119.33	112.13
1	A	47	GLU	N-CA-C	-5.17	100.54	109.58
1	B	150	ASP	N-CA-C	-5.14	102.59	110.14
1	A	150	ASP	N-CA-C	-5.13	102.60	110.14
1	B	47	GLU	N-CA-C	-5.10	100.66	109.58
1	B	192	GLY	N-CA-C	-5.07	106.35	112.48

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3959	0	3914	17	0
1	B	3950	0	3900	18	0
2	C	26	0	0	0	0
2	D	26	0	0	1	0
3	A	10	0	9	0	0
3	B	10	0	9	0	0
4	A	4	0	6	1	0
5	B	7	0	10	0	0
6	A	275	0	0	2	0
6	B	253	0	0	0	0
All	All	8520	0	7848	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:405:GLY:HA3	1:B:406:SER:HB2	1.45	0.98
1:B:403:ALA:O	1:B:406:SER:HB3	1.75	0.87
1:A:67:LEU:HD12	1:A:175[B]:VAL:HG21	1.68	0.74

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:405:GLY:CA	1:B:406:SER:HB2	2.18	0.72
1:A:67:LEU:HD12	1:A:175[B]:VAL:CG2	2.29	0.62
1:A:370:HIS:HD2	6:A:749:HOH:O	1.87	0.58
1:B:372:MET:HE3	1:B:402:TYR:CE1	2.40	0.57
1:A:212[B]:VAL:HG13	1:A:215:ASP:HB3	1.86	0.57
1:A:372:MET:HE2	1:A:375:TYR:HB2	1.86	0.56
1:B:272:TYR:HB3	1:B:275:LEU:HD13	1.87	0.56
1:B:294:GLN:HE21	1:B:294:GLN:HA	1.73	0.54
1:B:403:ALA:O	1:B:406:SER:CB	2.53	0.53
1:B:50:ASN:HD21	1:B:139:ILE:HA	1.75	0.52
1:A:95:ILE:HD11	1:A:101:ARG:HB2	1.92	0.51
1:A:335:TYR:OH	1:A:422:GLN:NE2	2.45	0.50
1:B:403:ALA:O	1:B:405:GLY:N	2.44	0.50
1:B:372:MET:HE3	1:B:402:TYR:HE1	1.77	0.49
1:B:68:ILE:HD12	1:B:175[B]:VAL:HG23	1.95	0.48
1:B:335:TYR:OH	1:B:422:GLN:NE2	2.46	0.48
1:B:422:GLN:HE22	1:B:430:THR:H	1.61	0.48
1:A:136:ILE:HD11	1:A:143:LYS:HD3	1.96	0.47
1:A:199:LYS:HE2	4:A:602:EDO:H12	1.96	0.47
1:B:372:MET:HE1	2:D:2:4GL:O2	2.14	0.46
1:A:150:ASP:HB3	1:A:163:ARG:HG3	1.98	0.46
1:A:422:GLN:HE22	1:A:430:THR:H	1.62	0.46
1:B:403:ALA:C	1:B:405:GLY:N	2.72	0.44
1:B:150:ASP:HB3	1:B:163:ARG:HG3	1.98	0.44
1:A:37:LEU:HD22	1:A:257:ILE:HB	2.00	0.43
1:B:405:GLY:CA	1:B:406:SER:CB	2.90	0.43
1:A:246:ARG:HG2	1:A:258:ILE:HB	2.01	0.43
1:A:440:GLN:NE2	6:A:701:HOH:O	2.51	0.43
1:A:128:GLU:HA	1:A:134:GLU:CG	2.49	0.42
1:B:37:LEU:HD22	1:B:257:ILE:HB	2.00	0.42
1:A:68:ILE:HD12	1:A:175[B]:VAL:HG13	2.01	0.42
1:A:429:ALA:HA	1:A:430:THR:HA	1.80	0.42

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	492/499 (99%)	475 (96%)	17 (4%)	0	100	100
1	B	487/499 (98%)	469 (96%)	16 (3%)	2 (0%)	30	22
All	All	979/998 (98%)	944 (96%)	33 (3%)	2 (0%)	43	36

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	406	SER
1	B	405	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	423/428 (99%)	421 (100%)	2 (0%)	81	84
1	B	421/428 (98%)	419 (100%)	2 (0%)	81	84
All	All	844/856 (99%)	840 (100%)	4 (0%)	81	84

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	156	ASP
1	A	260	THR
1	B	82	ILE
1	B	260	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	293	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	370	HIS
1	A	420	ASN
1	A	422	GLN
1	A	440	GLN
1	A	442	GLN
1	A	454	ASN
1	A	491	GLN
1	B	50	ASN
1	B	294	GLN
1	B	334	ASN
1	B	396	ASN
1	B	420	ASN
1	B	422	GLN
1	B	438	ASN
1	B	440	GLN
1	B	491	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	TT7	C	1	2	11,15,16	2.31	2 (18%)	10,22,25	1.40	2 (20%)
2	4GL	C	2	2	11,11,12	1.76	2 (18%)	15,15,17	2.48	7 (46%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	TT7	D	1	2	11,15,16	2.43	3 (27%)	10,22,25	1.41	1 (10%)
2	4GL	D	2	2	11,11,12	1.86	2 (18%)	15,15,17	2.34	5 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TT7	C	1	2	-	3/7/28/29	0/1/1/1
2	4GL	C	2	2	-	2/2/19/22	0/1/1/1
2	TT7	D	1	2	-	3/7/28/29	0/1/1/1
2	4GL	D	2	2	-	1/2/19/22	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1	TT7	C4-C3	-6.01	1.39	1.53
2	C	1	TT7	C4-C3	-5.86	1.40	1.53
2	D	2	4GL	O5-C5	4.62	1.52	1.43
2	C	2	4GL	O5-C5	4.48	1.52	1.43
2	D	1	TT7	O2-C2	3.81	1.47	1.40
2	C	1	TT7	O2-C2	3.73	1.47	1.40
2	D	2	4GL	C4-C3	-2.67	1.45	1.52
2	C	2	4GL	C4-C3	-2.64	1.45	1.52
2	D	1	TT7	O5-C2	-2.27	1.39	1.43

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	4GL	C3-C4-C5	4.44	118.28	110.23
2	C	2	4GL	C2-C3-C4	4.16	118.19	110.86
2	D	2	4GL	C1-O5-C5	4.13	117.72	112.19
2	D	2	4GL	C2-C3-C4	3.86	117.66	110.86
2	D	2	4GL	C1-C2-C3	3.83	115.22	109.64
2	D	2	4GL	C3-C4-C5	3.73	116.99	110.23
2	C	2	4GL	C1-C2-C3	3.62	114.91	109.64
2	C	2	4GL	C1-O5-C5	3.45	116.81	112.19
2	C	2	4GL	O4-C4-C5	3.39	117.68	109.32
2	D	1	TT7	O1-C1-C2	3.05	118.41	111.67
2	C	1	TT7	O2-C2-O5	2.54	114.20	109.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	TT7	O1-C1-C2	2.44	117.06	111.67
2	D	2	4GL	O2-C2-C3	2.41	115.14	110.15
2	C	2	4GL	O6-C6-C5	2.24	118.95	111.33
2	C	2	4GL	C6-C5-C4	2.16	118.32	113.02

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	1	TT7	O1-C1-C2-O2
2	D	1	TT7	O1-C1-C2-C3
2	D	1	TT7	O1-C1-C2-O5
2	C	2	4GL	C4-C5-C6-O6
2	D	2	4GL	C4-C5-C6-O6
2	C	1	TT7	O1-C1-C2-O2
2	C	1	TT7	O1-C1-C2-C3
2	C	1	TT7	O1-C1-C2-O5
2	C	2	4GL	O5-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	2	4GL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	PEG	B	602	-	6,6,6	0.24	0	5,5,5	0.11	0
3	ARA	A	601	-	10,10,10	2.22	2 (20%)	14,14,14	1.11	1 (7%)
4	EDO	A	602	-	3,3,3	0.59	0	2,2,2	0.12	0
3	ARA	B	601	-	10,10,10	2.23	3 (30%)	14,14,14	1.35	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ARA	A	601	-	-	-	0/1/1/1
4	EDO	A	602	-	-	0/1/1/1	-
5	PEG	B	602	-	-	2/4/4/4	-
3	ARA	B	601	-	-	-	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	ARA	C4-C3	-5.95	1.43	1.52
3	B	601	ARA	C4-C3	-5.74	1.43	1.52
3	B	601	ARA	C3-C2	-3.04	1.44	1.52
3	A	601	ARA	C3-C2	-2.78	1.45	1.52
3	B	601	ARA	O3-C3	2.25	1.48	1.43

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	ARA	C4-C3-C2	3.08	116.28	110.86
3	B	601	ARA	C4-C3-C2	2.28	114.87	110.86
3	B	601	ARA	O4-C4-C5	-2.06	104.50	109.22
3	B	601	ARA	O2-C2-C1	-2.05	104.52	109.25

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	602	PEG	C1-C2-O2-C3
5	B	602	PEG	C4-C3-O2-C2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	602	EDO	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	490/499 (98%)	-0.03	5 (1%) 79 82	15, 33, 56, 90	4 (0%)
1	B	489/499 (97%)	0.06	4 (0%) 82 85	18, 34, 58, 92	2 (0%)
All	All	979/998 (98%)	0.02	9 (0%) 81 83	15, 33, 57, 92	6 (0%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	252	SER	3.2
1	B	521	GLY	3.2
1	A	521	GLY	2.8
1	B	223	ASP	2.5
1	A	252	SER	2.5
1	A	268	LEU	2.4
1	A	84	LEU	2.3
1	A	251	ILE	2.2
1	B	407	PHE	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	4GL	D	2	11/12	0.89	0.10	42,45,47,51	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	4GL	C	2	11/12	0.94	0.07	38,41,45,46	0
2	TT7	D	1	15/16	0.96	0.07	22,37,45,48	0
2	TT7	C	1	15/16	0.98	0.06	23,35,38,40	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	EDO	A	602	4/4	0.77	0.15	63,63,63,64	0
5	PEG	B	602	7/7	0.80	0.14	50,52,56,58	0
3	ARA	A	601	10/10	0.96	0.06	23,26,28,28	0
3	ARA	B	601	10/10	0.96	0.06	26,29,29,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.