



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 06:15 AM UTC

PDB ID : 6ZE7 / pdb_00006ze7
Title : Chaetomium thermophilum FAD-dependent oxidoreductase in complex with 4-nitrophenol
Authors : Svecova, L.; Skalova, T.; Kolenko, P.; Koval, T.; Oestergaard, L.H.; Dohnalek, J.
Deposited on : 2020-06-16
Resolution : 1.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	595	% 94% ..
1	B	595	2% 92% 6% .
2	C	2	50% 50%
2	D	2	50% 50%

2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 11203 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FAD-dependent oxidoreductase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	584	Total	C	N	O	S	0	38	0
			4665	2969	810	870	16			
1	B	586	Total	C	N	O	S	0	36	0
			4661	2965	807	874	15			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is DIHYDROFLAVINE-ADENINE DINUCLEOTIDE (CCD ID: FDA) (formula: C₂₇H₃₅N₉O₁₅P₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	1
			16	9	1	6		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is P-NITROPHENOL (CCD ID: NPO) (formula: $C_6H_5NO_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	1
			10	6	1	3		
5	A	1	Total	C	N	O	0	0
			10	6	1	3		
5	B	1	Total	C	N	O	0	1
			10	6	1	3		

- Molecule 6 is FORMIC ACID (CCD ID: FMT) (formula: CH_2O_2).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 3 1 2	0	0
6	A	1	Total C O 3 1 2	0	0
6	A	1	Total C O 3 1 2	0	0
6	B	1	Total C O 3 1 2	0	0
6	B	1	Total C O 3 1 2	0	0
6	B	1	Total C O 3 1 2	0	0
6	B	1	Total C O 3 1 2	0	0
6	B	1	Total C O 3 1 2	0	0

- Molecule 7 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			13	8	5		
8	B	1	Total	C	O	0	0
			13	8	5		

- Molecule 9 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	C	O	0	0
			10	6	4		
9	B	1	Total	C	O	0	0
			10	6	4		

- Molecule 10 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	3	Total	Mg	0	0
			3	3		
10	B	2	Total	Mg	0	0
			2	2		

- Molecule 11 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	1	Total	Na	0	0
			1	1		


- Molecule 12 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	A	735	Total	O	0	20
			752	752		
12	B	754	Total	O	0	15
			763	763		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: FAD-dependent oxidoreductase



- Molecule 1: FAD-dependent oxidoreductase



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	92.61Å 109.67Å 115.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.23 – 1.50 47.23 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.3 (47.23-1.50) 99.3 (47.23-1.50)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.05 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.155 , 0.182 0.156 , 0.161	Depositor DCC
R_{free} test set	9455 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	14.3	Xtriage
Anisotropy	0.249	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 40.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	11203	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 70.87 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.8998e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, NPO, PG4, MG, FDA, NAG, FMT, PGE, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.06	2/4887 (0.0%)	1.17	5/6664 (0.1%)
1	B	1.10	2/4876 (0.0%)	1.18	4/6649 (0.1%)
All	All	1.08	4/9763 (0.0%)	1.18	9/13313 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	234	VAL	C-O	5.27	1.30	1.24
1	A	558	ILE	C-O	5.20	1.29	1.24
1	B	627	HIS	CE1-NE2	5.11	1.37	1.32
1	B	472	SER	CA-CB	-5.08	1.44	1.53

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	172	PRO	O-C-N	7.24	124.64	121.31
1	B	556	GLU	CB-CG-CD	6.33	123.36	112.60
1	A	217	GLU	CB-CG-CD	6.04	122.86	112.60
1	A	132	ASN	CA-CB-CG	5.65	118.25	112.60
1	B	217	GLU	CB-CG-CD	5.46	121.87	112.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	566	CYS	Peptide
1	B	566	CYS	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4665	0	4702	20	0
1	B	4661	0	4691	27	0
2	C	28	0	25	1	0
2	D	28	0	25	1	0
3	A	53	0	33	4	0
3	B	53	0	33	4	0
4	A	44	0	34	0	0
4	B	42	0	39	0	0
5	A	20	0	9	0	0
5	B	10	0	4	0	0
6	A	9	0	3	0	0
6	B	15	0	5	0	0
7	A	4	0	3	0	0
7	B	4	0	3	1	0
8	A	13	0	18	2	0
8	B	13	0	18	0	0
9	A	10	0	14	0	0
9	B	10	0	14	1	0
10	A	3	0	0	0	0
10	B	2	0	0	0	0
11	A	1	0	0	0	0
12	A	752	0	0	3	0
12	B	763	0	0	8	0
All	All	11203	0	9673	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 47 close contacts within the same asymmetric unit are listed below, sorted by their

clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:244[A]:ARG:HG3	2:C:1:NAG:H83	1.68	0.75
1:A:146[B]:SER:HG	1:A:602:PHE:HZ	1.47	0.61
1:B:50:ILE:HD13	1:B:300[A]:VAL:HG13	1.83	0.61
1:B:132:ASN:HB2	3:B:701:FDA:C5X	2.33	0.59
1:A:132:ASN:HB2	3:A:701:FDA:C5X	2.33	0.57

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	620/595 (104%)	598 (96%)	21 (3%)	1 (0%)	43	22
1	B	620/595 (104%)	593 (96%)	26 (4%)	1 (0%)	43	22
All	All	1240/1190 (104%)	1191 (96%)	47 (4%)	2 (0%)	43	22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	469	THR
1	B	469	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	515/485 (106%)	513 (100%)	2 (0%)	84	71
1	B	514/485 (106%)	514 (100%)	0	100	100
All	All	1029/970 (106%)	1027 (100%)	2 (0%)	88	77

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	628[A]	ARG
1	A	628[B]	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	378	ASN
1	B	46	ASN
1	B	331	ASN
1	B	378	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	2,1	14,14,15	1.70	2 (14%)	17,19,21	2.44	4 (23%)
2	NAG	C	2	2	14,14,15	0.47	0	17,19,21	0.98	1 (5%)
2	NAG	D	1	2,1	14,14,15	1.83	4 (28%)	17,19,21	3.13	4 (23%)
2	NAG	D	2	2	14,14,15	0.63	0	17,19,21	1.51	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	NAG	D	1	2,1	-	1/6/23/26	0/1/1/1
2	NAG	D	2	2	-	0/6/23/26	0/1/1/1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	NAG	O5-C1	4.23	1.50	1.43
2	D	1	NAG	O4-C4	3.59	1.51	1.43
2	C	1	NAG	O4-C4	3.25	1.51	1.43
2	D	1	NAG	O5-C5	3.08	1.49	1.43
2	D	1	NAG	C1-C2	-3.05	1.48	1.52

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	NAG	C1-O5-C5	8.67	123.81	112.19
2	C	1	NAG	O5-C1-C2	-8.00	98.91	111.29
2	D	1	NAG	O5-C1-C2	-5.58	102.65	111.29
2	D	2	NAG	C1-O5-C5	5.27	119.25	112.19
2	D	1	NAG	O3-C3-C4	-4.86	98.92	110.38

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	2	NAG	O5-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	D	1	NAG	O5-C5-C6-O6


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	NAG	1	0
2	D	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 32 ligands modelled in this entry, 6 are monoatomic - leaving 26 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	704	1	14,14,15	0.94	0	17,19,21	1.30	1 (5%)
4	NAG	A	702	1	14,14,15	0.36	0	17,19,21	1.04	0
3	FDA	A	701	-	57,58,58	0.78	0	78,89,89	0.80	3 (3%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	FMT	A	707	-	2,2,2	0.37	0	1,1,1	0.39	0
6	FMT	B	709	-	2,2,2	0.43	0	1,1,1	0.13	0
4	NAG	B	704	1	14,14,15	0.74	0	17,19,21	1.17	1 (5%)
4	NAG	A	703[A]	-	14,14,15	0.89	0	17,19,21	1.20	2 (11%)
5	NPO	A	705[A]	-	10,10,10	0.72	0	11,13,13	0.75	0
3	FDA	B	701	-	57,58,58	0.85	2 (3%)	78,89,89	0.81	2 (2%)
5	NPO	A	706	-	10,10,10	0.66	0	11,13,13	0.83	0
6	FMT	A	708	-	2,2,2	0.28	0	1,1,1	0.04	0
9	PGE	B	713	-	9,9,9	0.28	0	8,8,8	0.32	0
8	PG4	B	712	-	12,12,12	0.21	0	11,11,11	0.18	0
6	FMT	B	707	-	2,2,2	0.31	0	1,1,1	0.04	0
8	PG4	A	711	-	12,12,12	0.30	0	11,11,11	0.22	0
5	NPO	B	705[A]	-	10,10,10	1.02	0	11,13,13	0.88	0
4	NAG	A	703[B]	-	14,14,15	0.88	0	17,19,21	1.25	2 (11%)
9	PGE	A	712	-	9,9,9	0.23	0	8,8,8	0.49	0
6	FMT	B	708	-	2,2,2	0.40	0	1,1,1	0.08	0
4	NAG	B	702	1	14,14,15	0.40	0	17,19,21	1.30	3 (17%)
7	ACT	A	710	-	3,3,3	0.97	0	3,3,3	0.86	0
7	ACT	B	711	-	3,3,3	0.96	0	3,3,3	0.82	0
4	NAG	B	703	1	14,14,15	1.39	2 (14%)	17,19,21	0.83	1 (5%)
6	FMT	A	709	-	2,2,2	0.48	0	1,1,1	0.11	0
6	FMT	B	710	-	2,2,2	0.46	0	1,1,1	0.19	0
6	FMT	B	706	-	2,2,2	0.28	0	1,1,1	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	704	1	-	0/6/23/26	0/1/1/1
8	PG4	B	712	-	-	0/10/10/10	-
9	PGE	A	712	-	-	3/7/7/7	-
9	PGE	B	713	-	-	4/7/7/7	-
4	NAG	A	703[A]	-	-	0/6/23/26	0/1/1/1
4	NAG	A	704	1	-	0/6/23/26	0/1/1/1
4	NAG	B	702	1	-	2/6/23/26	0/1/1/1
4	NAG	A	702	1	-	2/6/23/26	0/1/1/1
8	PG4	A	711	-	-	1/10/10/10	-
5	NPO	A	705[A]	-	-	0/2/4/4	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FDA	B	701	-	-	4/34/50/50	0/6/6/6
4	NAG	B	703	1	-	0/6/23/26	0/1/1/1
5	NPO	A	706	-	-	0/2/4/4	0/1/1/1
3	FDA	A	701	-	-	5/34/50/50	0/6/6/6
5	NPO	B	705[A]	-	-	0/2/4/4	0/1/1/1
4	NAG	A	703[B]	-	-	0/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	703	NAG	O4-C4	2.99	1.50	1.43
4	B	703	NAG	O5-C1	-2.69	1.39	1.43
3	B	701	FDA	O2'-C2'	-2.59	1.37	1.43
3	B	701	FDA	PA-O3P	-2.27	1.57	1.59

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	702	NAG	C4-C3-C2	-3.12	106.45	111.02
4	B	704	NAG	C6-C5-C4	3.08	120.57	113.02
4	A	703[A]	NAG	O5-C1-C2	-2.92	106.77	111.29
4	A	703[B]	NAG	O5-C1-C2	-2.92	106.77	111.29
3	B	701	FDA	O2P-P-O1P	2.77	125.34	112.44

There are no chirality outliers.

5 of 21 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	701	FDA	PA-O3P-P-O5'
3	B	701	FDA	PA-O3P-P-O5'
9	A	712	PGE	C3-C4-O3-C5
4	A	702	NAG	O5-C5-C6-O6
9	B	713	PGE	O2-C3-C4-O3

There are no ring outliers.

5 monomers are involved in 12 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	701	FDA	4	0
3	B	701	FDA	4	0



Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	B	713	PGE	1	0
8	A	711	PG4	2	0
7	B	711	ACT	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	584/595 (98%)	-0.16	6 (1%) 79 82	6, 14, 26, 64	42 (7%)
1	B	586/595 (98%)	-0.17	11 (1%) 66 70	6, 13, 26, 53	41 (6%)
All	All	1170/1190 (98%)	-0.17	17 (1%) 72 75	6, 14, 26, 64	83 (7%)

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	629	LEU	4.1
1	A	628[A]	ARG	3.8
1	B	296	THR	3.4
1	B	97	TRP	3.2
1	B	630	ALA	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	C	2	14/15	0.68	0.17	34,45,50,51	14
2	NAG	D	2	14/15	0.73	0.16	40,47,50,51	14
2	NAG	D	1	14/15	0.89	0.12	15,20,28,36	0
2	NAG	C	1	14/15	0.90	0.11	15,19,25,30	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	B	702	14/15	0.70	0.15	43,55,60,64	0
7	ACT	A	710	4/4	0.74	0.18	54,58,58,60	0
5	NPO	A	706	10/10	0.75	0.16	33,35,39,39	10
4	NAG	A	702	14/15	0.77	0.13	44,50,57,62	0
7	ACT	B	711	4/4	0.80	0.15	43,43,44,47	0
6	FMT	A	707	3/3	0.83	0.16	26,26,29,31	0
9	PGE	A	712	10/10	0.83	0.13	36,39,43,43	0
9	PGE	B	713	10/10	0.83	0.16	41,43,47,49	0
8	PG4	B	712	13/13	0.85	0.13	31,33,38,39	13
6	FMT	B	708	3/3	0.88	0.12	45,45,49,51	0
6	FMT	A	709	3/3	0.88	0.18	31,31,36,37	0
10	MG	B	714	1/1	0.88	0.12	44,44,44,44	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	FMT	B	710	3/3	0.89	0.15	27,27,34,34	0
5	NPO	B	705[A]	10/10	0.90	0.10	17,19,20,20	10
8	PG4	A	711	13/13	0.90	0.11	23,24,36,36	0
6	FMT	B	709	3/3	0.90	0.10	36,36,42,44	0
6	FMT	B	706	3/3	0.91	0.14	28,28,31,32	0
5	NPO	A	705[A]	10/10	0.92	0.09	17,19,21,22	10
6	FMT	A	708	3/3	0.93	0.11	16,16,27,30	0
10	MG	A	713	1/1	0.94	0.09	23,23,23,23	1
6	FMT	B	707	3/3	0.94	0.08	15,15,22,29	0
4	NAG	A	704	14/15	0.95	0.07	16,18,28,33	0
10	MG	B	715	1/1	0.95	0.19	37,37,37,37	0
10	MG	A	714	1/1	0.96	0.24	34,34,34,34	0
4	NAG	B	703	14/15	0.96	0.06	13,15,17,22	0
4	NAG	B	704	14/15	0.96	0.07	15,17,27,36	0
3	FDA	B	701	53/53	0.97	0.05	10,12,14,15	0
4	NAG	A	703[A]	14/15	0.98	0.05	10,12,15,18	2
4	NAG	A	703[B]	14/15	0.98	0.05	10,12,14,14	2
3	FDA	A	701	53/53	0.98	0.05	10,13,15,16	0
10	MG	A	715	1/1	0.99	0.03	12,12,12,12	0
11	NA	A	716	1/1	0.99	0.07	18,18,18,18	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NPO A 706:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NPO B 705 (A):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around NPO A 705 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FDA B 701:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.