



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:16 AM UTC

PDB ID : 6ZEF / pdb_00006zef
Title : Structure of PP1(7-300) bound to Phactr1 (516-580) at pH 5.25
Authors : Mouilleron, S.; Treisman, R.; Fedoryshchak, R.; Lee, R.; Butler, A.M.; Prechova, M.
Deposited on : 2020-06-16
Resolution : 1.94 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	299	6% 93% • •
1	B	299	5% 92% 6% •
2	C	70	34% 76% 16% • 7%
2	D	70	29% 70% 20% • 9%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 11557 atoms, of which 5595 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein phosphatase PP1-alpha catalytic subunit.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	293	Total	C	H	N	O	S	0	3	0
			4645	1508	2299	389	430	19			
1	B	293	Total	C	H	N	O	S	0	1	0
			4621	1500	2285	389	428	19			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	GLY	-	expression tag	UNP P62136
A	3	HIS	-	expression tag	UNP P62136
A	4	MET	-	expression tag	UNP P62136
A	5	GLY	-	expression tag	UNP P62136
A	6	SER	-	expression tag	UNP P62136
B	2	GLY	-	expression tag	UNP P62136
B	3	HIS	-	expression tag	UNP P62136
B	4	MET	-	expression tag	UNP P62136
B	5	GLY	-	expression tag	UNP P62136
B	6	SER	-	expression tag	UNP P62136

- Molecule 2 is a protein called Phosphatase and actin regulator.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	C	65	Total	C	H	N	O	S	0	0	0
			993	314	486	95	97	1			
2	D	64	Total	C	H	N	O	S	0	0	0
			971	307	477	92	94	1			

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	511	GLY	-	expression tag	UNP Q4VY12

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	512	PRO	-	expression tag	UNP Q4VY12
C	513	LEU	-	expression tag	UNP Q4VY12
C	514	GLY	-	expression tag	UNP Q4VY12
C	515	SER	-	expression tag	UNP Q4VY12
D	511	GLY	-	expression tag	UNP Q4VY12
D	512	PRO	-	expression tag	UNP Q4VY12
D	513	LEU	-	expression tag	UNP Q4VY12
D	514	GLY	-	expression tag	UNP Q4VY12
D	515	SER	-	expression tag	UNP Q4VY12

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	B	1	Total	C	H	O	0	0
			10	2	6	2		
3	B	1	Total	C	H	O	0	0
			10	2	6	2		
3	B	1	Total	C	H	O	0	0
			10	2	6	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	C	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		

- Molecule 5 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Mn	0	0
			2	2		
5	B	2	Total	Mn	0	0
			2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	95	Total	O	0	0
			95	95		
6	B	122	Total	O	0	0
			122	122		
6	C	15	Total	O	0	0
			15	15		
6	D	10	Total	O	0	0
			10	10		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serine/threonine-protein phosphatase PP1-alpha catalytic subunit



- Molecule 1: Serine/threonine-protein phosphatase PP1-alpha catalytic subunit



- Molecule 2: Phosphatase and actin regulator



- Molecule 2: Phosphatase and actin regulator



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	47.51Å 57.55Å 89.61Å 78.08° 74.67° 81.66°	Depositor
Resolution (Å)	56.05 – 1.94 56.05 – 1.94	Depositor EDS
% Data completeness (in resolution range)	98.5 (56.05-1.94) 98.5 (56.05-1.94)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.41 (at 1.94Å)	Xtriage
Refinement program	PHENIX 1.18_3845, PHENIX 1.18_3845	Depositor
R, R_{free}	0.181 , 0.211 0.187 , 0.215	Depositor DCC
R_{free} test set	3133 reflections (4.74%)	wwPDB-VP
Wilson B-factor (Å ²)	30.1	Xtriage
Anisotropy	0.600	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 45.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11557	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN, CL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.23	0/2409	0.50	0/3259
1	B	0.22	0/2393	0.50	0/3237
2	C	0.20	0/514	0.42	0/692
2	D	0.21	0/500	0.47	0/674
All	All	0.22	0/5816	0.49	0/7862

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2346	2299	2297	10	0
1	B	2336	2285	2282	14	0
2	C	507	486	486	7	0
2	D	494	477	477	8	1
3	A	16	24	24	2	0
3	B	12	18	18	0	0
3	C	4	6	6	0	0
4	A	1	0	0	0	0
5	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	2	0	0	0	0
6	A	95	0	0	1	0
6	B	122	0	0	1	0
6	C	15	0	0	0	0
6	D	10	0	0	1	0
All	All	5962	5595	5590	33	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:535:ASP:OD1	2:D:537:ARG:HG2	2.03	0.57
2:C:515:SER:OG	2:C:519:LEU:HD11	2.06	0.56
2:C:567:VAL:HG23	2:C:569:GLU:H	1.72	0.55
1:B:127[B]:CYS:SG	1:B:195:VAL:HG21	2.47	0.55
1:B:288:THR:OG1	2:D:521:ARG:NH2	2.40	0.54
2:D:533:ASP:O	6:D:601:HOH:O	2.17	0.50
1:B:74:ARG:NH1	2:D:530:ASP:O	2.41	0.50
1:A:56:GLU:H	3:A:401:EDO:C1	2.25	0.49
2:C:570:LEU:O	2:C:571:SER:OG	2.25	0.48
1:A:56:GLU:H	3:A:401:EDO:H12	1.80	0.47
2:D:515:SER:OG	2:D:519:LEU:HD11	2.14	0.46
1:A:127[B]:CYS:SG	1:A:195:VAL:HG21	2.55	0.46
1:B:188:ARG:NH1	2:C:514:GLY:O	2.49	0.46
1:B:78:TYR:CD2	1:B:296:LEU:CD1	2.99	0.44
1:A:98:LYS:NZ	2:C:537:ARG:O	2.45	0.44
1:A:208:ASP:O	1:A:226:THR:HA	2.17	0.44
1:A:271:ASN:HB3	2:C:536:ARG:HD2	1.99	0.44
2:D:567:VAL:HG23	2:D:569:GLU:H	1.83	0.44
1:B:75:LEU:HD23	1:B:296:LEU:HD13	2.00	0.44
2:C:569:GLU:O	2:C:569:GLU:HG3	2.17	0.44
1:A:142:ARG:NH1	6:A:506:HOH:O	2.50	0.44
1:B:78:TYR:CD2	1:B:296:LEU:HD12	2.54	0.43
1:A:172:CYS:O	1:A:246:ARG:HA	2.19	0.42
1:B:19:VAL:HG13	1:B:70:TYR:OH	2.19	0.42
1:B:19:VAL:O	1:B:19:VAL:CG1	2.67	0.42
1:A:23:ARG:O	1:A:24:PRO:C	2.63	0.42
6:B:607:HOH:O	2:D:568:HIS:CB	2.68	0.42
1:A:60:PRO:HA	1:A:285:VAL:O	2.19	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:208:ASP:O	1:B:226:THR:HA	2.20	0.41
1:B:290:MET:HE2	2:D:521:ARG:NH1	2.36	0.41
1:B:45:ILE:HG21	1:B:111:LYS:HE3	2.04	0.40
1:B:161:ILE:HD12	1:B:201:LEU:HD11	2.03	0.40
1:B:286:ASP:C	1:B:286:ASP:OD1	2.65	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:524:ASP:O	2:D:575:THR:OG1[1_655]	1.95	0.25

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	294/299 (98%)	280 (95%)	14 (5%)	0	100	100
1	B	292/299 (98%)	281 (96%)	11 (4%)	0	100	100
2	C	63/70 (90%)	60 (95%)	3 (5%)	0	100	100
2	D	62/70 (89%)	59 (95%)	3 (5%)	0	100	100
All	All	711/738 (96%)	680 (96%)	31 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	255/262 (97%)	254 (100%)	1 (0%)	84	84
1	B	253/262 (97%)	251 (99%)	2 (1%)	73	71
2	C	50/61 (82%)	46 (92%)	4 (8%)	11	2
2	D	48/61 (79%)	44 (92%)	4 (8%)	10	2
All	All	606/646 (94%)	595 (98%)	11 (2%)	51	43

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	246	ARG
1	B	246	ARG
1	B	296	LEU
2	C	521	ARG
2	C	564	GLU
2	C	570	LEU
2	C	572	ARG
2	D	564	GLU
2	D	572	ARG
2	D	574	LEU
2	D	575	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	ASN
1	A	198	GLN
1	A	262	GLN
1	B	99	GLN
1	B	181	GLN
1	B	262	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 5 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	B	402	-	3,3,3	0.40	0	2,2,2	0.46	0
3	EDO	A	403	-	3,3,3	0.44	0	2,2,2	0.36	0
3	EDO	A	401	-	3,3,3	0.47	0	2,2,2	0.28	0
3	EDO	B	403	-	3,3,3	0.46	0	2,2,2	0.37	0
3	EDO	C	601	-	3,3,3	0.46	0	2,2,2	0.21	0
3	EDO	A	404	-	3,3,3	0.43	0	2,2,2	0.39	0
3	EDO	B	401	-	3,3,3	0.45	0	2,2,2	0.31	0
3	EDO	A	402	-	3,3,3	0.46	0	2,2,2	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	402	-	-	0/1/1/1	-
3	EDO	A	403	-	-	1/1/1/1	-
3	EDO	A	401	-	-	0/1/1/1	-
3	EDO	B	403	-	-	1/1/1/1	-
3	EDO	C	601	-	-	0/1/1/1	-
3	EDO	A	404	-	-	1/1/1/1	-
3	EDO	B	401	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	402	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	403	EDO	O1-C1-C2-O2
3	B	403	EDO	O1-C1-C2-O2
3	A	404	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	401	EDO	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	293/299 (97%)	0.41	17 (5%)	29 32	17, 38, 63, 94	3 (1%)
1	B	293/299 (97%)	0.42	14 (4%)	35 40	16, 39, 66, 102	1 (0%)
2	C	65/70 (92%)	2.05	24 (36%)	1 0	45, 59, 94, 112	0
2	D	64/70 (91%)	2.04	20 (31%)	1 1	44, 59, 88, 113	0
All	All	715/738 (96%)	0.71	75 (10%)	11 12	16, 42, 75, 113	4 (0%)

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	568	HIS	8.8
2	D	570	LEU	8.3
2	D	512	PRO	6.5
2	C	576	ARG	6.0
2	C	571	SER	5.9
2	D	571	SER	5.9
2	D	569	GLU	5.7
2	D	568	HIS	5.6
1	A	23	ARG	5.5
2	C	570	LEU	5.5
2	D	575	THR	5.5
1	A	299	ALA	5.2
2	C	575	THR	5.1
2	D	574	LEU	5.1
1	A	24	PRO	5.0
2	C	512	PRO	4.9
2	D	566	GLU	4.9
1	B	23	ARG	4.8
1	B	7	LEU	4.5
2	C	573	HIS	4.3
2	D	573	HIS	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	569	GLU	4.0
1	B	25	GLY	4.0
1	B	299	ALA	4.0
2	C	567	VAL	4.0
2	C	566	GLU	3.9
1	A	25	GLY	3.9
2	D	567	VAL	3.9
1	A	7	LEU	3.8
2	C	574	LEU	3.6
2	D	563	THR	3.6
2	D	526	VAL	3.5
2	C	526	VAL	3.4
1	A	21	GLY	3.3
1	B	24	PRO	3.3
2	C	525	TYR	3.3
1	B	78	TYR	3.2
1	B	21	GLY	3.0
1	B	20	GLN	3.0
2	C	572	ARG	2.9
2	C	563	THR	2.9
2	D	525	TYR	2.9
2	D	572	ARG	2.9
1	B	19	VAL	2.8
2	D	514	GLY	2.8
1	B	297	LYS	2.8
1	A	8	ASN	2.7
2	C	514	GLY	2.7
2	D	532	GLN	2.7
1	A	78	TYR	2.6
1	A	179	ASP	2.6
1	A	20	GLN	2.6
2	C	513	LEU	2.6
2	C	531	ALA	2.6
2	C	558	ASN	2.6
2	C	538	ALA	2.5
2	C	517	LYS	2.5
2	C	548	ALA	2.4
1	A	19	VAL	2.4
1	A	70	TYR	2.4
1	A	14	GLY	2.4
1	A	297	LYS	2.3
2	C	532	GLN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	60(A)	LEU	2.3
1	B	296	LEU	2.3
1	A	277	ASP	2.3
2	D	520	ILE	2.2
2	D	517	LYS	2.2
2	D	513	LEU	2.2
1	B	277	ASP	2.1
2	C	528	VAL	2.1
2	D	564	GLU	2.0
1	B	294	GLN	2.0
1	A	213	VAL	2.0
1	B	8	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MN	B	405	1/1	0.67	0.22	87,87,87,87	0
5	MN	A	407	1/1	0.80	0.15	93,93,93,93	0
3	EDO	A	402	4/4	0.83	0.18	33,47,56,56	0
5	MN	B	404	1/1	0.84	0.25	65,65,65,65	0
5	MN	A	406	1/1	0.89	0.26	64,64,64,64	0
3	EDO	A	403	4/4	0.89	0.14	37,46,54,65	0
3	EDO	A	401	4/4	0.90	0.13	31,42,50,60	0
3	EDO	B	402	4/4	0.90	0.20	41,49,55,55	0
3	EDO	A	404	4/4	0.92	0.12	44,53,57,60	0
3	EDO	C	601	4/4	0.92	0.12	45,55,60,63	0
3	EDO	B	401	4/4	0.92	0.11	35,45,47,54	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	EDO	B	403	4/4	0.93	0.10	32,39,47,51	0
4	CL	A	405	1/1	0.96	0.11	38,38,38,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.