



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 28, 2026 – 10:51 PM UTC

PDB ID : 3ZFX / pdb_00003zfx
Title : Crystal structure of EphB1
Authors : Debreczeni, J.E.; Overman, R.; Truman, C.; McAlister, M.; Attwood, T.K.
Deposited on : 2012-12-12
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	% 75% 14% • 10%
1	B	298	79% 7% 15%
1	C	298	% 75% 7% • 17%
1	D	298	79% 9% • 11%
1	E	298	% 77% 11% • 12%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	298	% 78%7%15%
1	G	298	75%10%15%
1	H	298	77%11%• 11%
1	I	298	2% 79%9%• 12%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 18158 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EPHRIN TYPE-B RECEPTOR 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	S	0	0	0
			2075	1330	350	380	15			
1	B	254	Total	C	N	O	S	0	0	0
			1961	1257	330	359	15			
1	C	248	Total	C	N	O	S	0	0	0
			1909	1223	318	353	15			
1	D	265	Total	C	N	O	S	0	0	0
			2052	1314	345	378	15			
1	E	262	Total	C	N	O	S	0	0	0
			2021	1294	339	373	15			
1	F	254	Total	C	N	O	S	0	0	0
			1946	1248	317	366	15			
1	G	253	Total	C	N	O	S	0	0	0
			1974	1261	332	366	15			
1	H	266	Total	C	N	O	S	0	0	0
			2051	1316	344	376	15			
1	I	263	Total	C	N	O	S	0	0	0
			2030	1300	343	372	15			

There are 27 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	599	GLY	-	expression tag	UNP P54762
A	600	SER	-	expression tag	UNP P54762
A	601	SER	-	expression tag	UNP P54762
B	599	GLY	-	expression tag	UNP P54762
B	600	SER	-	expression tag	UNP P54762
B	601	SER	-	expression tag	UNP P54762
C	599	GLY	-	expression tag	UNP P54762
C	600	SER	-	expression tag	UNP P54762
C	601	SER	-	expression tag	UNP P54762
D	599	GLY	-	expression tag	UNP P54762
D	600	SER	-	expression tag	UNP P54762

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	601	SER	-	expression tag	UNP P54762
E	599	GLY	-	expression tag	UNP P54762
E	600	SER	-	expression tag	UNP P54762
E	601	SER	-	expression tag	UNP P54762
F	599	GLY	-	expression tag	UNP P54762
F	600	SER	-	expression tag	UNP P54762
F	601	SER	-	expression tag	UNP P54762
G	599	GLY	-	expression tag	UNP P54762
G	600	SER	-	expression tag	UNP P54762
G	601	SER	-	expression tag	UNP P54762
H	599	GLY	-	expression tag	UNP P54762
H	600	SER	-	expression tag	UNP P54762
H	601	SER	-	expression tag	UNP P54762
I	599	GLY	-	expression tag	UNP P54762
I	600	SER	-	expression tag	UNP P54762
I	601	SER	-	expression tag	UNP P54762

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	52	Total 52	O 52	0	0
3	B	7	Total 7	O 7	0	0
3	C	1	Total 1	O 1	0	0
3	D	39	Total 39	O 39	0	0
3	E	3	Total 3	O 3	0	0
3	F	4	Total 4	O 4	0	0
3	G	5	Total 5	O 5	0	0
3	H	17	Total 17	O 17	0	0
3	I	6	Total 6	O 6	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: EPHRIN TYPE-B RECEPTOR 1




• Molecule 1: EPHRIN TYPE-B RECEPTOR 1



• Molecule 1: EPHRIN TYPE-B RECEPTOR 1





• Molecule 1: EPHRIN TYPE-B RECEPTOR 1









• Molecule 1: EPHRIN TYPE-B RECEPTOR 1





• Molecule 1: EPHRIN TYPE-B RECEPTOR 1




• Molecule 1: EPHRIN TYPE-B RECEPTOR 1




• Molecule 1: EPHRIN TYPE-B RECEPTOR 1



• Molecule 1: EPHRIN TYPE-B RECEPTOR 1





4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	195.98Å 195.98Å 60.24Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	15.00 – 2.50 15.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.8 (15.00-2.50) 99.5 (15.00-2.50)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.93 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.188 , 0.217 0.203 , 0.226	Depositor DCC
R_{free} test set	4677 reflections (5.22%)	wwPDB-VP
Wilson B-factor (Å ²)	32.7	Xtriage
Anisotropy	0.149	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 38.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.000 for -h,-k,l 0.000 for h,-h-k,-l 0.097 for -k,-h,-l	Xtriage
Reported twinning fraction	0.031 for H, K, L 0.030 for -K, -H, -L 0.432 for -h,-k,l 0.507 for K, H, -L	Depositor
Outliers	0 of 89479 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	18158	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.38 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.0458e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	2/2116 (0.1%)	1.02	8/2863 (0.3%)
1	B	0.74	4/1998 (0.2%)	0.84	0/2705
1	C	0.60	0/1946	0.90	3/2638 (0.1%)
1	D	0.82	0/2090	0.95	3/2828 (0.1%)
1	E	0.75	5/2059 (0.2%)	0.84	2/2790 (0.1%)
1	F	0.61	0/1983	0.81	1/2692 (0.0%)
1	G	0.80	8/2011 (0.4%)	0.89	1/2723 (0.0%)
1	H	0.90	5/2092 (0.2%)	0.96	4/2833 (0.1%)
1	I	0.65	3/2070 (0.1%)	0.84	5/2804 (0.2%)
All	All	0.76	27/18365 (0.1%)	0.90	27/24876 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	748	ARG	CZ-NH2	-10.84	1.19	1.33
1	B	734	TYR	CE1-CZ	-7.91	1.19	1.38
1	B	734	TYR	CG-CD2	-7.35	1.24	1.39
1	B	734	TYR	CG-CD1	-7.05	1.24	1.39
1	H	763	PHE	CG-CD2	-6.91	1.24	1.38

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	710	ARG	NE-CZ-NH2	-12.49	107.96	119.20
1	C	710	ARG	CD-NE-CZ	12.14	141.40	124.40
1	A	748	ARG	NH1-CZ-NH2	-9.66	106.75	119.30
1	A	748	ARG	NE-CZ-NH1	8.45	129.95	121.50
1	D	789	ARG	NE-CZ-NH2	-7.64	112.32	119.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	710	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2075	0	2028	24	1
1	B	1961	0	1896	8	1
1	C	1909	0	1831	12	0
1	D	2052	0	1999	16	2
1	E	2021	0	1953	17	1
1	F	1946	0	1850	10	1
1	G	1974	0	1910	17	0
1	H	2051	0	1993	21	0
1	I	2030	0	1969	21	0
2	A	5	0	0	0	0
3	A	52	0	0	4	0
3	B	7	0	0	0	0
3	C	1	0	0	0	0
3	D	39	0	0	0	0
3	E	3	0	0	0	0
3	F	4	0	0	0	0
3	G	5	0	0	0	0
3	H	17	0	0	0	0
3	I	6	0	0	0	0
All	All	18158	0	17429	140	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 140 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:741:VAL:HG23	1:I:769:LEU:HD11	1.49	0.94
1:D:651:LYS:NZ	1:D:668:GLU:OE1	2.09	0.84
1:G:741:VAL:HG12	1:G:743:ARG:HG2	1.60	0.83
1:A:816:GLU:OE2	3:A:2026:HOH:O	2.03	0.75
1:I:741:VAL:CG2	1:I:769:LEU:HD11	2.17	0.75

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:734:TYR:OH	1:B:884:ASN:ND2[1_556]	1.93	0.27
1:D:734:TYR:OH	1:E:884:ASN:OD1[3_554]	2.09	0.11
1:D:865:ARG:NH1	1:F:768:TYR:OH[1_554]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	259/298 (87%)	248 (96%)	11 (4%)	0	100	100
1	B	244/298 (82%)	237 (97%)	7 (3%)	0	100	100
1	C	238/298 (80%)	230 (97%)	8 (3%)	0	100	100
1	D	255/298 (86%)	245 (96%)	10 (4%)	0	100	100
1	E	252/298 (85%)	240 (95%)	11 (4%)	1 (0%)	30	49
1	F	244/298 (82%)	237 (97%)	7 (3%)	0	100	100
1	G	243/298 (82%)	237 (98%)	6 (2%)	0	100	100
1	H	258/298 (87%)	251 (97%)	7 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	I	255/298 (86%)	246 (96%)	9 (4%)	0	100	100
All	All	2248/2682 (84%)	2171 (97%)	76 (3%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	788	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	214/256 (84%)	209 (98%)	5 (2%)	44	72
1	B	200/256 (78%)	195 (98%)	5 (2%)	42	69
1	C	195/256 (76%)	191 (98%)	4 (2%)	47	74
1	D	211/256 (82%)	206 (98%)	5 (2%)	43	70
1	E	207/256 (81%)	200 (97%)	7 (3%)	32	60
1	F	198/256 (77%)	195 (98%)	3 (2%)	57	80
1	G	205/256 (80%)	198 (97%)	7 (3%)	32	60
1	H	210/256 (82%)	206 (98%)	4 (2%)	50	76
1	I	209/256 (82%)	205 (98%)	4 (2%)	50	76
All	All	1849/2304 (80%)	1805 (98%)	44 (2%)	43	70

5 of 44 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	893	THR
1	G	890	THR
1	G	617	SER
1	G	739	ASN
1	H	668	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	831	GLN
1	I	661	GLN
1	G	739	ASN
1	E	839	GLN
1	G	831	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	1890	-	4,4,4	0.78	0	6,6,6	0.29	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	267/298 (89%)	0.04	2 (0%) 84 81	17, 26, 40, 44	0
1	B	254/298 (85%)	0.23	0 100 100	28, 35, 46, 49	0
1	C	248/298 (83%)	0.29	3 (1%) 76 73	35, 41, 48, 52	0
1	D	265/298 (88%)	-0.19	1 (0%) 88 86	14, 23, 33, 42	0
1	E	262/298 (87%)	0.18	2 (0%) 82 80	28, 36, 47, 53	0
1	F	254/298 (85%)	0.37	2 (0%) 82 80	34, 40, 52, 58	0
1	G	253/298 (84%)	0.12	0 100 100	26, 34, 47, 52	0
1	H	266/298 (89%)	-0.21	0 100 100	14, 24, 34, 42	0
1	I	263/298 (88%)	0.50	5 (1%) 66 62	37, 43, 51, 56	0
All	All	2332/2682 (86%)	0.14	15 (0%) 85 83	14, 35, 48, 58	0

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	627	ALA	3.1
1	E	655	ALA	2.8
1	C	644	ARG	2.8
1	D	893	THR	2.7
1	C	867	SER	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	A	1890	5/5	0.96	0.11	25,28,32,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.