



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 06:19 PM UTC

PDB ID : 4ZFF / pdb_00004zff
Title : Dual-acting Fab 5A12 in complex with VEGF
Authors : Harris, S.F.; Wu, P.
Deposited on : 2015-04-21
Resolution : 2.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	228	 15% 70% 18% 10%
1	H	228	 73% 17% 7%
2	B	215	 20% 80% 15% 2%
2	L	215	 80% 17% 2%
3	C	99	 2% 81% 14% 2%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	99	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into two segments: a green segment on the left labeled '83%' and a yellow segment on the right labeled '16%'. A small red square is at the beginning of the bar, and a small black dot is at the end. A '%' symbol is positioned above the start of the bar.</p>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8374 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fragment antigen binding (Fab) 5A12 Heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	206	Total	C	N	O	S	0	1	0
			1554	997	252	299	6			
1	H	211	Total	C	N	O	S	0	0	0
			1589	1019	259	305	6			

- Molecule 2 is a protein called Fragment antigen binding (Fab) 5A12 Light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	210	Total	C	N	O	S	0	0	0
			1600	1006	266	323	5			
2	L	213	Total	C	N	O	S	0	0	0
			1622	1020	270	327	5			

- Molecule 3 is a protein called Vascular endothelial growth factor A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	96	Total	C	N	O	S	0	0	0
			775	486	130	146	13			
3	D	99	Total	C	N	O	S	0	0	0
			800	501	137	149	13			


- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	L	1	Total	O S	0	0
			5	4 1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	18	Total	O	0	0
			18	18		
5	B	29	Total	O	0	0
			29	29		
5	C	43	Total	O	0	0
			43	43		
5	D	49	Total	O	0	0
			49	49		
5	H	138	Total	O	0	0
			138	138		
5	L	152	Total	O	0	0
			152	152		

Chain L:  80% 17% ..




• Molecule 3: Vascular endothelial growth factor A

Chain C:  2% 81% 14% ...



• Molecule 3: Vascular endothelial growth factor A

Chain D:  83% 16% .



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	87.41Å 313.89Å 51.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.95 – 2.75 26.95 – 2.75	Depositor EDS
% Data completeness (in resolution range)	99.9 (26.95-2.75) 99.8 (26.95-2.75)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.53 (at 2.77Å)	Xtrriage
Refinement program	BUSTER 2.11.5	Depositor
R, R_{free}	0.197 , 0.238 0.202 , 0.248	Depositor DCC
R_{free} test set	1934 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	51.1	Xtrriage
Anisotropy	0.258	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 81.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8374	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.70	0/1595	1.12	5/2178 (0.2%)
1	H	0.84	0/1632	1.16	6/2230 (0.3%)
2	B	0.73	0/1634	1.15	0/2215
2	L	0.86	2/1657 (0.1%)	1.18	4/2248 (0.2%)
3	C	0.82	0/793	1.20	5/1069 (0.5%)
3	D	0.85	0/820	1.24	7/1106 (0.6%)
All	All	0.80	2/8131 (0.0%)	1.17	27/11046 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	L	4	MET	SD-CE	-6.33	1.63	1.79
2	L	140	TYR	CA-C	5.20	1.58	1.52

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	61	CYS	N-CA-C	-8.76	95.48	109.59
3	D	20	VAL	N-CA-CB	6.51	118.17	110.55
3	D	60	CYS	CA-C-N	6.38	132.19	122.93
3	D	60	CYS	C-N-CA	6.38	132.19	122.93
1	H	67	PHE	CA-CB-CG	6.28	120.08	113.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1554	0	1516	15	0
1	H	1589	0	1553	17	0
2	B	1600	0	1567	19	0
2	L	1622	0	1590	16	0
3	C	775	0	742	13	0
3	D	800	0	758	11	0
4	L	5	0	0	0	0
5	A	18	0	0	0	0
5	B	29	0	0	0	0
5	C	43	0	0	0	0
5	D	49	0	0	1	0
5	H	138	0	0	2	0
5	L	152	0	0	2	0
All	All	8374	0	7726	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 79 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:60:CYS:HG	3:D:51:CYS:HG	0.88	0.85
3:C:89:GLN:HE22	1:H:97:PHE:H	1.28	0.78
1:H:195:ILE:HG22	1:H:210:LYS:HA	1.68	0.74
3:C:50:SER:H	3:D:62:ASN:ND2	1.90	0.69
2:B:148:TRP:CE3	2:B:179:LEU:HD11	2.30	0.66

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	201/228 (88%)	194 (96%)	6 (3%)	1 (0%)	24	43
1	H	207/228 (91%)	201 (97%)	5 (2%)	1 (0%)	24	43
2	B	206/215 (96%)	195 (95%)	8 (4%)	3 (2%)	8	15
2	L	211/215 (98%)	204 (97%)	6 (3%)	1 (0%)	24	43
3	C	94/99 (95%)	93 (99%)	0	1 (1%)	11	22
3	D	97/99 (98%)	92 (95%)	4 (4%)	1 (1%)	12	24
All	All	1016/1084 (94%)	979 (96%)	29 (3%)	8 (1%)	16	30

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	D	26	CYS
1	A	144	ASP
2	B	211	ARG
2	L	211	ARG
2	B	138	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/188 (90%)	149 (88%)	20 (12%)	5	9
1	H	173/188 (92%)	157 (91%)	16 (9%)	8	17
2	B	183/187 (98%)	169 (92%)	14 (8%)	12	22
2	L	185/187 (99%)	172 (93%)	13 (7%)	14	26
3	C	90/94 (96%)	86 (96%)	4 (4%)	25	47
3	D	92/94 (98%)	88 (96%)	4 (4%)	26	47
All	All	892/938 (95%)	821 (92%)	71 (8%)	11	20

5 of 71 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	195	ILE
2	L	48	ILE
2	L	143	GLU
2	B	122	ASP
2	B	100	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
3	D	98	GLN
1	H	164	HIS
2	L	155	GLN
1	H	199	ASN
1	H	13	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	L	301	-	4,4,4	0.41	0	6,6,6	0.25	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	206/228 (90%)	1.12	34 (16%) 4 4	29, 111, 146, 158	1 (0%)
1	H	211/228 (92%)	-0.33	0 100 100	24, 42, 80, 101	0
2	B	210/215 (97%)	1.27	42 (20%) 3 2	56, 101, 143, 150	0
2	L	213/215 (99%)	-0.45	1 (0%) 87 87	24, 38, 68, 85	0
3	C	96/99 (96%)	-0.11	2 (2%) 63 61	30, 49, 89, 100	0
3	D	99/99 (100%)	-0.29	1 (1%) 79 79	31, 47, 79, 105	0
All	All	1035/1084 (95%)	0.28	80 (7%) 19 19	24, 59, 140, 158	1 (0%)

The worst 5 of 80 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	56[A]	TYR	5.3
1	A	158	ALA	4.4
1	A	119	PRO	4.1
2	B	148	TRP	4.0
2	B	196	VAL	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SO4	L	301	5/5	0.97	0.06	47,49,51,52	0

6.5 Other polymers [i](#)

There are no such residues in this entry.