



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 5, 2026 – 12:07 PM UTC

PDB ID : 1ZGP / pdb_00001zgp
Title : Crystal Structure of the Discosoma Red Fluorescent Protein (DsRed) Variant K70M
Authors : Tubbs, J.L.; Tainer, J.A.; Getzoff, E.D.
Deposited on : 2005-04-21
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	223	 9% 87% 11% .
1	B	223	 7% 87% 9% ..
1	C	223	 9% 87% 9% ..
1	D	223	 9% 86% 11% ..

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7491 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Red fluorescent protein drFP583.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	218	Total	C	N	O	S	0	0	0
			1790	1157	297	328	8			
1	B	218	Total	C	N	O	S	0	0	0
			1790	1157	297	328	8			
1	C	218	Total	C	N	O	S	0	0	0
			1790	1157	297	328	8			
1	D	218	Total	C	N	O	S	0	0	0
			1790	1157	297	328	8			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	66	CRQ	GLN	chromophore	UNP Q9U6Y8
A	66	CRQ	TYR	chromophore	UNP Q9U6Y8
A	66	CRQ	GLY	chromophore	UNP Q9U6Y8
A	70	MET	LYS	engineered mutation	UNP Q9U6Y8
B	66	CRQ	GLN	chromophore	UNP Q9U6Y8
B	66	CRQ	TYR	chromophore	UNP Q9U6Y8
B	66	CRQ	GLY	chromophore	UNP Q9U6Y8
B	70	MET	LYS	engineered mutation	UNP Q9U6Y8
C	66	CRQ	GLN	chromophore	UNP Q9U6Y8
C	66	CRQ	TYR	chromophore	UNP Q9U6Y8
C	66	CRQ	GLY	chromophore	UNP Q9U6Y8
C	70	MET	LYS	engineered mutation	UNP Q9U6Y8
D	66	CRQ	GLN	chromophore	UNP Q9U6Y8
D	66	CRQ	TYR	chromophore	UNP Q9U6Y8
D	66	CRQ	GLY	chromophore	UNP Q9U6Y8
D	70	MET	LYS	engineered mutation	UNP Q9U6Y8


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	76	Total 76	O 76	0	0
2	B	80	Total 80	O 80	0	0
2	C	83	Total 83	O 83	0	0
2	D	92	Total 92	O 92	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Red fluorescent protein drFP583




- Molecule 1: Red fluorescent protein drFP583




- Molecule 1: Red fluorescent protein drFP583



- Molecule 1: Red fluorescent protein drFP583





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	57.28Å 57.23Å 71.54Å 89.74° 100.25° 92.86°	Depositor
Resolution (Å)	32.51 – 1.90 32.51 – 1.90	Depositor EDS
% Data completeness (in resolution range)	81.8 (32.51-1.90) 81.8 (32.51-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.78 (at 1.89Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.226 , 0.263 0.229 , 0.262	Depositor DCC
R_{free} test set	2903 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	21.3	Xtriage
Anisotropy	0.772	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 30.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.049 for -h,k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7491	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CRQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	0/1815	1.04	9/2444 (0.4%)
1	B	0.67	1/1815 (0.1%)	1.06	11/2444 (0.5%)
1	C	0.67	0/1815	1.05	9/2444 (0.4%)
1	D	0.67	0/1815	1.06	11/2444 (0.5%)
All	All	0.67	1/7260 (0.0%)	1.05	40/9776 (0.4%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	70	MET	SD-CE	-6.12	1.64	1.79

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	69	SER	CA-C-N	8.99	132.70	120.38
1	D	69	SER	C-N-CA	8.99	132.70	120.38
1	A	69	SER	CA-C-N	8.91	132.59	120.38
1	A	69	SER	C-N-CA	8.91	132.59	120.38
1	C	69	SER	CA-C-N	8.34	132.56	120.38
1	C	69	SER	C-N-CA	8.34	132.56	120.38
1	B	69	SER	CA-C-N	7.09	130.72	120.38
1	B	69	SER	C-N-CA	7.09	130.72	120.38
1	B	36	ARG	CA-C-N	6.20	125.69	119.24
1	B	36	ARG	C-N-CA	6.20	125.69	119.24
1	D	36	ARG	CA-C-N	6.14	125.63	119.24
1	D	36	ARG	C-N-CA	6.14	125.63	119.24
1	A	206	GLU	N-CA-C	6.05	118.37	111.11
1	A	144	GLU	N-CA-C	-5.98	102.47	110.55
1	B	144	GLU	N-CA-C	-5.88	102.61	110.55
1	D	7	VAL	N-CA-C	5.84	116.62	110.72

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	144	GLU	N-CA-C	-5.78	102.74	110.55
1	B	20	GLY	N-CA-C	5.68	119.39	111.10
1	B	146	SER	N-CA-C	5.67	118.03	109.41
1	B	206	GLU	N-CA-C	5.67	117.91	111.11
1	D	146	SER	N-CA-C	5.60	117.92	109.41
1	C	36	ARG	CA-C-N	5.55	125.64	119.32
1	C	36	ARG	C-N-CA	5.55	125.64	119.32
1	A	70	MET	N-CA-C	5.54	118.03	111.33
1	A	174	LEU	N-CA-C	5.48	118.48	109.76
1	A	77	ALA	N-CA-C	5.38	117.84	111.33
1	C	146	SER	N-CA-C	5.31	117.49	109.41
1	D	174	LEU	N-CA-C	5.31	118.58	110.20
1	D	20	GLY	N-CA-C	5.29	119.00	111.12
1	D	144	GLU	N-CA-C	-5.26	102.73	110.52
1	B	201	ILE	N-CA-C	-5.25	100.19	108.23
1	B	168	LYS	N-CA-C	5.23	118.76	112.38
1	D	206	GLU	N-CA-C	5.16	117.30	111.11
1	D	70	MET	N-CA-C	5.15	117.56	111.33
1	C	174	LEU	N-CA-C	5.11	118.28	110.20
1	C	20	GLY	N-CA-C	5.08	118.70	111.12
1	C	59	ASP	N-CA-C	5.06	118.23	111.75
1	A	146	SER	N-CA-C	5.05	117.09	109.41
1	B	59	ASP	N-CA-C	5.02	118.17	111.75
1	A	100	GLU	N-CA-C	5.01	118.49	112.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1790	0	1738	5	0
1	B	1790	0	1738	8	0
1	C	1790	0	1738	6	0
1	D	1790	0	1738	7	0
2	A	76	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	80	0	0	1	0
2	C	83	0	0	1	0
2	D	92	0	0	1	0
All	All	7491	0	6952	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:203:SER:HB3	1:D:212:GLU:HB2	1.82	0.61
1:C:203:SER:HB3	1:C:212:GLU:HB2	1.83	0.60
1:B:203:SER:HB3	1:B:212:GLU:HB2	1.86	0.56
1:B:83:LYS:HE2	2:B:304:HOH:O	2.04	0.56
1:B:94:GLU:HG3	1:B:180:ILE:HB	1.90	0.53
1:B:66:CRQ:HE1	1:B:199:LEU:HB2	1.91	0.53
1:A:203:SER:HB3	1:A:212:GLU:HB2	1.91	0.52
1:D:66:CRQ:HE1	1:D:199:LEU:HB2	1.93	0.51
1:C:66:CRQ:HE1	1:C:199:LEU:HB2	1.93	0.51
1:B:66:CRQ:O2	1:B:70:MET:HE1	2.13	0.49
1:C:70:MET:O	1:C:73:VAL:HG12	2.15	0.47
1:A:91:PHE:HZ	1:A:181:TYR:HB3	1.80	0.45
1:C:151:TYR:HE2	1:C:160:GLU:OE1	1.98	0.45
1:D:217:THR:HG22	2:D:286:HOH:O	2.16	0.45
1:D:59:ASP:HB3	1:D:165:LEU:HD21	1.99	0.45
1:B:83:LYS:H	1:B:83:LYS:NZ	2.14	0.45
1:A:66:CRQ:HE1	1:A:199:LEU:HB2	1.97	0.44
1:B:59:ASP:HB3	1:B:165:LEU:HD21	1.99	0.44
1:B:70:MET:HG3	1:B:91:PHE:CE2	2.52	0.44
1:A:59:ASP:HB3	1:A:165:LEU:HD21	1.99	0.43
1:C:59:ASP:HB3	1:C:165:LEU:HD21	2.00	0.43
1:C:70:MET:H	1:C:70:MET:HG2	1.55	0.43
1:A:225:LEU:HD11	2:C:251:HOH:O	2.19	0.43
1:D:54:LEU:HA	1:D:55:PRO:HD3	1.87	0.42
1:D:40:GLY:HA2	1:D:72:TYR:O	2.21	0.41
1:D:109:GLN:HG3	1:D:122:VAL:HG22	2.01	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/223 (96%)	209 (98%)	3 (1%)	1 (0%)	24	16
1	B	213/223 (96%)	208 (98%)	3 (1%)	2 (1%)	14	6
1	C	213/223 (96%)	209 (98%)	3 (1%)	1 (0%)	24	16
1	D	213/223 (96%)	209 (98%)	3 (1%)	1 (0%)	24	16
All	All	852/892 (96%)	835 (98%)	12 (1%)	5 (1%)	21	13

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	186	PRO
1	D	186	PRO
1	A	186	PRO
1	B	170	GLY
1	C	186	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/196 (97%)	184 (96%)	7 (4%)	30	22
1	B	191/196 (97%)	185 (97%)	6 (3%)	35	29
1	C	191/196 (97%)	185 (97%)	6 (3%)	35	29
1	D	191/196 (97%)	186 (97%)	5 (3%)	40	35
All	All	764/784 (97%)	740 (97%)	24 (3%)	35	29

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	ASN
1	A	32	GLU
1	A	42	ASN
1	A	71	VAL
1	A	160	GLU
1	A	161	ILE
1	A	169	ASP
1	B	83	LYS
1	B	113	LEU
1	B	160	GLU
1	B	161	ILE
1	B	187	VAL
1	B	206	GLU
1	C	7	VAL
1	C	16	VAL
1	C	42	ASN
1	C	54	LEU
1	C	71	VAL
1	C	161	ILE
1	D	70	MET
1	D	71	VAL
1	D	160	GLU
1	D	161	ILE
1	D	200	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	ASN
1	A	98	ASN
1	B	204	HIS
1	C	98	ASN
1	C	114	GLN
1	D	98	ASN
1	D	188	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CRQ	D	66	1	25,25,26	3.09	11 (44%)	28,34,36	1.77	6 (21%)
1	CRQ	B	66	1	25,25,26	3.16	11 (44%)	28,34,36	1.78	4 (14%)
1	CRQ	C	66	1	25,25,26	3.04	11 (44%)	28,34,36	1.83	6 (21%)
1	CRQ	A	66	1	25,25,26	3.10	11 (44%)	28,34,36	1.83	6 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRQ	D	66	1	-	2/10/32/33	0/2/2/2
1	CRQ	B	66	1	-	1/10/32/33	0/2/2/2
1	CRQ	C	66	1	-	1/10/32/33	0/2/2/2
1	CRQ	A	66	1	-	1/10/32/33	0/2/2/2

All (44) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66	CRQ	CB2-CA2	6.95	1.41	1.35
1	B	66	CRQ	CB2-CA2	6.77	1.41	1.35
1	D	66	CRQ	CB2-CA2	6.42	1.41	1.35
1	B	66	CRQ	CE2-CZ	6.04	1.50	1.39
1	A	66	CRQ	CE2-CZ	5.88	1.50	1.39
1	D	66	CRQ	CE2-CZ	5.88	1.50	1.39
1	C	66	CRQ	CB2-CA2	5.85	1.40	1.35
1	C	66	CRQ	CE2-CZ	5.71	1.49	1.39
1	B	66	CRQ	CE1-CZ	5.46	1.49	1.39
1	B	66	CRQ	OH-CZ	-5.44	1.24	1.37
1	D	66	CRQ	CE1-CZ	5.41	1.49	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	66	CRQ	OH-CZ	-5.36	1.25	1.37
1	A	66	CRQ	OH-CZ	-5.33	1.25	1.37
1	D	66	CRQ	OH-CZ	-5.33	1.25	1.37
1	C	66	CRQ	CE1-CZ	5.32	1.49	1.39
1	A	66	CRQ	CE1-CZ	5.29	1.48	1.39
1	A	66	CRQ	CD1-CG2	5.22	1.50	1.39
1	D	66	CRQ	CD1-CG2	5.21	1.50	1.39
1	B	66	CRQ	CD1-CG2	5.20	1.50	1.39
1	C	66	CRQ	CD1-CG2	5.14	1.49	1.39
1	B	66	CRQ	CD2-CG2	4.61	1.48	1.39
1	A	66	CRQ	CD2-CG2	4.40	1.48	1.39
1	C	66	CRQ	CD2-CG2	4.38	1.48	1.39
1	D	66	CRQ	CD2-CG2	4.36	1.48	1.39
1	B	66	CRQ	CA1-N1	4.20	1.36	1.27
1	C	66	CRQ	CA1-N1	3.92	1.36	1.27
1	A	66	CRQ	CA1-N1	3.91	1.36	1.27
1	D	66	CRQ	CA1-N1	3.81	1.36	1.27
1	C	66	CRQ	CG2-CB2	-3.39	1.40	1.46
1	D	66	CRQ	C1-CA1	-3.21	1.43	1.48
1	B	66	CRQ	CA3-N3	3.10	1.53	1.47
1	B	66	CRQ	CG2-CB2	-3.07	1.41	1.46
1	D	66	CRQ	CA3-N3	2.95	1.52	1.47
1	C	66	CRQ	C1-CA1	-2.94	1.43	1.48
1	D	66	CRQ	CG2-CB2	-2.94	1.41	1.46
1	C	66	CRQ	CA3-N3	2.91	1.52	1.47
1	A	66	CRQ	CA3-N3	2.85	1.52	1.47
1	A	66	CRQ	CG2-CB2	-2.76	1.41	1.46
1	B	66	CRQ	C1-CA1	-2.72	1.44	1.48
1	A	66	CRQ	C1-CA1	-2.69	1.44	1.48
1	D	66	CRQ	C1-N3	2.40	1.42	1.38
1	C	66	CRQ	C1-N3	2.38	1.42	1.38
1	A	66	CRQ	C1-N3	2.28	1.41	1.38
1	B	66	CRQ	C1-N3	2.06	1.41	1.38

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	CRQ	C3-CA3-N3	6.50	127.22	112.43
1	C	66	CRQ	C3-CA3-N3	6.38	126.95	112.43
1	B	66	CRQ	C3-CA3-N3	6.07	126.24	112.43
1	D	66	CRQ	C3-CA3-N3	6.02	126.14	112.43
1	B	66	CRQ	CA3-N3-C1	-2.77	122.94	128.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	66	CRQ	CA3-N3-C2	2.66	129.52	123.67
1	A	66	CRQ	CA3-N3-C1	-2.55	123.36	128.39
1	C	66	CRQ	CA3-N3-C1	-2.54	123.38	128.39
1	C	66	CRQ	CA3-N3-C2	2.50	129.17	123.67
1	A	66	CRQ	O3-C3-CA3	-2.48	114.29	125.47
1	C	66	CRQ	O3-C3-CA3	-2.47	114.36	125.47
1	A	66	CRQ	CA3-N3-C2	2.43	129.01	123.67
1	D	66	CRQ	CA3-N3-C2	2.42	128.99	123.67
1	B	66	CRQ	O3-C3-CA3	-2.38	114.77	125.47
1	D	66	CRQ	O3-C3-CA3	-2.36	114.84	125.47
1	D	66	CRQ	CA3-N3-C1	-2.36	123.75	128.39
1	C	66	CRQ	CD2-CE2-CZ	2.16	122.16	119.88
1	C	66	CRQ	C2-CA2-N2	2.14	110.48	108.95
1	D	66	CRQ	CD2-CE2-CZ	2.12	122.12	119.88
1	D	66	CRQ	CE2-CZ-CE1	-2.03	116.54	119.77
1	A	66	CRQ	CD2-CE2-CZ	2.02	122.01	119.88
1	A	66	CRQ	CE2-CZ-CE1	-2.01	116.57	119.77

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	D	66	CRQ	C1-CA1-CB1-CG1
1	B	66	CRQ	CA1-CB1-CG1-CD3
1	C	66	CRQ	CA1-CB1-CG1-CD3
1	A	66	CRQ	CA1-CB1-CG1-CD3
1	D	66	CRQ	CA1-CB1-CG1-CD3

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	D	66	CRQ	1	0
1	B	66	CRQ	2	0
1	C	66	CRQ	1	0
1	A	66	CRQ	1	0

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	217/223 (97%)	0.79	20 (9%) 14 15	16, 27, 51, 85	0
1	B	217/223 (97%)	0.77	16 (7%) 20 22	17, 27, 54, 87	0
1	C	217/223 (97%)	0.81	21 (9%) 13 14	16, 29, 61, 82	0
1	D	217/223 (97%)	0.72	19 (8%) 15 16	16, 28, 51, 84	0
All	All	868/892 (97%)	0.77	76 (8%) 15 16	16, 28, 54, 87	0

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	170	GLY	6.4
1	B	70	MET	4.6
1	B	225	LEU	4.4
1	A	225	LEU	4.2
1	B	169	ASP	4.1
1	C	225	LEU	4.0
1	C	70	MET	4.0
1	C	7	VAL	3.9
1	B	168	LYS	3.9
1	D	70	MET	3.6
1	A	89	GLU	3.6
1	D	225	LEU	3.6
1	D	170	GLY	3.5
1	A	170	GLY	3.5
1	D	168	LYS	3.4
1	C	77	ALA	3.4
1	C	110	ASP	3.4
1	C	168	LYS	3.3
1	D	69	SER	3.3
1	D	88	PRO	3.3
1	C	170	GLY	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	88	PRO	3.1
1	C	8	ILE	3.1
1	C	69	SER	3.0
1	D	90	GLY	3.0
1	C	6	ASN	2.9
1	B	186	PRO	2.9
1	A	70	MET	2.9
1	C	141	MET	2.8
1	B	116	GLY	2.8
1	B	223	LEU	2.8
1	A	168	LYS	2.8
1	C	90	GLY	2.8
1	D	77	ALA	2.7
1	C	94	GLU	2.7
1	A	77	ALA	2.7
1	C	113	LEU	2.7
1	C	169	ASP	2.7
1	C	89	GLU	2.7
1	A	216	ARG	2.6
1	B	89	GLU	2.6
1	C	71	VAL	2.6
1	C	78	ASP	2.6
1	B	185	LYS	2.6
1	D	200	ASP	2.5
1	A	90	GLY	2.5
1	A	223	LEU	2.5
1	B	113	LEU	2.5
1	A	45	LYS	2.5
1	A	69	SER	2.4
1	D	89	GLU	2.4
1	B	6	ASN	2.3
1	D	169	ASP	2.3
1	A	6	ASN	2.3
1	D	6	ASN	2.3
1	B	47	LYS	2.3
1	B	184	LYS	2.3
1	C	215	GLU	2.2
1	D	160	GLU	2.2
1	B	141	MET	2.2
1	D	116	GLY	2.2
1	D	137	GLN	2.2
1	D	91	PHE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	169	ASP	2.2
1	D	132	ASP	2.2
1	A	113	LEU	2.1
1	A	91	PHE	2.1
1	B	88	PRO	2.1
1	C	186	PRO	2.1
1	A	8	ILE	2.1
1	D	115	ASP	2.1
1	A	217	THR	2.1
1	C	112	SER	2.0
1	D	38	TYR	2.0
1	A	117	CYS	2.0
1	A	41	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CRQ	C	66	24/25	0.87	0.12	29,33,35,38	0
1	CRQ	B	66	24/25	0.88	0.12	28,31,33,34	0
1	CRQ	A	66	24/25	0.88	0.12	26,31,34,38	0
1	CRQ	D	66	24/25	0.88	0.11	30,34,36,38	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.