



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 06:51 PM UTC

PDB ID : 2ZHY / pdb_00002zhy
Title : Crystal structure of a pduO-type ATP:cobalamin adenosyltransferase from Burkholderia thailandensis
Authors : Moon, J.H.; Park, A.K.; Jang, E.H.; Kim, H.S.; Chi, Y.M.
Deposited on : 2008-02-11
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	183	13% 69% 10% • 20%
1	B	183	7% 81% 11% • 7%
1	C	183	14% 73% 17% • 9%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3803 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP:cob(I)alamin adenosyltransferase, putative.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	147	Total	C	N	O	S	0	0	0
			1095	687	205	200	3			
1	B	171	Total	C	N	O	S	0	0	0
			1266	785	240	238	3			
1	C	167	Total	C	N	O	S	0	0	0
			1242	771	236	232	3			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	64	Total	O	0	0
			64	64		
2	B	75	Total	O	0	0
			75	75		
2	C	61	Total	O	0	0
			61	61		

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	52.58Å 149.11Å 157.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.30 – 1.80 47.30 – 1.80	Depositor EDS
% Data completeness (in resolution range)	93.9 (47.30-1.80) 94.2 (47.30-1.80)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.33 (at 1.81Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.229 , 0.252 0.229 , 0.252	Depositor DCC
R_{free} test set	2749 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	19.4	Xtriage
Anisotropy	0.205	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 42.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3803	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/1114	0.82	1/1518 (0.1%)
1	B	0.37	0/1285	0.85	1/1746 (0.1%)
1	C	0.37	0/1261	0.83	3/1713 (0.2%)
All	All	0.38	0/3660	0.83	5/4977 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	78	ILE	N-CA-C	5.77	113.78	107.60
1	A	153	ASN	N-CA-C	-5.65	105.20	111.36
1	C	55	PRO	N-CA-C	-5.28	102.98	111.38
1	C	53	PRO	N-CA-C	-5.19	103.04	111.19
1	C	28	LYS	N-CA-C	5.13	118.64	112.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1095	0	1103	16	0
1	B	1266	0	1273	19	0
1	C	1242	0	1252	25	0
2	A	64	0	0	2	0
2	B	75	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	61	0	0	2	0
All	All	3803	0	3628	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:85:THR:H	1:C:88:HIS:HD2	1.12	0.94
1:C:21:GLY:HA3	1:C:78:ILE:HG12	1.59	0.84
1:A:85:THR:H	1:A:88:HIS:HD2	1.25	0.82
1:B:85:THR:H	1:B:88:HIS:HD2	1.27	0.79
1:A:107:GLU:HG2	1:A:108:GLU:HG2	1.64	0.79
1:B:85:THR:H	1:B:88:HIS:CD2	2.04	0.76
1:C:22:ASP:OD1	1:C:24:SER:HB3	1.88	0.74
1:A:110:ILE:HG22	1:A:176:LEU:HD23	1.75	0.69
1:C:85:THR:H	1:C:88:HIS:CD2	2.03	0.67
1:A:85:THR:H	1:A:88:HIS:CD2	2.11	0.65
1:A:107:GLU:H	1:A:107:GLU:CD	2.05	0.65
1:C:107:GLU:HB3	1:C:108:GLU:OE1	1.97	0.63
1:B:75:GLU:HG3	1:B:82:ALA:HA	1.83	0.59
1:B:147:ALA:HB3	1:B:148:PRO:HD3	1.85	0.59
1:A:144:LEU:HD23	2:A:227:HOH:O	2.06	0.54
1:C:7:LYS:H	1:C:7:LYS:HD2	1.73	0.54
1:C:11:ARG:HD2	2:C:229:HOH:O	2.08	0.52
1:C:145:ASN:HD21	1:C:147:ALA:HB3	1.72	0.52
1:B:32:ARG:HH21	1:B:142:GLU:HG2	1.75	0.51
1:C:164:ARG:O	1:C:168:ARG:HG2	2.11	0.51
1:C:147:ALA:HB3	1:C:148:PRO:HD3	1.93	0.50
1:B:110:ILE:HG22	1:B:176:LEU:HD23	1.93	0.49
1:C:72:LEU:HD23	1:C:72:LEU:O	2.13	0.49
1:A:98:HIS:CE1	1:A:102:GLN:HE21	2.31	0.48
1:B:7:LYS:HD2	1:B:7:LYS:N	2.28	0.48
1:A:116:ARG:NH2	1:C:116:ARG:NH2	2.60	0.48
1:C:17:THR:O	1:C:28:LYS:HE3	2.14	0.47
1:C:72:LEU:HD23	1:C:72:LEU:C	2.39	0.47
1:A:129:ARG:HH12	1:A:153:ASN:HD22	1.62	0.47
1:B:78:ILE:O	1:B:81:HIS:HB3	2.14	0.47
1:B:86:ASP:OD2	1:C:7:LYS:HD2	2.14	0.47
1:B:110:ILE:HD11	1:B:160:PHE:CE1	2.50	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:149:ARG:HG2	2:C:219:HOH:O	2.15	0.46
1:A:103:LEU:CD2	1:A:168:ARG:HE	2.29	0.46
1:A:32:ARG:HH11	1:A:77:CYS:HA	1.80	0.46
1:B:154:ARG:HH11	1:B:154:ARG:HG3	1.82	0.45
1:B:96:LEU:O	1:B:100:ASN:HB2	2.17	0.45
1:A:103:LEU:HD23	1:A:168:ARG:HE	1.80	0.45
1:C:21:GLY:HA3	1:C:78:ILE:CG1	2.39	0.44
1:C:162:LEU:O	1:C:166:LEU:HG	2.16	0.44
1:C:154:ARG:HG3	1:C:154:ARG:HH11	1.83	0.44
1:C:12:THR:HG22	1:C:25:ARG:HH22	1.82	0.43
1:A:98:HIS:NE2	1:A:102:GLN:NE2	2.66	0.43
1:C:96:LEU:O	1:C:100:ASN:HB2	2.19	0.42
1:A:33:ILE:HG12	2:A:190:HOH:O	2.18	0.42
1:B:106:LEU:HD11	1:B:110:ILE:CG2	2.49	0.42
1:C:78:ILE:CG2	1:C:81:HIS:HB2	2.50	0.42
1:A:145:ASN:O	1:A:148:PRO:HD2	2.19	0.42
1:B:75:GLU:HG3	1:B:81:HIS:O	2.20	0.42
1:B:85:THR:HA	2:B:216:HOH:O	2.19	0.42
1:B:106:LEU:HD11	1:B:110:ILE:HG23	2.01	0.41
1:C:78:ILE:O	1:C:79:PRO:C	2.63	0.41
1:C:58:VAL:HG11	1:C:166:LEU:HD21	2.02	0.41
1:A:129:ARG:HH12	1:A:153:ASN:ND2	2.19	0.41
1:C:92:LEU:HD11	1:C:151:TYR:CE1	2.56	0.41
1:B:154:ARG:HG3	1:B:154:ARG:NH1	2.36	0.40
1:C:22:ASP:OD1	1:C:24:SER:CB	2.65	0.40
1:B:7:LYS:HD2	1:B:7:LYS:H	1.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	145/183 (79%)	143 (99%)	2 (1%)	0	100	100
1	B	169/183 (92%)	168 (99%)	1 (1%)	0	100	100
1	C	165/183 (90%)	161 (98%)	4 (2%)	0	100	100
All	All	479/549 (87%)	472 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	109/137 (80%)	107 (98%)	2 (2%)	51	43
1	B	127/137 (93%)	123 (97%)	4 (3%)	35	23
1	C	125/137 (91%)	123 (98%)	2 (2%)	55	47
All	All	361/411 (88%)	353 (98%)	8 (2%)	45	34

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	107	GLU
1	A	120	LEU
1	B	49	LEU
1	B	100	ASN
1	B	144	LEU
1	B	168	ARG
1	C	5	LEU
1	C	100	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	81	HIS
1	A	88	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	100	ASN
1	A	102	GLN
1	A	153	ASN
1	B	88	HIS
1	B	100	ASN
1	B	145	ASN
1	B	153	ASN
1	C	81	HIS
1	C	88	HIS
1	C	100	ASN
1	C	102	GLN
1	C	145	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	147/183 (80%)	0.50	24 (16%) 4 3	14, 23, 50, 59	0
1	B	171/183 (93%)	0.18	13 (7%) 20 18	15, 25, 46, 53	0
1	C	167/183 (91%)	0.60	26 (15%) 5 4	15, 26, 57, 65	0
All	All	485/549 (88%)	0.42	63 (12%) 7 6	14, 25, 51, 65	0

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	80	GLY	8.8
1	A	79	PRO	7.3
1	A	81	HIS	6.4
1	C	106	LEU	6.0
1	C	21	GLY	5.6
1	C	81	HIS	5.3
1	A	141	SER	5.1
1	A	78	ILE	5.1
1	C	23	GLY	5.0
1	B	176	LEU	4.8
1	C	24	SER	4.5
1	C	110	ILE	4.4
1	A	143	PRO	4.3
1	C	108	GLU	4.2
1	C	171	GLY	4.1
1	A	82	ALA	4.0
1	A	77	CYS	4.0
1	C	109	PHE	4.0
1	A	140	ALA	3.8
1	C	107	GLU	3.7
1	C	80	GLY	3.5
1	A	144	LEU	3.5
1	A	109	PHE	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	107	GLU	3.3
1	B	175	VAL	3.3
1	A	145	ASN	3.3
1	B	108	GLU	3.1
1	A	176	LEU	3.0
1	A	106	LEU	3.0
1	C	105	PRO	3.0
1	A	102	GLN	3.0
1	A	107	GLU	2.9
1	C	82	ALA	2.9
1	C	145	ASN	2.8
1	B	171	GLY	2.8
1	B	79	PRO	2.8
1	B	82	ALA	2.7
1	A	30	ASP	2.7
1	B	173	ALA	2.6
1	C	20	LEU	2.5
1	C	56	ASP	2.5
1	A	171	GLY	2.4
1	B	106	LEU	2.4
1	B	168	ARG	2.4
1	C	101	GLY	2.4
1	B	109	PHE	2.4
1	C	22	ASP	2.3
1	C	78	ILE	2.3
1	C	10	THR	2.3
1	C	169	ALA	2.3
1	A	173	ALA	2.2
1	A	142	GLU	2.2
1	A	101	GLY	2.2
1	C	116	ARG	2.2
1	A	175	VAL	2.2
1	B	169	ALA	2.2
1	A	104	PRO	2.2
1	A	105	PRO	2.2
1	C	79	PRO	2.2
1	B	7	LYS	2.1
1	C	7	LYS	2.1
1	C	5	LEU	2.1
1	C	103	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.