



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 11:53 PM UTC

PDB ID : 2ZIW / pdb_00002ziw
Title : Crystal structure of the Mus81-Eme1 complex
Authors : Chang, J.H.; Kim, J.J.; Choi, J.M.; Lee, J.H.; Cho, Y.
Deposited on : 2008-02-25
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	311	5% 42% 32% 11% • 13%
2	B	341	9% 32% 26% 8% • 33%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3869 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mus81 protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	271	Total	C	N	O	S	Se	0	0	0
			2155	1368	373	399	11	4			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	302	MSE	-	initiating methionine	UNP Q6GML8

- Molecule 2 is a protein called Crossover junction endonuclease EME1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	228	Total	C	N	O	S	Se	0	0	0
			1714	1079	304	321	6	4			

There are 16 discrepancies between the modelled and reference sequences:



Chain	Residue	Modelled	Actual	Comment	Reference
B	230	MSE	-	expression tag	UNP Q96AY2
B	231	GLY	-	expression tag	UNP Q96AY2
B	232	SER	-	expression tag	UNP Q96AY2
B	233	SER	-	expression tag	UNP Q96AY2
B	234	HIS	-	expression tag	UNP Q96AY2
B	235	HIS	-	expression tag	UNP Q96AY2
B	236	HIS	-	expression tag	UNP Q96AY2
B	237	HIS	-	expression tag	UNP Q96AY2
B	238	HIS	-	expression tag	UNP Q96AY2
B	239	HIS	-	expression tag	UNP Q96AY2
B	240	SER	-	expression tag	UNP Q96AY2
B	241	GLN	-	expression tag	UNP Q96AY2
B	242	ASP	-	expression tag	UNP Q96AY2
B	243	PRO	-	expression tag	UNP Q96AY2
B	244	ASN	-	expression tag	UNP Q96AY2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	245	SER	-	expression tag	UNP Q96AY2

- Molecule 1: Mus81 protein





4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	88.96Å 88.96Å 169.33Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	37.56 – 2.80 37.56 – 2.80	Depositor EDS
% Data completeness (in resolution range)	95.7 (37.56-2.80) 98.2 (37.56-2.80)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.04 (at 2.72Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.272 , 0.327 0.292 , 0.343	Depositor DCC
R_{free} test set	1863 reflections (4.56%)	wwPDB-VP
Wilson B-factor (Å ²)	52.8	Xtriage
Anisotropy	0.177	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 32.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.023 for -h,-k,l	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	3869	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	5/2184 (0.2%)	1.07	14/2934 (0.5%)
2	B	0.63	3/1730 (0.2%)	1.06	9/2340 (0.4%)
All	All	0.64	8/3914 (0.2%)	1.06	23/5274 (0.4%)

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	541	MSE	SE-CE	-9.07	1.68	1.95
1	A	396	MSE	SE-CE	-6.95	1.74	1.95
1	A	475	MSE	SE-CE	-6.17	1.76	1.95
2	B	557	MSE	SE-CE	-5.74	1.78	1.95
2	B	276	MSE	SE-CE	-5.71	1.78	1.95

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	483	TYR	N-CA-C	-9.82	103.43	114.62
2	B	309	GLU	N-CA-C	8.86	127.41	109.60
1	A	328	CYS	N-CA-C	-8.82	100.64	111.40
2	B	422	GLN	N-CA-C	-8.29	99.36	110.55
1	A	588	LYS	N-CA-C	-7.04	101.24	110.33

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2155	0	2198	149	0
2	B	1714	0	1663	119	0
All	All	3869	0	3861	258	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 33.

The worst 5 of 258 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:312:TRP:HE1	1:A:314:LEU:HB2	1.05	1.13
1:A:316:PRO:HD3	1:A:487:THR:HA	1.30	1.08
2:B:566:LEU:HD12	2:B:566:LEU:H	1.13	1.06
1:A:487:THR:HG23	1:A:488:LEU:H	1.28	0.99
2:B:555:LEU:HD21	2:B:564:LEU:HD21	1.45	0.97

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	263/311 (85%)	218 (83%)	31 (12%)	14 (5%)	1	5
2	B	214/341 (63%)	179 (84%)	23 (11%)	12 (6%)	1	4
All	All	477/652 (73%)	397 (83%)	54 (11%)	26 (6%)	1	4

5 of 26 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	316	PRO
1	A	341	LEU
1	A	482	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	484	GLN
1	A	488	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	241/264 (91%)	212 (88%)	29 (12%)	5	17
2	B	173/282 (61%)	153 (88%)	20 (12%)	5	18
All	All	414/546 (76%)	365 (88%)	49 (12%)	5	17

5 of 49 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	591	ARG
2	B	312	VAL
2	B	252	ILE
2	B	287	VAL
2	B	316	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	416	GLN
2	B	427	GLN
2	B	556	GLN
2	B	524	GLN
1	A	452	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	267/311 (85%)	0.66	17 (6%) 25 18	21, 47, 68, 85	0
2	B	224/341 (65%)	0.83	30 (13%) 7 5	20, 48, 79, 86	0
All	All	491/652 (75%)	0.73	47 (9%) 13 10	20, 48, 72, 86	0

The worst 5 of 47 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	349	ASP	6.0
1	A	329	GLU	5.7
1	A	508	ALA	4.6
2	B	352	ALA	3.5
2	B	404	SER	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.