



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 10:40 PM UTC

PDB ID : 7ZIQ / pdb_00007ziq
Title : BK Polyomavirus VP1 in complex with 6'-Sialyllactose glycomacromolecules (aromatic linker)
Authors : Freytag, J.; Mueller, J.C.; Stehle, T.
Deposited on : 2022-04-08
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	275	3% 89% 6% 5%
1	BBB	275	2% 85% 8% 7%
1	CCC	275	2% 83% 5% 12%
1	DDD	275	3% 89% 8% .
1	EEE	275	3% 86% 9% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	AaA	3	 67% 33%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 10790 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Capsid protein VP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	260	Total	C	N	O	S	0	4	0
			2009	1264	347	385	13			
1	BBB	255	Total	C	N	O	S	0	3	0
			1970	1241	343	373	13			
1	CCC	243	Total	C	N	O	S	0	2	0
			1881	1181	328	359	13			
1	DDD	266	Total	C	N	O	S	0	2	0
			2044	1286	352	392	14			
1	EEE	260	Total	C	N	O	S	0	6	0
			2030	1278	351	388	13			

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
AAA	26	GLY	-	expression tag	UNP Q65613
AAA	27	SER	-	expression tag	UNP Q65613
AAA	28	HIS	-	expression tag	UNP Q65613
AAA	29	MET	-	expression tag	UNP Q65613
BBB	26	GLY	-	expression tag	UNP Q65613
BBB	27	SER	-	expression tag	UNP Q65613
BBB	28	HIS	-	expression tag	UNP Q65613
BBB	29	MET	-	expression tag	UNP Q65613
CCC	26	GLY	-	expression tag	UNP Q65613
CCC	27	SER	-	expression tag	UNP Q65613
CCC	28	HIS	-	expression tag	UNP Q65613
CCC	29	MET	-	expression tag	UNP Q65613
DDD	26	GLY	-	expression tag	UNP Q65613
DDD	27	SER	-	expression tag	UNP Q65613
DDD	28	HIS	-	expression tag	UNP Q65613
DDD	29	MET	-	expression tag	UNP Q65613
EEE	26	GLY	-	expression tag	UNP Q65613
EEE	27	SER	-	expression tag	UNP Q65613
EEE	28	HIS	-	expression tag	UNP Q65613

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
EEE	29	MET	-	expression tag	UNP Q65613

- Molecule 2 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	AaA	3	Total	C	N	O	0	0	0
			43	23	1	19			

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	BBB	1	Total	C	O	0	0
			4	2	2		
3	CCC	1	Total	C	O	0	0
			4	2	2		
3	EEE	1	Total	C	O	0	0
			4	2	2		
3	EEE	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is N-acetyl-alpha-neuraminic acid (CCD ID: SIA) (formula: $C_{11}H_{19}NO_9$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	BBB	1	Total	C	N	O	0	0
			21	11	1	9		
4	DDD	1	Total	C	N	O	0	0
			21	11	1	9		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	CCC	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	AAA	145	Total	O	0	0
			145	145		
6	BBB	150	Total	O	0	0
			150	150		
6	CCC	133	Total	O	0	0
			133	133		
6	DDD	158	Total	O	0	0
			158	158		
6	EEE	162	Total	O	0	0
			162	162		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Capsid protein VP1

Chain AAA: 




- Molecule 1: Capsid protein VP1

Chain BBB: 




- Molecule 1: Capsid protein VP1

Chain CCC: 




- Molecule 1: Capsid protein VP1

Chain DDD: 



- Molecule 1: Capsid protein VP1

Chain EEE: 



● Molecule 2: N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	144.33Å 151.84Å 63.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.50 – 1.90 48.50 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.50-1.90) 100.0 (48.50-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.09 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.187 , 0.226 0.195 , 0.232	Depositor DCC
R_{free} test set	3302 reflections (3.00%)	wwPDB-VP
Wilson B-factor (Å ²)	27.9	Xtriage
Anisotropy	0.063	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 33.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.019 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10790	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, BGC, EDO, GAL, SIA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	AAA	0.99	0/2066	1.21	2/2812 (0.1%)
1	BBB	0.98	0/2023	1.18	0/2750
1	CCC	0.96	0/1932	1.17	0/2627
1	DDD	1.00	0/2096	1.22	2/2853 (0.1%)
1	EEE	1.00	0/2089	1.18	1/2838 (0.0%)
All	All	0.98	0/10206	1.19	5/13880 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	98	LEU	CA-C-O	-5.61	114.78	121.11
1	AAA	181	ASN	CB-CA-C	5.50	115.88	110.71
1	DDD	104	CYS	CA-C-N	5.30	125.86	119.98
1	DDD	104	CYS	C-N-CA	5.30	125.86	119.98
1	EEE	255	LYS	CA-C-O	-5.23	115.01	120.55

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	2009	0	1942	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	BBB	1970	0	1910	16	0
1	CCC	1881	0	1814	7	0
1	DDD	2044	0	1987	13	0
1	EEE	2030	0	1971	10	0
2	AaA	43	0	37	0	0
3	BBB	4	0	6	0	0
3	CCC	4	0	6	0	0
3	EEE	8	0	12	0	0
4	BBB	21	0	18	0	0
4	DDD	21	0	18	0	0
5	CCC	7	0	10	0	0
6	AAA	145	0	0	1	0
6	BBB	150	0	0	3	0
6	CCC	133	0	0	0	0
6	DDD	158	0	0	4	0
6	EEE	162	0	0	0	0
All	All	10790	0	9731	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:BBB:109:MET:HE2	1:BBB:295:VAL:HG11	1.56	0.85
1:BBB:109:MET:CE	1:BBB:295:VAL:HG11	2.24	0.68
1:BBB:44:ILE:C	1:BBB:44:ILE:HD12	2.20	0.67
1:BBB:58:PRO:HA	1:BBB:83:LYS:HD2	1.80	0.64
1:BBB:296:LYS:O	1:BBB:297:ASN:HB2	1.97	0.64
1:DDD:32:VAL:N	6:DDD:603:HOH:O	2.32	0.62
1:BBB:109:MET:HE2	1:BBB:295:VAL:CG1	2.32	0.58
1:DDD:216:GLU:CG	6:DDD:712:HOH:O	2.51	0.57
1:BBB:44:ILE:HD12	1:BBB:44:ILE:O	2.07	0.54
1:EEE:113[B]:VAL:HG12	1:EEE:245:LEU:HD21	1.91	0.52
1:EEE:160[B]:GLU:OE2	1:EEE:214[B]:ARG:HB3	2.09	0.52
1:EEE:259:LEU:HD21	1:EEE:288:ILE:HD13	1.92	0.51
1:BBB:134:LYS:NZ	6:BBB:604:HOH:O	2.45	0.48
1:BBB:259:LEU:HD21	1:BBB:288:ILE:HD13	1.96	0.48
1:DDD:56:GLY:O	1:DDD:83:LYS:HD2	2.13	0.48
1:DDD:46:GLU:HG2	1:DDD:289:ARG:HG2	1.96	0.47
1:CCC:166:MET:HA	1:CCC:166:MET:HE2	1.95	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:BBB:109:MET:HE1	1:BBB:253:LEU:HD13	1.97	0.46
1:BBB:160:GLU:HB3	1:BBB:214[B]:ARG:CZ	2.46	0.46
1:AAA:114:THR:HB	1:AAA:241:THR:CG2	2.45	0.46
1:AAA:190:ASN:HA	6:AAA:412:HOH:O	2.16	0.46
1:DDD:113[A]:VAL:HG22	1:DDD:289:ARG:O	2.16	0.46
1:BBB:199:LYS:NZ	6:BBB:603:HOH:O	2.36	0.45
1:DDD:98:LEU:HD21	1:DDD:110:TRP:CZ2	2.51	0.45
1:EEE:98:LEU:HD21	1:EEE:110:TRP:CZ2	2.51	0.45
1:EEE:82[B]:ARG:NH2	1:EEE:201:ASN:OD1	2.46	0.45
1:CCC:259:LEU:HD21	1:CCC:288:ILE:HD13	1.98	0.44
1:AAA:113[B]:VAL:HG22	1:AAA:289:ARG:O	2.18	0.44
1:DDD:43:ALA:CB	1:DDD:292:LYS:HE3	2.47	0.44
1:AAA:46:GLU:OE1	1:AAA:289:ARG:NH2	2.51	0.44
1:AAA:100:GLU:HA	1:AAA:107:LEU:HD13	1.99	0.44
1:DDD:292:LYS:NZ	6:DDD:610:HOH:O	2.50	0.44
1:CCC:142:LYS:HE3	1:DDD:228:GLU:OE1	2.18	0.44
1:CCC:293:ARG:HG2	1:CCC:294:SER:N	2.33	0.43
1:EEE:114:THR:HB	1:EEE:241:THR:CG2	2.49	0.43
1:DDD:114:THR:HB	1:DDD:241:THR:CG2	2.49	0.43
1:CCC:114:THR:HB	1:CCC:241:THR:CG2	2.48	0.43
1:CCC:146:GLY:HA2	1:CCC:227:GLY:O	2.19	0.43
1:AAA:146:GLY:HA2	1:AAA:227:GLY:O	2.19	0.42
1:EEE:49[B]:CYS:SG	1:EEE:261:VAL:HG21	2.59	0.42
1:BBB:256:ALA:N	6:BBB:601:HOH:O	2.30	0.42
1:EEE:58:PRO:HA	1:EEE:83:LYS:HE3	2.02	0.42
1:AAA:259:LEU:HD21	1:AAA:288:ILE:HD13	2.02	0.42
1:CCC:117:THR:HA	1:CCC:285:TYR:O	2.21	0.41
1:DDD:109:MET:HE1	1:DDD:255:LYS:HD3	2.01	0.41
1:BBB:44:ILE:C	1:BBB:44:ILE:CD1	2.90	0.41
1:BBB:117:THR:HA	1:BBB:285:TYR:O	2.20	0.41
1:DDD:259:LEU:HD21	1:DDD:288:ILE:HD13	2.03	0.41
1:EEE:146:GLY:HA2	1:EEE:227:GLY:O	2.19	0.41
1:DDD:83:LYS:HG2	6:DDD:714:HOH:O	2.20	0.40
1:BBB:146:GLY:HA2	1:BBB:227:GLY:O	2.21	0.40
1:EEE:117:THR:HA	1:EEE:285:TYR:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	260/275 (94%)	250 (96%)	9 (4%)	1 (0%)	30	22
1	BBB	252/275 (92%)	239 (95%)	12 (5%)	1 (0%)	30	22
1	CCC	241/275 (88%)	231 (96%)	9 (4%)	1 (0%)	30	22
1	DDD	266/275 (97%)	251 (94%)	14 (5%)	1 (0%)	30	22
1	EEE	260/275 (94%)	248 (95%)	11 (4%)	1 (0%)	30	22
All	All	1279/1375 (93%)	1219 (95%)	55 (4%)	5 (0%)	30	22

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	188	VAL
1	BBB	188	VAL
1	CCC	188	VAL
1	DDD	188	VAL
1	EEE	188	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	220/236 (93%)	219 (100%)	1 (0%)	81	84
1	BBB	215/236 (91%)	212 (99%)	3 (1%)	59	59
1	CCC	206/236 (87%)	204 (99%)	2 (1%)	68	70
1	DDD	225/236 (95%)	224 (100%)	1 (0%)	84	87

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	EEE	223/236 (94%)	221 (99%)	2 (1%)	70	73
All	All	1089/1180 (92%)	1080 (99%)	9 (1%)	76	75

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	99	ASN
1	BBB	49[A]	CYS
1	BBB	49[B]	CYS
1	BBB	181	ASN
1	CCC	247	GLU
1	CCC	273	SER
1	DDD	101	ASP
1	EEE	137	GLU
1	EEE	273	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	AaA	1	2	12,12,12	0.43	0	17,17,17	0.51	0
2	GAL	AaA	2	2	11,11,12	0.25	0	15,15,17	0.51	0
2	SIA	AaA	3	2	20,20,21	0.60	0	21,28,31	1.14	2 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	AaA	1	2	-	0/2/22/22	0/1/1/1
2	GAL	AaA	2	2	-	0/2/19/22	0/1/1/1
2	SIA	AaA	3	2	-	0/18/34/38	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	AaA	3	SIA	O1B-C1-C2	2.95	120.38	112.71
2	AaA	3	SIA	O1A-C1-C2	-2.07	118.37	122.85


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	EEE	502	-	3,3,3	0.17	0	2,2,2	0.06	0
3	EDO	EEE	501	-	3,3,3	0.20	0	2,2,2	0.37	0
3	EDO	BBB	501	-	3,3,3	0.13	0	2,2,2	0.27	0
4	SIA	BBB	502	-	21,21,21	1.49	3 (14%)	24,31,31	1.34	2 (8%)
3	EDO	CCC	401	-	3,3,3	0.10	0	2,2,2	0.06	0
4	SIA	DDD	501	-	21,21,21	1.49	4 (19%)	24,31,31	1.64	4 (16%)
5	PEG	CCC	402	-	6,6,6	0.16	0	5,5,5	0.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	EEE	502	-	-	0/1/1/1	-
3	EDO	EEE	501	-	-	0/1/1/1	-
3	EDO	BBB	501	-	-	1/1/1/1	-
4	SIA	BBB	502	-	-	1/20/38/38	0/1/1/1
3	EDO	CCC	401	-	-	0/1/1/1	-
4	SIA	DDD	501	-	-	4/20/38/38	0/1/1/1
5	PEG	CCC	402	-	-	2/4/4/4	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	BBB	502	SIA	O2-C2	4.14	1.45	1.39
4	DDD	501	SIA	O2-C2	4.03	1.45	1.39
4	BBB	502	SIA	O6-C2	3.29	1.46	1.43
4	BBB	502	SIA	C3-C2	2.74	1.55	1.51
4	DDD	501	SIA	C3-C2	2.43	1.55	1.51
4	DDD	501	SIA	O1A-C1	2.27	1.29	1.22
4	DDD	501	SIA	C5-N5	2.03	1.49	1.45

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	DDD	501	SIA	O1A-C1-C2	-4.90	115.68	123.85
4	BBB	502	SIA	O1A-C1-C2	-3.72	117.64	123.85
4	DDD	501	SIA	O6-C6-C5	2.70	112.32	109.84
4	DDD	501	SIA	C3-C4-C5	2.36	113.37	109.72
4	DDD	501	SIA	C11-C10-N5	2.25	119.85	116.12
4	BBB	502	SIA	O6-C6-C7	2.10	109.94	106.65

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	DDD	501	SIA	C11-C10-N5-C5
4	DDD	501	SIA	O10-C10-N5-C5
3	BBB	501	EDO	O1-C1-C2-O2
4	DDD	501	SIA	O1A-C1-C2-O6
5	CCC	402	PEG	O1-C1-C2-O2

Continued on next page...


Continued from previous page...



Mol	Chain	Res	Type	Atoms
5	CCC	402	PEG	O2-C3-C4-O4
4	BBB	502	SIA	O1A-C1-C2-C3
4	DDD	501	SIA	O1B-C1-C2-C3


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	260/275 (94%)	0.03	7 (2%) 56 60	16, 30, 46, 92	4 (1%)
1	BBB	255/275 (92%)	-0.00	6 (2%) 59 63	15, 31, 56, 72	3 (1%)
1	CCC	243/275 (88%)	0.08	5 (2%) 63 67	19, 33, 54, 75	2 (0%)
1	DDD	266/275 (96%)	0.04	8 (3%) 52 56	16, 30, 57, 82	2 (0%)
1	EEE	260/275 (94%)	-0.05	7 (2%) 56 60	15, 29, 52, 73	6 (2%)
All	All	1284/1375 (93%)	0.02	33 (2%) 57 61	15, 30, 54, 92	17 (1%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	CCC	98	LEU	4.4
1	BBB	43	ALA	4.1
1	EEE	40	GLY	3.9
1	EEE	101	ASP	3.8
1	EEE	44	ILE	3.8
1	AAA	32	VAL	3.7
1	BBB	44	ILE	3.7
1	DDD	104	CYS	3.4
1	EEE	32	VAL	3.3
1	CCC	43	ALA	3.1
1	CCC	294	SER	3.1
1	AAA	107	LEU	3.1
1	DDD	102	LEU	3.0
1	CCC	108	LEU	3.0
1	BBB	38	LYS	2.9
1	AAA	297	ASN	2.9
1	AAA	100	GLU	2.9
1	AAA	99	ASN	2.4
1	BBB	255	LYS	2.3
1	DDD	37	VAL	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	EEE	39	THR	2.3
1	EEE	297	ASN	2.3
1	AAA	295	VAL	2.3
1	DDD	103	THR	2.3
1	DDD	106	ASN	2.2
1	DDD	297	ASN	2.1
1	CCC	44	ILE	2.1
1	EEE	100	GLU	2.1
1	AAA	296	LYS	2.0
1	DDD	50	PHE	2.0
1	BBB	297	ASN	2.0
1	BBB	98	LEU	2.0
1	DDD	107	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BGC	AaA	1	12/12	-	-	20,20,20,20	0
2	GAL	AaA	2	11/12	-	-	20,20,20,20	0
2	SIA	AaA	3	20/21	-	-	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain AaA:

2mF_o-DF_c (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
 and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	PEG	CCC	402	7/7	0.77	0.13	58,60,65,68	0
4	SIA	DDD	501	21/21	0.83	0.12	39,52,60,66	0
4	SIA	BBB	502	21/21	0.83	0.12	37,61,66,71	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	EDO	EEE	501	4/4	0.85	0.17	43,48,51,57	0
3	EDO	BBB	501	4/4	0.86	0.16	50,57,59,61	0
3	EDO	CCC	401	4/4	0.86	0.11	56,57,57,57	0
3	EDO	EEE	502	4/4	0.88	0.12	43,44,47,49	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around SIA DDD 501:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SIA BBB 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around EDO EEE 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around EDO BBB 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around EDO CCC 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around EDO EEE 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.