



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 02:41 PM UTC

PDB ID : 2ZJT / pdb_00002zjt
Title : Crystal structure of dna gyrase B' domain sheds lights on the mechanism for T-segment navigation
Authors : Fu, G.S.; Zhu, D.Y.; Hu, Y.L.; Wang, D.C.
Deposited on : 2008-03-10
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	247	4% 36% 32% 6% 25%
1	B	247	4% 37% 31% 6% 25%

2 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 2962 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA gyrase subunit B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	185	Total	C	N	O	S	0	0	0
			1481	946	254	275	6			
1	B	185	Total	C	N	O	S	0	0	0
			1481	946	254	275	6			

There are 38 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	481	MET	-	expression tag	UNP P0C5C5
A	482	THR	-	expression tag	UNP P0C5C5
A	483	PRO	-	expression tag	UNP P0C5C5
A	484	TYR	-	expression tag	UNP P0C5C5
A	485	ALA	-	expression tag	UNP P0C5C5
A	486	PHE	-	expression tag	UNP P0C5C5
A	715	LYS	-	expression tag	UNP P0C5C5
A	716	LEU	-	expression tag	UNP P0C5C5
A	717	ALA	-	expression tag	UNP P0C5C5
A	718	ALA	-	expression tag	UNP P0C5C5
A	719	ALA	-	expression tag	UNP P0C5C5
A	720	LEU	-	expression tag	UNP P0C5C5
A	721	GLU	-	expression tag	UNP P0C5C5
A	722	HIS	-	expression tag	UNP P0C5C5
A	723	HIS	-	expression tag	UNP P0C5C5
A	724	HIS	-	expression tag	UNP P0C5C5
A	725	HIS	-	expression tag	UNP P0C5C5
A	726	HIS	-	expression tag	UNP P0C5C5
A	727	HIS	-	expression tag	UNP P0C5C5
B	481	MET	-	expression tag	UNP P0C5C5
B	482	THR	-	expression tag	UNP P0C5C5
B	483	PRO	-	expression tag	UNP P0C5C5
B	484	TYR	-	expression tag	UNP P0C5C5
B	485	ALA	-	expression tag	UNP P0C5C5
B	486	PHE	-	expression tag	UNP P0C5C5

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	715	LYS	-	expression tag	UNP P0C5C5
B	716	LEU	-	expression tag	UNP P0C5C5
B	717	ALA	-	expression tag	UNP P0C5C5
B	718	ALA	-	expression tag	UNP P0C5C5
B	719	ALA	-	expression tag	UNP P0C5C5
B	720	LEU	-	expression tag	UNP P0C5C5
B	721	GLU	-	expression tag	UNP P0C5C5
B	722	HIS	-	expression tag	UNP P0C5C5
B	723	HIS	-	expression tag	UNP P0C5C5
B	724	HIS	-	expression tag	UNP P0C5C5
B	725	HIS	-	expression tag	UNP P0C5C5
B	726	HIS	-	expression tag	UNP P0C5C5
B	727	HIS	-	expression tag	UNP P0C5C5


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DNA gyrase subunit B



• Molecule 1: DNA gyrase subunit B



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.83Å 52.76Å 192.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.03 – 2.80 19.03 – 2.80	Depositor EDS
% Data completeness (in resolution range)	91.8 (19.03-2.80) 95.7 (19.03-2.80)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.36 (at 2.80Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.244 , 0.280 0.255 , 0.291	Depositor DCC
R_{free} test set	679 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	61.8	Xtriage
Anisotropy	0.059	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 56.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.480 for k,h,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	2962	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/1506	1.04	6/2027 (0.3%)
1	B	0.52	0/1506	1.04	7/2027 (0.3%)
All	All	0.52	0/3012	1.04	13/4054 (0.3%)

There are no bond length outliers.

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	659	GLU	N-CA-C	-11.40	98.17	112.88
1	B	659	GLU	N-CA-C	-11.29	98.32	112.88
1	A	660	LEU	N-CA-C	-10.17	101.42	114.04
1	B	660	LEU	N-CA-C	-10.01	101.62	114.04
1	B	581	THR	N-CA-C	-6.20	105.87	113.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1481	0	1498	108	0
1	B	1481	0	1498	108	0
All	All	2962	0	2996	208	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 35.

The worst 5 of 208 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:577:GLN:HB3	1:B:580:SER:HB3	1.37	1.04
1:B:657:ALA:O	1:B:658:LYS:HB3	1.59	1.03
1:A:577:GLN:HB3	1:A:580:SER:HB3	1.39	1.01
1:A:657:ALA:O	1:A:658:LYS:HB3	1.59	0.99
1:A:686:LEU:O	1:A:690:LEU:HG	1.73	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	179/247 (72%)	154 (86%)	19 (11%)	6 (3%)	3	11
1	B	179/247 (72%)	154 (86%)	19 (11%)	6 (3%)	3	11
All	All	358/494 (72%)	308 (86%)	38 (11%)	12 (3%)	3	11

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	575	ASP
1	A	656	ASP
1	B	575	ASP
1	B	656	ASP
1	A	576	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	158/207 (76%)	143 (90%)	15 (10%)	8	26
1	B	158/207 (76%)	143 (90%)	15 (10%)	8	26
All	All	316/414 (76%)	286 (90%)	30 (10%)	8	26

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	672	LEU
1	B	663	THR
1	B	562	ARG
1	B	672	LEU
1	B	652	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	604	GLN
1	B	613	GLN
1	B	674	GLN
1	A	613	GLN
1	A	604	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	185/247 (74%)	0.34	10 (5%) 31 24	30, 56, 87, 96	0
1	B	185/247 (74%)	0.35	11 (5%) 28 21	31, 56, 87, 96	0
All	All	370/494 (74%)	0.34	21 (5%) 29 22	30, 56, 88, 96	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	615	SER	4.0
1	A	498	GLU	3.7
1	B	576	GLY	3.2
1	A	652	LEU	3.1
1	B	498	GLU	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.