



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 11:17 AM UTC

PDB ID : 3ZJG / pdb_00003zjg
Title : A20 OTU domain with irreversibly oxidised Cys103 from 60 min H₂O₂ soak.
Authors : Kulathu, Y.; Garcia, F.J.; Mevissen, T.E.T.; Busch, M.; Arnaudo, N.; Carroll, K.S.; Barford, D.; Komander, D.
Deposited on : 2013-01-17
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1188 (1.92-1.92)
Clashscore	190562	1209 (1.92-1.92)
Ramachandran outliers	187476	1195 (1.92-1.92)
Sidechain outliers	187428	1195 (1.92-1.92)
RSRZ outliers	180081	1188 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	366	8% 77% 10% • 11%
2	B	366	12% 77% 11% • 10%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5495 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TUMOR NECROSIS FACTOR ALPHA-INDUCED PROTEIN 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	327	Total	C	N	O	S	0	6	0
			2622	1687	450	469	16			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	114	SER	GLY	engineered mutation	UNP P21580

- Molecule 2 is a protein called TUMOR NECROSIS FACTOR ALPHA-INDUCED PROTEIN 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	328	Total	C	N	O	S	0	2	0
			2611	1680	454	462	15			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	114	SER	GLY	engineered mutation	UNP P21580

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	3	Total	Cl	0	0
			3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	136	Total	O	0	0
			136	136		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	123	Total 123	O 123	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: TUMOR NECROSIS FACTOR ALPHA-INDUCED PROTEIN 3



• Molecule 2: TUMOR NECROSIS FACTOR ALPHA-INDUCED PROTEIN 3



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	43.80Å 69.17Å 85.00Å 98.57° 100.51° 97.10°	Depositor
Resolution (Å)	34.00 – 1.92 34.00 – 1.92	Depositor EDS
% Data completeness (in resolution range)	95.3 (34.00-1.92) 95.3 (34.00-1.92)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 1.92Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.184 , 0.220 0.186 , 0.221	Depositor DCC
R_{free} test set	3528 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	31.6	Xtriage
Anisotropy	0.320	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 66.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5495	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSD, OCS, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.03	0/2678	1.07	4/3641 (0.1%)
2	B	0.93	0/2666	1.06	5/3622 (0.1%)
All	All	0.98	0/5344	1.06	9/7263 (0.1%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	212	ASP	N-CA-C	6.51	118.45	111.36
2	B	57	CYS	CA-C-N	5.94	127.27	119.84
2	B	57	CYS	C-N-CA	5.94	127.27	119.84
1	A	57	CYS	CA-C-N	5.24	126.39	119.84
1	A	57	CYS	C-N-CA	5.24	126.39	119.84
2	B	278	ARG	N-CA-C	-5.19	102.00	110.20
1	A	344	ASP	N-CA-C	-5.16	105.74	111.36
1	A	322	THR	N-CA-C	5.15	116.90	109.07
2	B	85	TRP	N-CA-C	-5.06	106.96	113.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2622	0	2508	22	0
2	B	2611	0	2491	28	0
3	B	3	0	0	0	0
4	A	136	0	0	1	0
4	B	123	0	0	1	0
All	All	5495	0	4999	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:66:LYS:O	2:B:96:LYS:NZ	2.07	0.88
2:B:50:MET:HE2	2:B:51:PHE:H	1.53	0.72
2:B:237:HIS:HD2	4:B:2004:HOH:O	1.78	0.66
2:B:34:ASN:HD21	2:B:36:ILE:HG22	1.60	0.66
2:B:137:ASN:HD22	2:B:237:HIS:CE1	2.14	0.65
2:B:11:TYR:HA	2:B:341:LEU:HD23	1.79	0.65
1:A:91:LYS:NZ	4:A:2054:HOH:O	2.31	0.64
1:A:134:ASP:OD1	1:A:136:ARG:HD3	1.97	0.63
2:B:282:ARG:NE	2:B:284:GLU:OE2	2.32	0.62
2:B:52:ARG:HE	2:B:53:THR:H	1.45	0.62
2:B:34:ASN:ND2	2:B:36:ILE:HG22	2.17	0.58
1:A:16:ARG:HG2	2:B:344:ASP:OD1	2.04	0.57
1:A:214[A]:MET:SD	1:A:214[A]:MET:N	2.75	0.57
1:A:49:GLU:HB2	1:A:333:ALA:HB2	1.86	0.56
2:B:52:ARG:HE	2:B:53:THR:N	2.03	0.56
1:A:7:PRO:HG2	1:A:345:TYR:CE2	2.42	0.55
1:A:241:GLN:H	1:A:241:GLN:CD	2.15	0.54
2:B:11:TYR:HA	2:B:341:LEU:CD2	2.37	0.54
1:A:351:HIS:ND1	2:B:23:GLU:OE2	2.39	0.54
1:A:348:LEU:HD11	2:B:16:ARG:HA	1.91	0.51
2:B:50:MET:CE	2:B:50:MET:HA	2.41	0.51
2:B:50:MET:HA	2:B:50:MET:HE3	1.93	0.50
2:B:214:MET:SD	2:B:287:LYS:HE3	2.52	0.50
2:B:214:MET:HE1	2:B:287:LYS:HG3	1.95	0.49
1:A:270:ILE:O	1:A:270:ILE:HG13	2.13	0.49
2:B:33:THR:O	2:B:34:ASN:HB3	2.12	0.48
2:B:214:MET:HB2	2:B:230:GLY:HA3	1.93	0.48
2:B:49:GLU:OE2	2:B:271:ARG:NH1	2.47	0.47
1:A:17:LYS:NZ	1:A:344:ASP:OD2	2.39	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:335:LEU:HD22	2:B:339:ILE:HG13	1.96	0.47
1:A:103:CSD:OD1	1:A:256:HIS:HA	2.15	0.47
2:B:48:LEU:HD11	2:B:342:VAL:HG21	1.97	0.46
1:A:143:GLN:HE21	1:A:164:TRP:CG	2.34	0.46
2:B:49:GLU:OE2	2:B:262[A]:THR:OG1	2.32	0.46
1:A:20:LYS:HZ3	1:A:24:ARG:HD2	1.81	0.45
1:A:69:ILE:HD12	1:A:93[B]:VAL:CG2	2.47	0.45
2:B:209:VAL:HG11	2:B:229:VAL:HG22	1.99	0.44
1:A:293:ASP:O	1:A:297:GLU:HG2	2.17	0.44
1:A:20:LYS:NZ	1:A:24:ARG:HD2	2.32	0.44
2:B:200:CYS:HB2	2:B:205:ARG:O	2.16	0.44
2:B:32:PRO:HG2	2:B:36:ILE:HG12	2.01	0.43
1:A:11:TYR:HA	1:A:341:LEU:HD13	2.02	0.42
1:A:119:ASP:O	1:A:120:LEU:HB2	2.20	0.42
2:B:275:LEU:C	2:B:286[B]:LEU:HD13	2.46	0.41
1:A:17:LYS:HZ2	1:A:344:ASP:CG	2.25	0.40
1:A:48:LEU:HD11	1:A:342:VAL:CG2	2.51	0.40
1:A:48:LEU:HD11	1:A:342:VAL:HG21	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	324/366 (88%)	315 (97%)	7 (2%)	2 (1%)	21	10
2	B	321/366 (88%)	312 (97%)	6 (2%)	3 (1%)	14	5
All	All	645/732 (88%)	627 (97%)	13 (2%)	5 (1%)	16	6

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	319	HIS
2	B	34	ASN
1	A	338	GLU
2	B	265	ASP
2	B	58	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	272/335 (81%)	253 (93%)	19 (7%)	14	4
2	B	267/335 (80%)	254 (95%)	13 (5%)	22	8
All	All	539/670 (80%)	507 (94%)	32 (6%)	18	5

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	LEU
1	A	20	LYS
1	A	24	ARG
1	A	36	ILE
1	A	89	VAL
1	A	90	ARG
1	A	120	LEU
1	A	134	ASP
1	A	136	ARG
1	A	147	LEU
1	A	179	THR
1	A	214[A]	MET
1	A	214[B]	MET
1	A	241	GLN
1	A	245	ARG
1	A	282	ARG
1	A	325	ILE
1	A	335	LEU
1	A	343	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	12	LEU
2	B	31	LYS
2	B	36	ILE
2	B	50	MET
2	B	52	ARG
2	B	58	PRO
2	B	120	LEU
2	B	186	LEU
2	B	229	VAL
2	B	282	ARG
2	B	300	GLU
2	B	335	LEU
2	B	347	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	110	GLN
1	A	241	GLN
2	B	34	ASN
2	B	189	ASN
2	B	237	HIS
2	B	315	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSD	A	103	1	4,7,8	4.89	1 (25%)	1,8,10	1.62	0
2	OCS	B	103	2	6,8,9	1.46	1 (16%)	7,11,13	1.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	A	103	1	-	1/2/6/8	-
2	OCS	B	103	2	-	0/4/7/9	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	103	CSD	OD1-SG	-9.57	1.39	1.47
2	B	103	OCS	CB-CA	2.89	1.55	1.53

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	103	CSD	CA-CB-SG-OD1

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	103	CSD	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	326/366 (89%)	0.34	28 (8%) 16 19	11, 40, 77, 97	6 (1%)
2	B	327/366 (89%)	0.51	45 (13%) 6 9	13, 45, 81, 100	2 (0%)
All	All	653/732 (89%)	0.43	73 (11%) 10 13	11, 42, 80, 100	8 (1%)

All (73) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	266	SER	8.0
1	A	11	TYR	6.3
1	A	227	LEU	6.0
2	B	227	LEU	5.1
1	A	6	LEU	5.0
1	A	356	TRP	4.9
2	B	134	ASP	4.9
2	B	58	PRO	4.8
1	A	321	THR	4.7
2	B	166	ASP	4.7
1	A	214[A]	MET	4.7
2	B	214	MET	4.5
1	A	319	HIS	4.1
2	B	133	THR	3.9
2	B	33	THR	3.9
1	A	320	GLY	3.9
2	B	8	GLN	3.8
2	B	213	LYS	3.7
2	B	153	VAL	3.6
2	B	152	PHE	3.6
1	A	218	LEU	3.5
2	B	9	ALA	3.4
2	B	266	SER	3.4
1	A	54	CYS	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	357	GLN	3.2
2	B	265	ASP	3.2
2	B	186	LEU	3.2
2	B	60	PHE	3.2
2	B	55	GLN	3.1
2	B	341	LEU	3.0
1	A	267	GLY	3.0
2	B	337	LYS	3.0
2	B	59	GLN	2.9
2	B	5	VAL	2.9
2	B	132	GLU	2.9
2	B	11	TYR	2.9
2	B	180	PRO	2.8
1	A	160	ASP	2.8
1	A	57	CYS	2.8
1	A	161	THR	2.8
1	A	215	LEU	2.7
2	B	356	TRP	2.7
2	B	64	ILE	2.7
1	A	8	GLN	2.7
2	B	169	ASP	2.7
1	A	341	LEU	2.6
2	B	335	LEU	2.6
2	B	338	GLU	2.6
2	B	54	CYS	2.5
2	B	57	CYS	2.5
1	A	316	GLY	2.5
2	B	165	ASN	2.5
1	A	63	ILE	2.5
2	B	170	ASN	2.5
2	B	336	PRO	2.4
2	B	6	LEU	2.3
1	A	254	SER	2.3
2	B	282	ARG	2.3
1	A	59	GLN	2.3
1	A	186	LEU	2.3
1	A	9	ALA	2.2
1	A	217	SER	2.2
2	B	63	ILE	2.2
2	B	61	ARG	2.2
2	B	164	TRP	2.2
2	B	177	THR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	185	GLY	2.1
2	B	340	ASN	2.1
1	A	133	THR	2.1
2	B	179	THR	2.1
1	A	148	LYS	2.1
2	B	36	ILE	2.1
1	A	55	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	OCS	B	103	9/10	0.96	0.09	34,35,52,54	0
1	CSD	A	103	8/9	0.97	0.06	18,23,45,48	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CL	B	1358	1/1	0.98	0.05	23,23,23,23	0
3	CL	B	1360	1/1	0.99	0.07	28,28,28,28	0
3	CL	B	1359	1/1	1.00	0.06	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.