



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 10:03 AM UTC

PDB ID : 4ZJV / pdb_00004zjv
Title : crystal structure of EGFR kinase domain in complex with Mitogen-inducible gene 6 protein
Authors : Eck, M.J.; Park, E.; Lee, B.
Deposited on : 2015-04-29
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	76% 8% 15%
1	B	331	71% 10% 18%
2	C	70	7% 63% 6% 31%
2	D	70	6% 49% 11% 37%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5274 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Epidermal growth factor receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	282	Total	C	N	O	S	0	0	0
			2255	1455	381	404	15			
1	B	273	Total	C	N	O	S	0	1	0
			2198	1425	371	387	15			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	692	GLY	-	expression tag	UNP P00533
A	693	SER	-	expression tag	UNP P00533
A	694	THR	-	expression tag	UNP P00533
B	692	GLY	-	expression tag	UNP P00533
B	693	SER	-	expression tag	UNP P00533
B	694	THR	-	expression tag	UNP P00533

- Molecule 2 is a protein called ERBB receptor feedback inhibitor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	48	Total	C	N	O	P S	0	0	0
			377	243	56	73	2 3			
2	D	44	Total	C	N	O	P S	0	0	0
			348	225	50	68	2 3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	390	CYS	SER	engineered mutation	UNP Q9UJM3
D	390	CYS	SER	engineered mutation	UNP Q9UJM3


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	41	Total 41	O 41	0	0
3	B	38	Total 38	O 38	0	0
3	C	6	Total 6	O 6	0	0
3	D	11	Total 11	O 11	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Epidermal growth factor receptor




- Molecule 1: Epidermal growth factor receptor



- Molecule 2: ERBB receptor feedback inhibitor 1



- Molecule 2: ERBB receptor feedback inhibitor 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.46Å 81.86Å 96.19Å 90.00° 106.67° 90.00°	Depositor
Resolution (Å)	50.00 – 2.70 50.00 – 2.70	Depositor EDS
% Data completeness (in resolution range)	89.8 (50.00-2.70) 89.9 (50.00-2.70)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, R_{free}	0.179 , 0.232 0.186 , 0.236	Depositor DCC
R_{free} test set	1099 reflections (4.56%)	wwPDB-VP
Wilson B-factor (Å ²)	45.3	Xtriage
Anisotropy	0.160	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 41.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.011 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5274	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/2304	0.79	0/3116
1	B	0.60	0/2250	0.78	0/3043
2	C	0.78	0/353	0.82	0/480
2	D	0.96	0/324	0.98	0/442
All	All	0.66	0/5231	0.80	0/7081

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2255	0	2309	20	0
1	B	2198	0	2267	24	0
2	C	377	0	368	1	0
2	D	348	0	336	10	0
3	A	41	0	0	0	0
3	B	38	0	0	0	0
3	C	6	0	0	0	0
3	D	11	0	0	0	0
All	All	5274	0	5280	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:934:PRO:HG2	2:D:346:MET:HE2	1.57	0.87
2:D:389:VAL:HG12	2:D:390:CYS:H	1.63	0.64
1:B:926:ILE:HG23	1:B:931:GLU:HB2	1.81	0.62
1:B:762:GLU:HG3	1:B:859:ALA:HB3	1.82	0.61
2:D:382:ILE:HG22	2:D:389:VAL:HG11	1.83	0.60
1:A:847:THR:HG22	1:A:850:HIS:HB3	1.85	0.58
1:B:714:LYS:NZ	1:B:787:GLN:OE1	2.37	0.58
1:B:722:ALA:HB1	1:B:875:LYS:HD2	1.88	0.54
1:A:982:GLN:HG3	1:B:813:TYR:CE2	2.44	0.53
1:A:789:ILE:HD12	1:A:789:ILE:N	2.24	0.53
2:D:361:SER:O	2:D:362:LYS:HD2	2.08	0.53
1:A:839:ALA:HB3	1:A:841:ARG:HG2	1.92	0.51
1:B:715:ILE:HG12	1:B:729:GLY:HA2	1.92	0.50
1:A:982:GLN:NE2	1:B:808:ASN:O	2.46	0.49
1:B:823:LYS:HA	1:B:965:ILE:HD11	1.95	0.49
1:A:841:ARG:NH2	1:A:877:PRO:HB3	2.29	0.48
1:A:736:GLU:HB3	1:A:738:VAL:HG12	1.96	0.48
1:B:769:VAL:HG22	1:B:776:ARG:HA	1.96	0.48
1:A:832:ARG:NH2	1:A:863:GLY:O	2.44	0.47
1:A:783:THR:OG1	1:A:784:SER:N	2.47	0.47
1:A:808:ASN:O	1:A:808:ASN:OD1	2.32	0.47
1:A:871:ALA:HB3	1:A:889:ARG:HG2	1.95	0.47
1:B:882:ALA:HA	1:B:898:TRP:CD2	2.49	0.47
1:A:747:LEU:HD11	1:A:856:PHE:O	2.15	0.46
2:D:389:VAL:HG12	2:D:390:CYS:N	2.28	0.46
1:A:823:LYS:HA	1:A:965:ILE:HD11	1.97	0.46
1:B:769:VAL:O	1:B:776:ARG:HG3	2.16	0.45
1:A:841:ARG:HH22	1:A:877:PRO:CB	2.30	0.45
1:B:877:PRO:O	1:B:881:MET:HG3	2.16	0.44
1:B:832:ARG:NH2	1:B:863:GLY:O	2.48	0.44
1:B:898:TRP:CE3	1:B:951:TRP:HA	2.53	0.44
1:A:762:GLU:HG3	1:A:859:ALA:HB3	2.00	0.44
2:D:379:ILE:HA	2:D:395:PTR:O	2.18	0.43
1:A:775:CYS:SG	1:A:854:THR:HG23	2.58	0.43
1:A:882:ALA:HA	1:A:898:TRP:CD2	2.54	0.43
1:A:982:GLN:CG	1:B:813:TYR:CE2	3.02	0.43
1:B:980:VAL:HB	2:D:338:LEU:HD11	2.00	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:934:PRO:HG3	2:D:347:PRO:O	2.19	0.42
1:B:706:ILE:HD12	1:B:706:ILE:N	2.35	0.42
1:B:961:PHE:HA	1:B:964:LEU:HD12	2.01	0.42
1:B:806:LYS:HG2	1:B:910:PHE:HB3	2.02	0.41
1:A:952:MET:HE3	2:C:357:LYS:HD3	2.02	0.41
2:D:382:ILE:CG2	2:D:389:VAL:HG11	2.50	0.41
1:B:798:LEU:HD22	1:B:851:VAL:HG11	2.03	0.41
2:D:361:SER:O	2:D:362:LYS:CB	2.68	0.41
1:B:802:VAL:CG1	1:B:910:PHE:HA	2.51	0.40
1:A:856:PHE:CD1	1:A:856:PHE:N	2.89	0.40
1:B:747:LEU:N	1:B:747:LEU:CD1	2.84	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	278/331 (84%)	273 (98%)	5 (2%)	0	100	100
1	B	268/331 (81%)	258 (96%)	10 (4%)	0	100	100
2	C	40/70 (57%)	38 (95%)	2 (5%)	0	100	100
2	D	36/70 (51%)	33 (92%)	3 (8%)	0	100	100
All	All	622/802 (78%)	602 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/290 (85%)	244 (99%)	3 (1%)	63	84
1	B	243/290 (84%)	236 (97%)	7 (3%)	37	67
2	C	43/62 (69%)	42 (98%)	1 (2%)	44	73
2	D	40/62 (64%)	38 (95%)	2 (5%)	22	48
All	All	573/704 (81%)	560 (98%)	13 (2%)	44	73

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	726	VAL
1	A	751	THR
1	A	889	ARG
1	B	726	VAL
1	B	747	LEU
1	B	762	GLU
1	B	803	ARG
1	B	804	GLU
1	B	875	LYS
1	B	926	ILE
2	C	388	LYS
2	D	361	SER
2	D	389	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	805	HIS
1	A	826	ASN
1	A	888	HIS
1	B	826	ASN
1	B	888	HIS
1	B	894	GLN
2	C	343	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	PTR	D	394	2	15,16,17	0.75	0	17,22,24	0.83	0
2	PTR	D	395	2	15,16,17	0.74	0	17,22,24	1.05	0
2	PTR	C	394	2	15,16,17	0.68	0	17,22,24	0.99	1 (5%)
2	PTR	C	395	2	15,16,17	0.64	0	17,22,24	1.01	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PTR	D	394	2	-	0/10/11/13	0/1/1/1
2	PTR	D	395	2	-	4/10/11/13	0/1/1/1
2	PTR	C	394	2	-	0/10/11/13	0/1/1/1
2	PTR	C	395	2	-	2/10/11/13	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	395	PTR	OH-P-O1P	-2.26	101.94	109.48
2	C	394	PTR	OH-P-O1P	-2.17	102.24	109.48

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	395	PTR	CZ-OH-P-O1P
2	D	395	PTR	C-CA-CB-CG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	D	395	PTR	CZ-OH-P-O1P
2	D	395	PTR	N-CA-CB-CG
2	C	395	PTR	CZ-OH-P-O3P
2	D	395	PTR	CZ-OH-P-O2P

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	395	PTR	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	282/331 (85%)	-0.33	3 (1%) 78 76	24, 40, 71, 94	0
1	B	273/331 (82%)	-0.30	1 (0%) 88 87	24, 42, 69, 86	1 (0%)
2	C	46/70 (65%)	0.03	5 (10%) 10 9	22, 41, 78, 84	0
2	D	42/70 (60%)	0.35	4 (9%) 14 11	29, 44, 78, 97	0
All	All	643/802 (80%)	-0.24	13 (2%) 65 62	22, 41, 73, 97	1 (0%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	389	VAL	4.1
2	D	390	CYS	3.6
2	C	389	VAL	3.3
2	D	382	ILE	3.1
2	C	376	VAL	3.0
1	A	703	LEU	2.6
1	A	841	ARG	2.6
2	C	336	LYS	2.3
1	A	808	ASN	2.3
1	B	739	LYS	2.1
2	C	398	PRO	2.1
2	C	390	CYS	2.1
2	D	398	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PTR	D	395	16/17	0.86	0.15	50,71,109,111	0
2	PTR	D	394	16/17	0.95	0.08	44,47,58,59	0
2	PTR	C	395	16/17	0.95	0.11	33,54,85,91	0
2	PTR	C	394	16/17	0.96	0.07	30,39,45,54	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.