



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 30, 2026 – 03:50 AM UTC

PDB ID : 6ZJK / pdb_00006zjk
Title : Ribonucleotide reductase R2 subunit from Clostridium botulinum
Authors : Martinez-Carranza, M.; Stenmark, P.
Deposited on : 2020-06-29
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	FAILED
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

MolProbity failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10892 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribonucleoside-diphosphate reductase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	316	Total	C	N	O	S	0	1	0
			2666	1716	441	498	11			
1	A	317	Total	C	N	O	S	0	1	0
			2673	1719	443	501	10			
1	D	315	Total	C	N	O	S	0	3	0
			2667	1718	440	498	11			
1	B	316	Total	C	N	O	S	0	0	0
			2661	1712	441	498	10			

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-21	MET	-	initiating methionine	UNP B1KYY8
C	-20	HIS	-	expression tag	UNP B1KYY8
C	-19	HIS	-	expression tag	UNP B1KYY8
C	-18	HIS	-	expression tag	UNP B1KYY8
C	-17	HIS	-	expression tag	UNP B1KYY8
C	-16	HIS	-	expression tag	UNP B1KYY8
C	-15	HIS	-	expression tag	UNP B1KYY8
C	-14	SER	-	expression tag	UNP B1KYY8
C	-13	SER	-	expression tag	UNP B1KYY8
C	-12	GLY	-	expression tag	UNP B1KYY8
C	-11	VAL	-	expression tag	UNP B1KYY8
C	-10	ASP	-	expression tag	UNP B1KYY8
C	-9	LEU	-	expression tag	UNP B1KYY8
C	-8	GLY	-	expression tag	UNP B1KYY8
C	-7	THR	-	expression tag	UNP B1KYY8
C	-6	GLU	-	expression tag	UNP B1KYY8
C	-5	ASN	-	expression tag	UNP B1KYY8
C	-4	LEU	-	expression tag	UNP B1KYY8
C	-3	TYR	-	expression tag	UNP B1KYY8
C	-2	PHE	-	expression tag	UNP B1KYY8
C	-1	GLN	-	expression tag	UNP B1KYY8

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	0	SER	-	expression tag	UNP B1KYY8
A	-21	MET	-	initiating methionine	UNP B1KYY8
A	-20	HIS	-	expression tag	UNP B1KYY8
A	-19	HIS	-	expression tag	UNP B1KYY8
A	-18	HIS	-	expression tag	UNP B1KYY8
A	-17	HIS	-	expression tag	UNP B1KYY8
A	-16	HIS	-	expression tag	UNP B1KYY8
A	-15	HIS	-	expression tag	UNP B1KYY8
A	-14	SER	-	expression tag	UNP B1KYY8
A	-13	SER	-	expression tag	UNP B1KYY8
A	-12	GLY	-	expression tag	UNP B1KYY8
A	-11	VAL	-	expression tag	UNP B1KYY8
A	-10	ASP	-	expression tag	UNP B1KYY8
A	-9	LEU	-	expression tag	UNP B1KYY8
A	-8	GLY	-	expression tag	UNP B1KYY8
A	-7	THR	-	expression tag	UNP B1KYY8
A	-6	GLU	-	expression tag	UNP B1KYY8
A	-5	ASN	-	expression tag	UNP B1KYY8
A	-4	LEU	-	expression tag	UNP B1KYY8
A	-3	TYR	-	expression tag	UNP B1KYY8
A	-2	PHE	-	expression tag	UNP B1KYY8
A	-1	GLN	-	expression tag	UNP B1KYY8
A	0	SER	-	expression tag	UNP B1KYY8
D	-21	MET	-	initiating methionine	UNP B1KYY8
D	-20	HIS	-	expression tag	UNP B1KYY8
D	-19	HIS	-	expression tag	UNP B1KYY8
D	-18	HIS	-	expression tag	UNP B1KYY8
D	-17	HIS	-	expression tag	UNP B1KYY8
D	-16	HIS	-	expression tag	UNP B1KYY8
D	-15	HIS	-	expression tag	UNP B1KYY8
D	-14	SER	-	expression tag	UNP B1KYY8
D	-13	SER	-	expression tag	UNP B1KYY8
D	-12	GLY	-	expression tag	UNP B1KYY8
D	-11	VAL	-	expression tag	UNP B1KYY8
D	-10	ASP	-	expression tag	UNP B1KYY8
D	-9	LEU	-	expression tag	UNP B1KYY8
D	-8	GLY	-	expression tag	UNP B1KYY8
D	-7	THR	-	expression tag	UNP B1KYY8
D	-6	GLU	-	expression tag	UNP B1KYY8
D	-5	ASN	-	expression tag	UNP B1KYY8
D	-4	LEU	-	expression tag	UNP B1KYY8
D	-3	TYR	-	expression tag	UNP B1KYY8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-2	PHE	-	expression tag	UNP B1KYY8
D	-1	GLN	-	expression tag	UNP B1KYY8
D	0	SER	-	expression tag	UNP B1KYY8
B	-21	MET	-	initiating methionine	UNP B1KYY8
B	-20	HIS	-	expression tag	UNP B1KYY8
B	-19	HIS	-	expression tag	UNP B1KYY8
B	-18	HIS	-	expression tag	UNP B1KYY8
B	-17	HIS	-	expression tag	UNP B1KYY8
B	-16	HIS	-	expression tag	UNP B1KYY8
B	-15	HIS	-	expression tag	UNP B1KYY8
B	-14	SER	-	expression tag	UNP B1KYY8
B	-13	SER	-	expression tag	UNP B1KYY8
B	-12	GLY	-	expression tag	UNP B1KYY8
B	-11	VAL	-	expression tag	UNP B1KYY8
B	-10	ASP	-	expression tag	UNP B1KYY8
B	-9	LEU	-	expression tag	UNP B1KYY8
B	-8	GLY	-	expression tag	UNP B1KYY8
B	-7	THR	-	expression tag	UNP B1KYY8
B	-6	GLU	-	expression tag	UNP B1KYY8
B	-5	ASN	-	expression tag	UNP B1KYY8
B	-4	LEU	-	expression tag	UNP B1KYY8
B	-3	TYR	-	expression tag	UNP B1KYY8
B	-2	PHE	-	expression tag	UNP B1KYY8
B	-1	GLN	-	expression tag	UNP B1KYY8
B	0	SER	-	expression tag	UNP B1KYY8

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0

- Molecule 3 is FE (III) ION (CCD ID: FE) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	2	Total Fe 2 2	0	0
3	A	2	Total Fe 2 2	0	0
3	D	2	Total Fe 2 2	0	0
3	B	2	Total Fe 2 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	75	Total O 75 75	0	0
4	A	40	Total O 40 40	0	0
4	D	38	Total O 38 38	0	0
4	B	40	Total O 40 40	0	0

MolProbity failed to run properly - this section is therefore empty.

3 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	210.83Å 61.85Å 154.34Å 90.00° 127.46° 90.00°	Depositor
Resolution (Å)	83.67 – 2.00 83.67 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.8 (83.67-2.00) 86.2 (83.67-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.72 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.18rc4-3812-000	Depositor
R, R_{free}	0.222 , 0.237 0.224 , 0.238	Depositor DCC
R_{free} test set	5488 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å ²)	34.0	Xtriage
Anisotropy	0.518	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 38.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10892	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

4.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	C	401	-	5,5,5	0.53	0	5,5,5	1.05	0
2	GOL	C	402	-	5,5,5	0.45	0	5,5,5	0.70	0
2	GOL	B	401	-	5,5,5	0.85	0	5,5,5	1.07	0
2	GOL	A	401	-	5,5,5	0.28	0	5,5,5	0.78	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	C	401	-	-	0/4/4/4	-
2	GOL	C	402	-	-	0/4/4/4	-
2	GOL	B	401	-	-	0/4/4/4	-
2	GOL	A	401	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data

5.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	317/364 (87%)	0.63	14 (4%) 39 38	29, 50, 74, 101	1 (0%)
1	B	316/364 (86%)	0.52	8 (2%) 58 58	39, 58, 83, 97	0
1	C	316/364 (86%)	0.25	4 (1%) 75 74	33, 46, 69, 93	1 (0%)
1	D	315/364 (86%)	0.56	7 (2%) 62 61	29, 56, 77, 101	3 (0%)
All	All	1264/1456 (86%)	0.49	33 (2%) 57 56	29, 53, 77, 101	5 (0%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	LEU	5.2
1	D	2	LEU	4.3
1	B	2	LEU	3.8
1	A	-1	GLN	3.7
1	A	270	LEU	3.6
1	B	315	LEU	3.6
1	A	0	SER	3.3
1	B	144	GLU	3.2
1	A	1	MET	3.1
1	D	252	MET	3.1
1	A	237	ILE	3.0
1	A	301	ILE	2.9
1	C	303	LYS	2.8
1	B	232	ASN	2.6
1	A	302	ASN	2.6
1	A	300	GLU	2.5
1	C	65	TYR	2.5
1	A	168	ILE	2.5
1	C	270	LEU	2.4
1	B	150	GLU	2.4
1	A	22	ASN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	57	ASN	2.3
1	C	2	LEU	2.3
1	D	270	LEU	2.3
1	D	237	ILE	2.2
1	D	137	LYS	2.2
1	D	272	LEU	2.2
1	A	235	SER	2.2
1	B	265	THR	2.1
1	A	167	GLU	2.0
1	A	236	HIS	2.0
1	B	169	HIS	2.0
1	D	254	VAL	2.0

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	B	401	6/6	0.85	0.15	59,62,65,70	0
3	FE	B	403	1/1	0.90	0.10	57,57,57,57	1
2	GOL	C	401	6/6	0.91	0.09	45,47,50,52	0
2	GOL	C	402	6/6	0.92	0.11	43,47,51,54	6
2	GOL	A	401	6/6	0.93	0.10	47,48,49,49	6
3	FE	D	401	1/1	0.94	0.09	58,58,58,58	1
3	FE	A	403	1/1	0.95	0.06	48,48,48,48	1
3	FE	B	402	1/1	0.97	0.06	52,52,52,52	1
3	FE	C	404	1/1	0.97	0.06	48,48,48,48	1
3	FE	A	402	1/1	0.98	0.03	46,46,46,46	1
3	FE	C	403	1/1	0.99	0.03	43,43,43,43	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	FE	D	402	1/1	0.99	0.04	52,52,52,52	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around FE D 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE A 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE C 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around FE A 402:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FE C 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





5.5 Other polymers ⓘ

There are no such residues in this entry.