



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 04:11 PM UTC

PDB ID : 2ZKL / pdb_00002zkl
Title : Crystal Structure of capsular polysaccharide assembling protein CapF from staphylococcus aureus
Authors : Miyafusa, T.; Tanaka, Y.; Yao, M.; Tanaka, I.; Tsumoto, K.
Deposited on : 2008-03-25
Resolution : 2.61 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4951 (2.64-2.60)
Clashscore	190562	5303 (2.64-2.60)
Ramachandran outliers	187476	5217 (2.64-2.60)
Sidechain outliers	187428	5217 (2.64-2.60)
RSRZ outliers	180081	4950 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	369	5% 59% 29% 9% ..

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2940 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Capsular polysaccharide synthesis enzyme Cap5F.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	361	Total	C	N	O	S	0	0	0
			2916	1866	482	561	7			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	17	Total	O	0	0
			17	17		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Capsular polysaccharide synthesis enzyme Cap5F



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	119.64Å 119.64Å 129.45Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	10.00 – 2.61 10.00 – 2.61	Depositor EDS
% Data completeness (in resolution range)	98.7 (10.00-2.61) 97.0 (10.00-2.61)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.85 (at 2.61Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.238 , 0.273 0.237 , 0.270	Depositor DCC
R_{free} test set	1682 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å ²)	63.8	Xtriage
Anisotropy	0.093	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 48.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.017 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2940	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/2981	1.01	19/4048 (0.5%)

There are no bond length outliers.

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	136	LYS	N-CA-C	13.76	125.97	110.97
1	A	47	ALA	N-CA-C	7.13	120.29	110.24
1	A	280	LYS	CA-C-N	7.13	128.75	119.84
1	A	280	LYS	C-N-CA	7.13	128.75	119.84
1	A	336	THR	N-CA-C	-6.31	101.60	110.50
1	A	227	LEU	N-CA-C	5.86	120.05	113.01
1	A	363	ASP	N-CA-C	-5.85	103.42	110.44
1	A	137	TRP	N-CA-C	5.79	119.87	113.21
1	A	283	ILE	N-CA-C	5.64	116.58	110.21
1	A	295	GLU	N-CA-C	5.52	117.02	109.18
1	A	313	ASP	N-CA-C	5.49	116.61	108.60
1	A	170	LEU	N-CA-C	5.44	117.96	109.52
1	A	293	LYS	N-CA-C	-5.36	102.35	110.28
1	A	48	ASP	N-CA-C	-5.36	107.40	114.31
1	A	122	GLY	N-CA-C	5.35	122.48	115.36
1	A	24	THR	N-CA-C	5.33	116.37	108.86
1	A	6	THR	N-CA-C	-5.32	102.61	110.48
1	A	361	GLN	CA-C-N	-5.13	114.95	120.03
1	A	361	GLN	C-N-CA	-5.13	114.95	120.03

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2916	0	2873	121	0
2	A	1	0	0	0	0
3	A	6	0	7	2	0
4	A	17	0	0	0	0
All	All	2940	0	2880	121	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

All (121) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:279:SER:HB3	1:A:283:ILE:HD11	1.29	1.13
1:A:95:ILE:HD12	1:A:131:PRO:HA	1.40	1.01
1:A:336:THR:HG21	1:A:367:LEU:HB3	1.56	0.87
1:A:301:SER:HB3	1:A:348:VAL:CG1	2.04	0.87
1:A:204:THR:HG22	1:A:207:GLU:H	1.40	0.86
1:A:279:SER:CB	1:A:283:ILE:HD11	2.07	0.85
1:A:168:LEU:HB3	1:A:170:LEU:HD21	1.61	0.81
1:A:338:ASN:HD22	1:A:338:ASN:C	1.94	0.75
1:A:131:PRO:CD	1:A:171:ASN:HD22	2.04	0.70
1:A:131:PRO:HD2	1:A:171:ASN:HD22	1.56	0.70
1:A:24:THR:HG23	1:A:26:HIS:H	1.56	0.69
1:A:309:ARG:HH11	1:A:369:VAL:HB	1.58	0.68
1:A:280:LYS:O	1:A:283:ILE:HD12	1.92	0.67
1:A:164:ARG:O	1:A:204:THR:HG23	1.94	0.67
1:A:101:ASN:HD21	1:A:103:TYR:HB3	1.60	0.66
1:A:336:THR:CG2	1:A:367:LEU:HB3	2.25	0.65
1:A:270:ASP:C	1:A:271:ARG:HG2	2.23	0.64
1:A:20:LEU:O	1:A:24:THR:HG22	1.97	0.64
1:A:78:ASP:O	1:A:81:THR:HB	1.97	0.64
1:A:265:PHE:O	1:A:266:ILE:HD12	1.97	0.64
1:A:75:HIS:HE1	1:A:79:ILE:HD11	1.63	0.63
1:A:301:SER:HB3	1:A:348:VAL:HG12	1.81	0.62
1:A:305:VAL:HG22	1:A:342:LEU:HD22	1.81	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:297:PHE:CE1	1:A:351:MET:HE3	2.36	0.61
1:A:36:LYS:HB2	1:A:39:GLU:OE1	2.00	0.61
1:A:51:VAL:HG12	1:A:53:LEU:HD13	1.82	0.60
1:A:305:VAL:HG22	1:A:342:LEU:CD2	2.31	0.60
1:A:204:THR:HG23	1:A:206:GLY:H	1.68	0.59
1:A:24:THR:HG23	1:A:26:HIS:N	2.17	0.59
1:A:168:LEU:HB3	1:A:170:LEU:CD2	2.32	0.58
1:A:4:VAL:CG2	1:A:50:ILE:HG13	2.34	0.58
1:A:309:ARG:NH1	1:A:369:VAL:HB	2.17	0.58
1:A:304:GLY:HA2	1:A:342:LEU:HB2	1.86	0.58
1:A:309:ARG:HD3	1:A:369:VAL:CG1	2.33	0.57
1:A:172:TYR:CD2	1:A:174:ASP:HB2	2.39	0.57
1:A:279:SER:HB3	1:A:283:ILE:CD1	2.19	0.56
1:A:23:THR:HG21	1:A:181:LYS:NZ	2.20	0.56
1:A:307:ARG:O	1:A:337:HIS:HA	2.06	0.56
1:A:368:GLU:CD	1:A:369:VAL:H	2.13	0.55
1:A:368:GLU:O	1:A:369:VAL:HB	2.07	0.55
1:A:369:VAL:OXT	1:A:369:VAL:HG12	2.07	0.55
1:A:77:LEU:HB3	1:A:117:TYR:CE2	2.41	0.54
1:A:95:ILE:CD1	1:A:131:PRO:HA	2.28	0.54
1:A:75:HIS:CE1	1:A:79:ILE:HD11	2.42	0.54
1:A:336:THR:HG21	1:A:367:LEU:CB	2.33	0.54
1:A:310:HIS:HD2	1:A:312:ASN:H	1.56	0.54
1:A:297:PHE:HE1	1:A:351:MET:HE3	1.73	0.53
1:A:294:ASN:C	1:A:295:GLU:HG3	2.34	0.53
1:A:51:VAL:HG12	1:A:53:LEU:CD1	2.39	0.53
1:A:309:ARG:HD3	1:A:369:VAL:HG11	1.91	0.53
1:A:23:THR:HG21	1:A:181:LYS:HZ1	1.72	0.53
1:A:259:ARG:HG2	1:A:259:ARG:HH11	1.76	0.51
1:A:17:LYS:HE2	1:A:30:GLU:OE2	2.11	0.51
1:A:316:ILE:HD11	1:A:369:VAL:HG13	1.93	0.51
1:A:83:ASN:OD1	1:A:85:LYS:HG3	2.11	0.50
1:A:101:ASN:ND2	1:A:103:TYR:HB3	2.25	0.50
1:A:292:THR:HA	1:A:333:VAL:HG11	1.93	0.50
1:A:191:GLU:O	1:A:192:ASN:HB2	2.12	0.49
1:A:204:THR:HB	1:A:207:GLU:OE2	2.11	0.49
1:A:257:ASP:OD1	1:A:257:ASP:C	2.56	0.49
1:A:2:ASN:ND2	1:A:27:HIS:HB3	2.29	0.48
1:A:131:PRO:O	1:A:133:LEU:HD13	2.14	0.48
1:A:259:ARG:HG2	1:A:259:ARG:NH1	2.28	0.48
1:A:4:VAL:HG12	1:A:29:PHE:HB2	1.95	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:305:VAL:HG13	1:A:320:TYR:CD2	2.48	0.48
1:A:316:ILE:CD1	1:A:369:VAL:HG13	2.43	0.48
1:A:369:VAL:CG1	1:A:369:VAL:OXT	2.62	0.47
1:A:95:ILE:HD13	1:A:95:ILE:H	1.80	0.47
1:A:234:ASP:O	1:A:238:THR:CG2	2.62	0.47
1:A:204:THR:CG2	1:A:207:GLU:HG3	2.45	0.47
1:A:103:TYR:O	1:A:107:LYS:HG2	2.14	0.47
1:A:258:ASP:O	1:A:280:LYS:HD2	2.15	0.47
1:A:204:THR:CG2	1:A:206:GLY:H	2.28	0.46
1:A:169:THR:C	1:A:170:LEU:HD22	2.40	0.46
1:A:24:THR:HG23	1:A:25:ASP:N	2.29	0.46
1:A:73:LEU:HD23	1:A:113:LEU:HD23	1.97	0.46
1:A:194:VAL:O	1:A:196:THR:HG22	2.15	0.46
1:A:74:ASP:OD1	3:A:3147:GOL:O2	2.33	0.46
1:A:191:GLU:HB3	1:A:196:THR:HG21	1.96	0.46
1:A:338:ASN:C	1:A:338:ASN:ND2	2.65	0.45
1:A:297:PHE:CD1	1:A:351:MET:HG2	2.51	0.45
1:A:240:LEU:O	1:A:243:LEU:HB2	2.17	0.45
1:A:41:GLU:O	1:A:45:LEU:HD22	2.17	0.45
1:A:218:SER:OG	1:A:223:THR:HG22	2.17	0.45
1:A:291:HIS:HB2	1:A:363:ASP:OD1	2.17	0.45
1:A:204:THR:HB	1:A:207:GLU:HG3	1.98	0.44
1:A:94:SER:C	1:A:96:GLN:H	2.24	0.44
1:A:78:ASP:HB2	3:A:3147:GOL:H2	1.99	0.44
1:A:304:GLY:HA3	1:A:340:GLU:O	2.18	0.44
1:A:15:ASN:ND2	1:A:136:LYS:NZ	2.66	0.44
1:A:223:THR:CG2	1:A:224:LEU:N	2.81	0.44
1:A:357:PHE:CE1	1:A:359:PRO:HG3	2.53	0.43
1:A:132:ASN:O	1:A:170:LEU:HA	2.18	0.43
1:A:204:THR:HG22	1:A:207:GLU:HG3	1.99	0.43
1:A:23:THR:O	1:A:23:THR:CG2	2.66	0.43
1:A:204:THR:HG22	1:A:207:GLU:N	2.21	0.43
1:A:259:ARG:HA	1:A:283:ILE:HG21	2.01	0.43
1:A:221:ASP:OD1	1:A:223:THR:HB	2.19	0.43
1:A:254:MET:HG2	1:A:263:THR:OG1	2.19	0.43
1:A:313:ASP:OD1	1:A:314:ASP:N	2.49	0.42
1:A:188:PRO:HG2	1:A:190:ILE:HD11	2.01	0.42
1:A:197:VAL:HA	1:A:198:PRO:HD3	1.77	0.42
1:A:360:ASN:C	1:A:362:PRO:HD3	2.44	0.42
1:A:111:GLU:O	1:A:115:ARG:HG3	2.20	0.42
1:A:176:ILE:HD13	1:A:197:VAL:HG21	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:4:VAL:HG22	1:A:50:ILE:HG13	2.01	0.42
1:A:94:SER:C	1:A:96:GLN:N	2.77	0.42
1:A:280:LYS:O	1:A:283:ILE:CD1	2.63	0.42
1:A:336:THR:HG23	1:A:367:LEU:HD23	2.03	0.41
1:A:305:VAL:O	1:A:339:ILE:HA	2.21	0.41
1:A:23:THR:O	1:A:23:THR:HG22	2.20	0.41
1:A:65:PHE:N	1:A:65:PHE:CD2	2.88	0.41
1:A:95:ILE:HD13	1:A:95:ILE:N	2.36	0.41
1:A:234:ASP:O	1:A:238:THR:HG22	2.20	0.41
1:A:101:ASN:O	1:A:102:PRO:C	2.64	0.41
1:A:265:PHE:CE1	1:A:266:ILE:HD13	2.56	0.41
1:A:191:GLU:HG2	1:A:192:ASN:OD1	2.19	0.40
1:A:296:LYS:HB2	1:A:352:TRP:HB3	2.02	0.40
1:A:307:ARG:HB2	1:A:338:ASN:ND2	2.36	0.40
1:A:259:ARG:HD3	1:A:259:ARG:H	1.86	0.40
1:A:326:LEU:HD12	1:A:326:LEU:HA	1.88	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	357/369 (97%)	334 (94%)	20 (6%)	3 (1%)	16 31

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	34	GLN
1	A	281	PRO
1	A	361	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	328/336 (98%)	281 (86%)	47 (14%)	3 6

All (47) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	LEU
1	A	24	THR
1	A	34	GLN
1	A	40	LEU
1	A	45	LEU
1	A	53	LEU
1	A	81	THR
1	A	90	LEU
1	A	95	ILE
1	A	100	ASP
1	A	105	GLU
1	A	120	GLU
1	A	133	LEU
1	A	139	LYS
1	A	167	GLU
1	A	168	LEU
1	A	170	LEU
1	A	196	THR
1	A	200	VAL
1	A	204	THR
1	A	211	LEU
1	A	212	LEU
1	A	219	ARG
1	A	220	LEU
1	A	223	THR
1	A	224	LEU
1	A	227	LEU
1	A	238	THR
1	A	240	LEU
1	A	243	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	252	LEU
1	A	266	ILE
1	A	271	ARG
1	A	274	VAL
1	A	281	PRO
1	A	283	ILE
1	A	291	HIS
1	A	305	VAL
1	A	314	ASP
1	A	326	LEU
1	A	336	THR
1	A	338	ASN
1	A	342	LEU
1	A	344	ASP
1	A	348	VAL
1	A	361	GLN
1	A	368	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	ASN
1	A	15	ASN
1	A	26	HIS
1	A	75	HIS
1	A	101	ASN
1	A	141	ASN
1	A	165	ASN
1	A	171	ASN
1	A	255	ASN
1	A	277	ASN
1	A	288	HIS
1	A	294	ASN
1	A	310	HIS
1	A	338	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	3147	-	5,5,5	0.99	0	5,5,5	0.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	3147	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	3147	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	3147	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	361/369 (97%)	0.03	20 (5%)	30 25	39, 60, 87, 98	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	38	GLU	4.1
1	A	143	ASN	3.9
1	A	344	ASP	3.6
1	A	55	GLY	3.6
1	A	78	ASP	2.9
1	A	314	ASP	2.9
1	A	160	GLN	2.8
1	A	281	PRO	2.8
1	A	27	HIS	2.8
1	A	360	ASN	2.5
1	A	158	GLU	2.4
1	A	361	GLN	2.4
1	A	191	GLU	2.4
1	A	368	GLU	2.3
1	A	56	VAL	2.3
1	A	35	THR	2.3
1	A	283	ILE	2.2
1	A	86	LYS	2.0
1	A	359	PRO	2.0
1	A	162	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	A	3147	6/6	0.59	0.23	90,92,94,95	0
2	ZN	A	400	1/1	0.97	0.04	57,57,57,57	0

6.5 Other polymers [i](#)

There are no such residues in this entry.