



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 05:49 PM UTC

PDB ID : 3ZKG / pdb_00003zkg
Title : BACE2 MUTANT APO STRUCTURE
Authors : Banner, D.W.; Kuglstatter, A.; Benz, J.; Stihle, M.; Ruf, A.
Deposited on : 2013-01-23
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	386	
1	B	386	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5954 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called BETA-SECRETASE 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	364	Total	C	N	O	S	0	3	0
			2848	1838	454	543	13			
1	B	359	Total	C	N	O	S	0	1	0
			2785	1793	445	534	13			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	269	ALA	GLU	engineered mutation	UNP Q9Y5Z0
B	269	ALA	GLU	engineered mutation	UNP Q9Y5Z0

- Molecule 2 is 2-[3-(2-HYDROXY-1,1-DIHYDROXYMETHYL-ETHYLAMINO)-PROPYLAMINO]-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: B3P) (formula: $C_{11}H_{26}N_2O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			19	11	2	6		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	199	Total	O	0	0
			199	199		
3	B	103	Total	O	0	0
			103	103		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: BETA-SECRETASE 2



• Molecule 1: BETA-SECRETASE 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	46.89Å 88.47Å 98.57Å 90.00° 96.58° 90.00°	Depositor
Resolution (Å)	32.82 – 1.90 32.82 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.2 (32.82-1.90) 97.2 (32.82-1.90)	Depositor EDS
R_{merge}	0.01	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.33 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.228 , 0.273 (Not available) , 0.286	Depositor DCC
R_{free} test set	3195 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	19.4	Xtriage
Anisotropy	0.088	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	5954	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: B3P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.77	0/2916	0.86	0/3961
1	B	0.57	0/2851	0.79	0/3873
All	All	0.68	0/5767	0.82	0/7834

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2848	0	2772	32	0
1	B	2785	0	2708	25	0
2	A	19	0	26	0	0
3	A	199	0	0	3	0
3	B	103	0	0	1	0
All	All	5954	0	5506	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:48:ASP:HB2	1:A:134[B]:ILE:HD11	1.56	0.88
1:B:48:ASP:HB3	1:B:134[B]:ILE:HD11	1.69	0.74
1:A:46:LEU:HD23	1:A:134[B]:ILE:HD13	1.71	0.72
1:B:316:GLN:HB2	1:B:379:VAL:HG23	1.70	0.72
1:A:48:ASP:CB	1:A:134[B]:ILE:HD11	2.18	0.71
1:A:170:MET:CE	1:A:351:GLU:HG3	2.22	0.69
1:A:46:LEU:HD23	1:A:134[B]:ILE:CD1	2.22	0.69
1:A:34:MET:HE1	1:A:135:LEU:HD22	1.76	0.68
1:A:170:MET:HE2	1:A:351:GLU:HG3	1.78	0.66
1:B:48:ASP:CB	1:B:134[B]:ILE:HD11	2.28	0.63
1:A:236:ASP:HB2	1:A:341:ASN:O	2.00	0.62
1:B:245:THR:O	1:B:348:THR:HG23	2.04	0.58
1:A:48:ASP:OD2	1:A:134[A]:ILE:HD11	2.03	0.58
1:B:317:LEU:HG	1:B:379:VAL:HG22	1.85	0.57
1:B:290:TRP:CD2	1:B:316:GLN:HG3	2.39	0.56
1:A:30:TYR:CE1	1:A:170:MET:HE3	2.40	0.56
1:B:320:GLN:HE22	1:B:348:THR:CG2	2.17	0.56
1:B:170:MET:HE1	1:B:347:ALA:HA	1.88	0.55
1:A:170:MET:HE1	1:A:347:ALA:O	2.07	0.54
1:A:34:MET:CE	1:A:135:LEU:HD22	2.39	0.53
1:A:210:TRP:CG	1:A:211:TYR:H	2.28	0.52
1:B:46:LEU:HD23	1:B:134[B]:ILE:HD13	1.92	0.51
1:A:320:GLN:HE22	1:A:348:THR:HG21	1.75	0.51
1:A:32:LEU:HD23	1:A:34:MET:HE2	1.93	0.51
1:A:141:THR:CG2	1:A:210:TRP:HA	2.41	0.51
1:A:33:GLU:OE1	3:A:2005:HOH:O	2.20	0.50
1:B:320:GLN:HE22	1:B:348:THR:HG22	1.76	0.50
1:A:280:LEU:HD12	1:B:280:LEU:HD12	1.94	0.49
1:A:333:ARG:NH1	3:A:2183:HOH:O	2.45	0.49
1:A:46:LEU:HB3	1:A:134[B]:ILE:HD12	1.94	0.48
1:B:271:SER:N	3:B:2085:HOH:O	2.47	0.48
1:B:220:GLU:HB3	1:B:297:SER:HB2	1.95	0.47
1:A:109:THR:HA	3:A:2081:HOH:O	2.13	0.47
1:B:192:ILE:HG23	1:B:354:TYR:HE2	1.80	0.47
1:B:111:PHE:CZ	1:B:156:LEU:HD13	2.50	0.46
1:A:56:VAL:CG2	1:A:115:ILE:HD11	2.45	0.45
1:B:341:ASN:OD1	1:B:341:ASN:N	2.50	0.45
1:B:316:GLN:HB2	1:B:379:VAL:CG2	2.44	0.44
1:B:35:LEU:HB3	1:B:40:PRO:HB3	1.99	0.44
1:A:265:SER:O	1:A:266:LEU:CB	2.66	0.43
1:A:170:MET:HE1	1:A:347:ALA:CA	2.48	0.43
1:A:95:PHE:O	1:A:116:ALA:HA	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:33:GLU:CD	1:A:42:LYS:HD2	2.44	0.43
1:B:234:ASN:O	1:B:235:ALA:C	2.61	0.43
1:A:170:MET:HE1	1:A:347:ALA:HB1	2.01	0.42
1:B:214:ILE:HD12	1:B:300:LEU:HD22	2.02	0.42
1:B:170:MET:HE3	1:B:351:GLU:HG3	2.01	0.42
1:A:312:THR:HB	1:A:381:GLU:HB2	2.01	0.42
1:B:290:TRP:CG	1:B:316:GLN:HG3	2.54	0.42
1:A:193:GLU:HA	1:A:194:PRO:HD2	1.96	0.42
1:A:170:MET:HE2	1:A:351:GLU:CG	2.47	0.42
1:A:104:PRO:O	1:A:108:ASN:ND2	2.51	0.42
1:B:134[B]:ILE:HD12	1:B:135:LEU:N	2.35	0.41
1:B:320:GLN:HE22	1:B:348:THR:HG21	1.85	0.41
1:A:265:SER:O	1:A:266:LEU:HB2	2.20	0.41
1:B:170:MET:HE1	1:B:350:MET:HB2	2.01	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	357/386 (92%)	344 (96%)	12 (3%)	1 (0%)	36	29
1	B	350/386 (91%)	337 (96%)	12 (3%)	1 (0%)	36	29
All	All	707/772 (92%)	681 (96%)	24 (3%)	2 (0%)	36	29

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	266	LEU
1	B	341	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	309/319 (97%)	300 (97%)	9 (3%)	37	31
1	B	302/319 (95%)	292 (97%)	10 (3%)	33	26
All	All	611/638 (96%)	592 (97%)	19 (3%)	35	29

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	86	LYS
1	A	132	ASN
1	A	183	ASN
1	A	198	LYS
1	A	252	LYS
1	A	270	PHE
1	A	278	SER
1	A	305	SER
1	A	391	VAL
1	B	65	ASP
1	B	71	GLU
1	B	98	GLU
1	B	109	THR
1	B	198	LYS
1	B	219	LEU
1	B	225	SER
1	B	271	SER
1	B	322	MET
1	B	333	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	89	GLN
1	A	183	ASN
1	A	320	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	251	GLN
1	B	320	GLN
1	B	361	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	B3P	A	1000	-	18,18,18	0.59	0	23,23,23	2.14	5 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	B3P	A	1000	-	-	14/28/28/28	-

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1000	B3P	C2-N2-C8	6.54	125.73	116.17
2	A	1000	B3P	C3-N1-C4	5.75	124.58	116.17
2	A	1000	B3P	C11-C8-C10	2.65	115.84	110.02
2	A	1000	B3P	O3-C11-C8	-2.28	107.05	111.68
2	A	1000	B3P	C9-C8-N2	2.24	115.69	109.02

There are no chirality outliers.


All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1000	B3P	N1-C4-C5-O4
2	A	1000	B3P	C6-C4-C5-O4
2	A	1000	B3P	C7-C4-C5-O4
2	A	1000	B3P	C9-C8-N2-C2
2	A	1000	B3P	C10-C8-N2-C2
2	A	1000	B3P	C11-C8-N2-C2
2	A	1000	B3P	O2-C10-C8-N2
2	A	1000	B3P	O2-C10-C8-C9
2	A	1000	B3P	O2-C10-C8-C11
2	A	1000	B3P	C3-C1-C2-N2
2	A	1000	B3P	O3-C11-C8-C9
2	A	1000	B3P	C1-C3-N1-C4
2	A	1000	B3P	O3-C11-C8-C10
2	A	1000	B3P	O3-C11-C8-N2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	364/386 (94%)	0.41	23 (6%) 26 27	6, 18, 37, 60	14 (3%)
1	B	359/386 (93%)	1.47	90 (25%) 1 1	11, 32, 53, 69	16 (4%)
All	All	723/772 (93%)	0.94	113 (15%) 5 5	6, 26, 49, 69	30 (4%)

All (113) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	288	THR	8.0
1	B	265	SER	7.4
1	B	235	ALA	6.0
1	B	17	ALA	5.9
1	A	323	MET	5.9
1	B	292	TYR	5.7
1	B	283	TRP	5.6
1	B	290	TRP	5.4
1	B	340	THR	5.1
1	B	111	PHE	4.6
1	A	292	TYR	4.5
1	A	270	PHE	4.5
1	A	267	ILE	4.5
1	B	16	LEU	4.4
1	B	375	ALA	4.2
1	A	16	LEU	4.2
1	B	183	ASN	4.1
1	A	15	PHE	4.0
1	B	224	GLN	3.9
1	B	374	ILE	3.9
1	B	391	VAL	3.9
1	B	161	ASN	3.8
1	B	108	ASN	3.8
1	B	263	ARG	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	266	LEU	3.8
1	B	392	ALA	3.8
1	B	372	ALA	3.7
1	B	261	VAL	3.7
1	A	283	TRP	3.7
1	B	306	SER	3.6
1	B	129	ILE	3.6
1	B	305	SER	3.6
1	B	262	ALA	3.5
1	B	107	PHE	3.5
1	B	377	ALA	3.5
1	B	230	CYS	3.5
1	B	291	SER	3.4
1	B	106	GLY	3.4
1	A	375	ALA	3.4
1	B	323	MET	3.4
1	B	229	ASP	3.3
1	A	183	ASN	3.3
1	B	105	LYS	3.1
1	B	228	LEU	3.1
1	B	388	THR	3.1
1	B	378	ALA	3.1
1	B	293	PHE	3.1
1	B	88	THR	3.1
1	B	109	THR	3.1
1	B	39	PRO	3.1
1	B	223	GLY	3.1
1	B	264	ALA	3.1
1	B	227	ASN	3.0
1	B	294	PRO	3.0
1	A	107	PHE	3.0
1	A	14	ASN	3.0
1	A	272	ASP	2.9
1	B	389	GLU	2.9
1	B	272	ASP	2.9
1	B	233	TYR	2.8
1	B	162	ILE	2.8
1	A	17	ALA	2.8
1	B	376	GLY	2.8
1	B	289	PRO	2.8
1	B	379	VAL	2.8
1	B	324	GLY	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	322	MET	2.7
1	B	304	ASN	2.7
1	B	370	PRO	2.7
1	B	341	ASN	2.7
1	B	332	TYR	2.7
1	B	236	ASP	2.6
1	B	315	PRO	2.6
1	B	221	ILE	2.6
1	B	260	ALA	2.6
1	B	281	ALA	2.6
1	B	353	PHE	2.6
1	B	280	LEU	2.6
1	B	396	VAL	2.6
1	A	330	GLU	2.5
1	B	259	GLU	2.5
1	B	157	VAL	2.5
1	A	397	PRO	2.5
1	B	390	ASP	2.5
1	B	173	ALA	2.5
1	B	256	ALA	2.5
1	B	314	LEU	2.4
1	B	330	GLU	2.4
1	B	18	MET	2.4
1	A	106	GLY	2.4
1	B	225	SER	2.4
1	B	77	ARG	2.3
1	B	254	PHE	2.3
1	B	226	LEU	2.3
1	A	288	THR	2.3
1	B	276	THR	2.3
1	B	190	GLY	2.3
1	B	78	SER	2.3
1	B	331	CYS	2.3
1	A	389	GLU	2.2
1	B	40	PRO	2.2
1	B	271	SER	2.2
1	B	222	GLY	2.2
1	B	339	SER	2.1
1	B	393	SER	2.1
1	A	271	SER	2.1
1	A	18	MET	2.1
1	B	321	PRO	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	264	ALA	2.1
1	B	258	VAL	2.1
1	B	126	LEU	2.1
1	B	386	PHE	2.0
1	A	108	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	B3P	A	1000	19/19	0.78	0.16	27,37,46,47	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.