



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:41 PM UTC

PDB ID : 3ZKJ / pdb_00003zkj
Title : Crystal Structure of Ankyrin Repeat and Socs Box-Containing Protein 9 (Asb9) in Complex with Elonginb and Elonginc
Authors : Muniz, J.R.C.; Guo, K.; Zhang, Y.; Ayinampudi, V.; Savitsky, P.; Keates, T.; Filippakopoulos, P.; Vollmar, M.; Yue, W.W.; Krojer, T.; Ugochukwu, E.; von Delft, F.; Knapp, S.; Weigelt, J.; Arrowsmith, C.H.; Edwards, A.M.; Bountra, C.; Bullock, A.N.
Deposited on : 2013-01-23
Resolution : 2.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4770 (2.60-2.56)
Clashscore	190562	5124 (2.60-2.56)
Ramachandran outliers	187476	5046 (2.60-2.56)
Sidechain outliers	187428	5046 (2.60-2.56)
RSRZ outliers	180081	4770 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	261	3% 85% 13% .
1	D	261	2% 85% 13% ..
2	B	96	15% 79% 6% 15%
2	E	96	8% 77% 7% 16%

Continued on next page...

Validation Pipeline (wwPDB-VP) : 2.49

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	118	21% 53% 7% 41%
3	F	118	8% 72% 16% • 11%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6472 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ANKYRIN REPEAT AND SOCS BOX PROTEIN 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	257	Total	C	N	O	S	0	1	0
			1921	1212	347	349	13			
1	D	259	Total	C	N	O	S	0	1	0
			1952	1230	354	355	13			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	34	MET	-	expression tag	UNP Q96DX5
D	34	MET	-	expression tag	UNP Q96DX5

- Molecule 2 is a protein called TRANSCRIPTION ELONGATION FACTOR B POLYPEPTIDE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	82	Total	C	N	O	S	0	0	0
			612	399	95	114	4			
2	E	81	Total	C	N	O	S	0	0	0
			628	413	98	112	5			

- Molecule 3 is a protein called TRANSCRIPTION ELONGATION FACTOR B POLYPEPTIDE 2.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	70	Total	C	N	O	S	0	0	0
			475	302	77	94	2			
3	F	105	Total	C	N	O	S	0	0	0
			805	511	136	154	4			

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Cl	0	0
			1	1		
6	F	1	Total	Cl	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	35	Total	O	0	0
			35	35		
7	B	1	Total	O	0	0
			1	1		
7	D	21	Total	O	0	0
			21	21		
7	F	5	Total	O	0	0
			5	5		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ANKYRIN REPEAT AND SOCS BOX PROTEIN 9




- Molecule 1: ANKYRIN REPEAT AND SOCS BOX PROTEIN 9



- Molecule 2: TRANSCRIPTION ELONGATION FACTOR B POLYPEPTIDE 1




- Molecule 2: TRANSCRIPTION ELONGATION FACTOR B POLYPEPTIDE 1




- Molecule 3: TRANSCRIPTION ELONGATION FACTOR B POLYPEPTIDE 2





● Molecule 3: TRANSCRIPTION ELONGATION FACTOR B POLYPEPTIDE 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 2 21	Depositor
Cell constants a, b, c, α , β , γ	102.86Å 110.74Å 113.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.47 – 2.58 50.47 – 2.58	Depositor EDS
% Data completeness (in resolution range)	99.4 (50.47-2.58) 99.4 (50.47-2.58)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 2.58Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.227 , 0.265 0.226 , 0.259	Depositor DCC
R_{free} test set	2058 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	60.9	Xtriage
Anisotropy	0.527	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 55.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.023 for -h,l,k	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6472	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, EDO, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.09	1/1966 (0.1%)	1.06	5/2673 (0.2%)
1	D	0.97	0/1996	1.01	5/2715 (0.2%)
2	B	0.89	0/624	0.99	1/846 (0.1%)
2	E	0.86	0/641	0.96	0/865
3	C	0.81	0/481	1.02	0/656
3	F	0.97	0/822	1.10	0/1116
All	All	0.98	1/6530 (0.0%)	1.03	11/8871 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	276	HIS	C-O	-5.05	1.17	1.24

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	243	SER	CA-C-N	8.71	128.35	119.82
1	A	243	SER	C-N-CA	8.71	128.35	119.82
1	A	147	VAL	CB-CA-C	-6.25	103.97	111.97
2	B	95	ILE	CB-CA-C	-5.91	103.67	110.41
1	D	128	VAL	N-CA-C	5.63	115.82	110.53
1	D	147	VAL	CB-CA-C	-5.51	104.70	112.14
1	D	243	SER	CA-C-N	5.10	126.22	119.84
1	D	243	SER	C-N-CA	5.10	126.22	119.84
1	A	147	VAL	N-CA-C	5.04	115.26	110.42
1	D	54	ASN	N-CA-C	5.04	116.47	111.07
1	A	243	SER	N-CA-C	5.03	116.17	109.93

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1921	0	1882	31	0
1	D	1952	0	1919	22	0
2	B	612	0	564	3	0
2	E	628	0	617	5	0
3	C	475	0	408	4	0
3	F	805	0	779	11	0
4	A	4	0	6	0	0
4	D	4	0	6	0	0
5	A	7	0	10	0	0
6	A	1	0	0	1	0
6	F	1	0	0	0	0
7	A	35	0	0	2	0
7	B	1	0	0	0	0
7	D	21	0	0	0	0
7	F	5	0	0	0	0
All	All	6472	0	6191	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:260:GLN:HE21	1:A:260:GLN:HA	1.42	0.83
2:E:20:LYS:HE3	2:E:28:GLU:CB	2.14	0.77
1:A:236:VAL:O	1:A:239:VAL:HG12	1.89	0.72
2:E:81:VAL:O	2:E:84:THR:HG22	1.93	0.69
3:F:30:ILE:O	3:F:34:ILE:HD13	1.91	0.69
3:F:99:LEU:HD12	3:F:100:PRO:HD2	1.77	0.67
1:D:209:THR:HG22	1:D:211:SER:N	2.10	0.66
3:F:33:GLY:C	3:F:34:ILE:HD12	2.21	0.66
1:A:209:THR:HG22	1:A:211:SER:N	2.11	0.66
1:D:209:THR:HG22	1:D:211:SER:H	1.62	0.64
1:A:209:THR:HG22	1:A:211:SER:H	1.64	0.63
1:D:37:TRP:HZ2	1:D:45:ILE:HD12	1.64	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:260:GLN:HA	1:A:260:GLN:NE2	2.14	0.60
1:A:115:TRP:HB2	1:A:148[A]:GLU:HG3	1.86	0.56
1:A:260:GLN:HE21	1:A:260:GLN:CA	2.13	0.53
1:D:130:PRO:HD2	1:D:137:PRO:HD2	1.90	0.53
1:A:212:GLU:N	6:A:1297:CL:CL	2.79	0.52
1:D:286:ASP:OD1	1:D:287:LEU:HG	2.10	0.52
1:D:40:MET:HE1	1:D:56:ILE:CG1	2.39	0.51
1:A:40:MET:HE1	1:A:56:ILE:HG13	1.91	0.51
1:A:115:TRP:HB2	1:A:148[A]:GLU:CG	2.41	0.51
1:D:209:THR:CG2	1:D:211:SER:HB2	2.41	0.50
1:D:40:MET:HE1	1:D:56:ILE:HG13	1.94	0.49
1:A:40:MET:HE1	1:A:56:ILE:CG1	2.42	0.49
1:A:50:LEU:HB2	1:D:212:GLU:OE1	2.13	0.49
3:F:34:ILE:HG22	3:F:35:LEU:N	2.27	0.49
1:D:37:TRP:CZ2	1:D:45:ILE:HD12	2.46	0.49
1:A:209:THR:CG2	1:A:211:SER:HB2	2.43	0.48
1:D:219:MET:HE1	1:D:252:ARG:HG3	1.96	0.48
3:F:61:GLY:O	3:F:66:THR:HG21	2.13	0.47
1:A:246:ALA:O	1:A:247:GLN:C	2.57	0.47
1:D:255:PRO:HG2	2:E:80:LYS:HD2	1.96	0.47
1:A:128:VAL:HG11	1:A:158:GLY:CA	2.44	0.46
1:A:130:PRO:HD2	1:A:137:PRO:HD2	1.96	0.46
3:F:99:LEU:HD12	3:F:100:PRO:CD	2.45	0.46
1:A:239:VAL:HG21	1:A:245:LEU:HB2	1.97	0.46
1:A:260:GLN:HE22	1:A:263:ARG:HE	1.64	0.45
3:F:38:PRO:HD2	3:F:41:GLU:OE1	2.16	0.45
1:A:115:TRP:NE1	1:A:119:ASN:ND2	2.65	0.45
1:D:37:TRP:CG	1:D:38:SER:H	2.35	0.44
1:A:115:TRP:NE1	1:A:119:ASN:HD21	2.16	0.44
1:D:115:TRP:HB2	1:D:148[B]:GLU:HG3	2.00	0.44
1:D:259:MET:HG2	2:E:104:LEU:HD12	1.99	0.44
1:A:145:GLY:HA2	1:A:182:CYS:SG	2.58	0.43
3:F:23:THR:HA	3:F:56:THR:HA	2.00	0.43
3:C:3:VAL:O	3:C:17:ASP:HA	2.19	0.43
1:A:293:HIS:HE1	7:A:2025:HOH:O	2.02	0.42
3:F:11:LYS:HG3	3:F:91:GLU:HG3	2.02	0.42
1:A:239:VAL:HG21	1:A:245:LEU:CB	2.49	0.42
3:C:63:THR:HG22	3:C:64:SER:N	2.35	0.42
1:A:83:SER:HB2	7:A:2005:HOH:O	2.19	0.42
2:B:78:THR:HG22	3:C:69:PRO:HB3	2.02	0.42
1:A:260:GLN:HE22	1:A:263:ARG:HH21	1.67	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:236:VAL:HA	1:A:249:PHE:HD2	1.85	0.41
3:F:79:PHE:O	3:F:86:GLU:HG2	2.20	0.41
1:A:265:ARG:NH2	2:B:97:PRO:HG3	2.35	0.41
1:D:161:ASP:O	1:D:162:HIS:C	2.64	0.41
2:E:20:LYS:CE	2:E:28:GLU:CB	2.93	0.41
1:A:255:PRO:HA	1:A:256:PRO:HD3	1.89	0.41
1:A:239:VAL:O	1:A:240:PRO:C	2.63	0.41
3:C:10:HIS:O	3:C:91:GLU:HB2	2.21	0.41
1:D:99:THR:HG23	1:D:103:HIS:O	2.21	0.41
1:D:128:VAL:HG11	1:D:158:GLY:CA	2.51	0.41
1:D:196:GLY:CA	1:D:202:PRO:HD3	2.51	0.41
1:A:236:VAL:O	1:A:239:VAL:CG1	2.62	0.40
1:D:265:ARG:HA	1:D:265:ARG:HD2	1.93	0.40
1:A:236:VAL:O	1:A:236:VAL:HG22	2.21	0.40
1:D:219:MET:O	1:D:220:ASP:C	2.62	0.40
2:B:96:ALA:O	2:B:99:ILE:HG13	2.21	0.40
3:F:3:VAL:HG13	3:F:67:ALA:HB3	2.02	0.40
1:D:286:ASP:OD1	1:D:287:LEU:N	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	254/261 (97%)	249 (98%)	5 (2%)	0	100	100
1	D	258/261 (99%)	249 (96%)	9 (4%)	0	100	100
2	B	76/96 (79%)	74 (97%)	2 (3%)	0	100	100
2	E	75/96 (78%)	74 (99%)	1 (1%)	0	100	100
3	C	60/118 (51%)	54 (90%)	6 (10%)	0	100	100
3	F	103/118 (87%)	94 (91%)	9 (9%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	826/950 (87%)	794 (96%)	32 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/218 (93%)	197 (98%)	5 (2%)	42	67
1	D	206/218 (94%)	200 (97%)	6 (3%)	37	63
2	B	60/85 (71%)	59 (98%)	1 (2%)	53	76
2	E	66/85 (78%)	65 (98%)	1 (2%)	57	78
3	C	43/103 (42%)	42 (98%)	1 (2%)	44	69
3	F	85/103 (82%)	82 (96%)	3 (4%)	32	57
All	All	662/812 (82%)	645 (97%)	17 (3%)	40	66

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	62	VAL
1	A	95	VAL
1	A	134	LEU
1	A	209	THR
1	A	242	GLU
2	B	31	VAL
3	C	16	THR
1	D	62	VAL
1	D	95	VAL
1	D	129	GLN
1	D	134	LEU
1	D	209	THR
1	D	233	LYS
2	E	31	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	F	12	THR
3	F	66	THR
3	F	94	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	119	ASN
1	A	260	GLN
2	B	61	ASN
1	D	48	HIS
1	D	119	ASN
1	D	129	GLN
2	E	61	ASN
2	E	108	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	D	1295	-	3,3,3	0.51	0	2,2,2	0.30	0
4	EDO	A	1295	-	3,3,3	0.57	0	2,2,2	0.14	0
5	PEG	A	1296	-	6,6,6	0.79	0	5,5,5	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	D	1295	-	-	1/1/1/1	-
4	EDO	A	1295	-	-	0/1/1/1	-
5	PEG	A	1296	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1296	PEG	O1-C1-C2-O2
5	A	1296	PEG	O2-C3-C4-O4
5	A	1296	PEG	C4-C3-O2-C2
4	D	1295	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	257/261 (98%)	0.05	8 (3%)	51	46	29, 55, 91, 124	2 (0%)
1	D	259/261 (99%)	0.20	6 (2%)	61	56	30, 65, 98, 129	2 (0%)
2	B	82/96 (85%)	0.94	14 (17%)	4	3	44, 88, 129, 139	0
2	E	81/96 (84%)	0.83	8 (9%)	13	10	67, 92, 128, 134	0
3	C	70/118 (59%)	1.73	25 (35%)	1	0	59, 123, 182, 211	0
3	F	105/118 (88%)	0.58	9 (8%)	16	13	58, 84, 129, 154	0
All	All	854/950 (89%)	0.46	70 (8%)	17	14	29, 70, 129, 211	4 (0%)

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	86	SER	8.0
3	F	77	LEU	4.8
3	C	25	PHE	4.8
3	C	24	VAL	4.7
1	D	36	ASP	4.7
2	B	18	TYR	4.5
3	C	65	GLN	4.5
3	F	78	ALA	4.4
1	A	130	PRO	4.3
2	B	58	ASN	4.3
1	A	129	GLN	4.1
3	C	62	PHE	3.9
3	C	18	ALA	3.9
3	C	16	THR	3.9
1	D	37	TRP	3.8
3	C	22	SER	3.8
2	B	26	GLY	3.7
2	E	89	GLU	3.7
3	F	106	GLN	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	131	GLU	3.4
3	C	12	THR	3.4
1	A	131	GLU	3.3
2	B	86	SER	3.3
3	C	31	VAL	3.3
2	B	89	GLU	3.3
3	C	8	ARG	3.2
1	A	250	LEU	3.1
3	C	19	LYS	3.1
3	C	50	LEU	3.0
3	C	10	HIS	3.0
3	C	63	THR	3.0
3	C	35	LEU	3.0
3	C	9	ARG	2.9
1	D	130	PRO	2.9
3	F	83	ASP	2.9
3	C	66	THR	2.8
3	C	48	ASP	2.8
3	C	3	VAL	2.8
2	B	27	HIS	2.7
2	E	26	GLY	2.7
2	B	83	TYR	2.7
2	E	84	THR	2.7
2	B	84	THR	2.6
3	F	81	ALA	2.6
2	B	23	SER	2.6
2	E	18	TYR	2.5
2	B	31	VAL	2.5
3	F	2	ASP	2.4
3	F	79	PHE	2.4
2	B	22	ILE	2.3
2	E	27	HIS	2.3
3	F	10	HIS	2.3
2	E	76	TYR	2.3
3	C	64	SER	2.3
3	C	46	LYS	2.3
1	A	58	GLN	2.2
1	A	241	PRO	2.2
3	C	39	PRO	2.2
1	A	128	VAL	2.1
2	B	64	GLU	2.1
1	D	241	PRO	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	45	MET	2.1
1	A	239	VAL	2.1
2	B	30	ILE	2.1
1	D	247	GLN	2.1
3	F	12	THR	2.1
3	C	68	ARG	2.0
2	E	28	GLU	2.0
3	C	17	ASP	2.0
3	C	11	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	PEG	A	1296	7/7	0.80	0.24	76,101,113,114	0
6	CL	F	1107	1/1	0.90	0.21	93,93,93,93	0
4	EDO	D	1295	4/4	0.91	0.13	87,93,93,97	0
6	CL	A	1297	1/1	0.95	0.21	97,97,97,97	0
4	EDO	A	1295	4/4	0.95	0.12	68,71,73,74	0

6.5 Other polymers [i](#)

There are no such residues in this entry.