



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:50 AM UTC

PDB ID : 2ZL7 / pdb_00002zl7
Title : Atomic resolution structural characterization of recognition of histo-blood group antigens by Norwalk virus
Authors : Choi, J.M.; Huston, A.M.; Estes, M.K.; Prasad, B.V.V.
Deposited on : 2008-04-02
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1216 (1.36-1.36)
Clashscore	190562	1232 (1.36-1.36)
Ramachandran outliers	187476	1220 (1.36-1.36)
Sidechain outliers	187428	1220 (1.36-1.36)
RSRZ outliers	180081	1214 (1.36-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	295	8% 89% 7% .
1	B	295	14% 88% 8% ..
2	C	3	33% 67%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NGA	C	3	X	-	-	-
5	ACT	A	520	-	-	X	-
5	ACT	B	521	-	-	X	-

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 5106 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 58 kd capsid protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	285	Total	C	N	O	S	0	1	0
			2154	1380	357	409	8			
1	B	286	Total	C	N	O	S	0	3	0
			2169	1390	359	412	8			

- Molecule 2 is an oligosaccharide called alpha-L-fucopyranose-(1-2)-[2-acetamido-2-deoxy-beta-D-galactopyranose-(1-3)]beta-D-galactopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	3	Total	C	N	O	0	0	0
			36	20	1	15			

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	B	1	Total	Mg	0	0
			1	1		

- Molecule 5 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	347	Total	O	0	0
			347	347		
6	B	389	Total	O	0	0
			389	389		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 58 kd capsid protein



- Molecule 1: 58 kd capsid protein



- Molecule 2: alpha-L-fucopyranose-(1-2)-[2-acetamido-2-deoxy-beta-D-galactopyranose-(1-3)]bet a-D-galactopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	83.07 Å 83.07 Å 164.32 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.07 – 1.35 30.07 – 1.36	Depositor EDS
% Data completeness (in resolution range)	98.5 (30.07-1.35) 98.5 (30.07-1.36)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.84 (at 1.36 Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.171 , 0.188 0.178 , 0.195	Depositor DCC
R_{free} test set	6957 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	13.1	Xtriage
Anisotropy	0.007	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 45.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.036 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5106	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, GAL, CA, FUC, MG, NGA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	0/2227	0.74	0/3054
1	B	0.63	0/2249	0.81	3/3084 (0.1%)
All	All	0.62	0/4476	0.78	3/6138 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	383	SER	N-CA-C	-13.25	88.02	109.76
1	B	348	THR	CA-C-N	-5.17	115.40	120.52
1	B	348	THR	C-N-CA	-5.17	115.40	120.52

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	382	PRO	Peptide
1	B	413	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2154	0	2075	30	0
1	B	2169	0	2099	34	0
2	C	36	0	33	0	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	4	0	3	13	0
5	B	4	0	3	15	0
6	A	347	0	0	8	0
6	B	389	0	0	4	0
All	All	5106	0	4213	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:304:LEU:HD11	5:A:520:ACT:H3	1.25	1.17
1:B:304:LEU:HD11	5:B:521:ACT:H3	1.20	1.14
1:B:304:LEU:HD11	5:B:521:ACT:CH3	1.95	0.94
1:B:304:LEU:CD1	5:B:521:ACT:H3	1.97	0.94
1:A:370:VAL:CG1	6:A:690:HOH:O	2.17	0.91
1:A:304:LEU:CD1	5:A:520:ACT:H3	2.00	0.90
1:A:304:LEU:HD21	5:A:520:ACT:CH3	2.02	0.90
1:A:370:VAL:HG12	6:A:690:HOH:O	1.74	0.86
1:B:414:PHE:O	1:B:414:PHE:HD2	1.59	0.85
1:A:304:LEU:HD11	5:A:520:ACT:CH3	2.09	0.78
1:B:291:ARG:HD2	1:B:370:VAL:HG22	1.66	0.77
1:A:396:GLY:O	6:A:862:HOH:O	2.03	0.76
1:B:304:LEU:HD13	6:B:959:HOH:O	1.88	0.73
1:B:414:PHE:O	1:B:414:PHE:CD2	2.42	0.72
6:A:690:HOH:O	1:B:426:PRO:HG3	1.90	0.70
1:B:304:LEU:HD21	5:B:521:ACT:CH3	2.22	0.70
1:B:329:HIS:HD2	1:B:344:ASP:OD1	1.77	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:291:ARG:CD	1:B:370:VAL:HG22	2.25	0.67
1:A:304:LEU:HD21	5:A:520:ACT:H2	1.76	0.67
1:B:329:HIS:HE1	1:B:377:SER:OG	1.79	0.66
1:A:329:HIS:HD2	1:A:344:ASP:OD1	1.79	0.66
6:A:521:HOH:O	1:B:286:HIS:HE1	1.79	0.64
1:B:413:GLY:HA3	1:B:414:PHE:CD1	2.32	0.64
1:A:286:HIS:HE1	6:B:663:HOH:O	1.79	0.64
1:B:289:LYS:HE2	5:B:521:ACT:H1	1.79	0.63
1:A:370:VAL:HG11	6:A:690:HOH:O	1.93	0.63
1:A:329:HIS:HE1	1:A:377:SER:OG	1.82	0.61
1:B:396:GLY:O	6:B:1036:HOH:O	2.16	0.59
1:A:304:LEU:HD21	5:A:520:ACT:H1	1.83	0.59
1:A:304:LEU:CD2	5:A:520:ACT:CH3	2.79	0.58
1:A:304:LEU:CG	5:A:520:ACT:H3	2.34	0.58
1:A:289:LYS:HE2	5:A:520:ACT:H1	1.84	0.58
1:B:286:HIS:HD2	6:B:662:HOH:O	1.87	0.57
1:A:286:HIS:HD2	6:A:537:HOH:O	1.87	0.56
1:A:314:GLY:O	1:A:355:HIS:HD2	1.87	0.56
1:A:399:ILE:H	1:A:399:ILE:HD12	1.70	0.55
1:A:289:LYS:CE	5:A:520:ACT:H1	2.37	0.53
1:A:399:ILE:HD12	1:A:399:ILE:N	2.23	0.53
1:B:399:ILE:HD12	1:B:399:ILE:C	2.34	0.53
1:B:289:LYS:CE	5:B:521:ACT:H1	2.38	0.52
1:B:304:LEU:HD21	5:B:521:ACT:H2	1.92	0.51
1:B:304:LEU:CD1	5:B:521:ACT:CH3	2.75	0.50
1:B:414:PHE:H	1:B:415:GLY:HA2	1.77	0.50
1:A:239:SER:HB3	1:A:496:ILE:HG23	1.93	0.50
1:A:289:LYS:NZ	5:A:520:ACT:H1	2.27	0.49
1:B:239:SER:HB3	1:B:497:ILE:HG23	1.95	0.49
6:A:720:HOH:O	1:B:444:HIS:HD2	1.95	0.49
1:A:346:ASP:OD2	1:A:381:HIS:HD2	1.95	0.49
1:B:413:GLY:O	1:B:416:GLU:HB2	2.13	0.49
1:A:304:LEU:CG	5:A:520:ACT:CH3	2.91	0.48
1:A:331:ASN:HD21	1:A:340:GLN:NE2	2.11	0.48
1:B:304:LEU:CG	5:B:521:ACT:H3	2.45	0.46
1:B:304:LEU:HD21	5:B:521:ACT:H1	1.97	0.45
1:B:304:LEU:CG	5:B:521:ACT:CH3	2.95	0.45
1:A:249:PRO:HG2	5:B:521:ACT:H2	1.98	0.44
1:B:304:LEU:CD2	5:B:521:ACT:CH3	2.95	0.43
1:A:474:LYS:HG3	1:A:512:LEU:HD21	2.01	0.42
1:B:232:THR:OG1	1:B:444:HIS:HE1	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:399:ILE:HD12	1:B:400:THR:N	2.35	0.42
1:B:289:LYS:NZ	5:B:521:ACT:H1	2.35	0.41
1:A:304:LEU:CD1	5:A:520:ACT:CH3	2.84	0.41
1:B:291:ARG:HD2	1:B:370:VAL:CG2	2.43	0.41
1:A:249:PRO:CG	5:B:521:ACT:H2	2.51	0.41
1:B:346:ASP:OD2	1:B:381:HIS:HD2	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	282/295 (96%)	276 (98%)	5 (2%)	1 (0%)	30	11
1	B	287/295 (97%)	280 (98%)	6 (2%)	1 (0%)	36	16
All	All	569/590 (96%)	556 (98%)	11 (2%)	2 (0%)	30	11

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	414	PHE
1	A	339	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	243/250 (97%)	242 (100%)	1 (0%)	84	69
1	B	246/250 (98%)	245 (100%)	1 (0%)	84	69
All	All	489/500 (98%)	487 (100%)	2 (0%)	84	69

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	432	ASN
1	B	452	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	286	HIS
1	A	329	HIS
1	A	340	GLN
1	A	355	HIS
1	A	381	HIS
1	A	444	HIS
1	A	449	GLN
1	B	286	HIS
1	B	329	HIS
1	B	334	GLN
1	B	381	HIS
1	B	394	ASN
1	B	439	GLN
1	B	444	HIS
1	B	493	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GAL	C	1	2	12,12,12	0.51	0	17,17,17	0.94	1 (5%)
2	FUC	C	2	2	10,10,11	0.40	0	14,14,16	0.34	0
2	NGA	C	3	2	14,14,15	2.39	2 (14%)	17,19,21	2.13	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GAL	C	1	2	-	0/2/22/22	0/1/1/1
2	FUC	C	2	2	-	-	0/1/1/1
2	NGA	C	3	2	1/1/5/7	2/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	3	NGA	O7-C7	8.38	1.42	1.23
2	C	3	NGA	C7-N2	2.75	1.43	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	3	NGA	O7-C7-N2	-6.40	110.66	121.98
2	C	3	NGA	O7-C7-C8	-4.07	114.80	122.05
2	C	1	GAL	O3-C3-C2	-2.54	104.39	110.38
2	C	3	NGA	C8-C7-N2	-2.50	111.98	116.12

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	C	3	NGA	C1


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	3	NGA	C1-C2-N2-C7
2	C	3	NGA	O7-C7-N2-C2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	ACT	A	520	-	3,3,3	0.70	0	3,3,3	1.05	0
5	ACT	B	521	-	3,3,3	0.76	0	3,3,3	0.97	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 28 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	520	ACT	13	0
5	B	521	ACT	15	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	285/295 (96%)	0.42	24 (8%)	17 23	7, 13, 27, 34	1 (0%)
1	B	286/295 (96%)	0.49	41 (14%)	6 10	6, 13, 29, 38	3 (1%)
All	All	571/590 (96%)	0.45	65 (11%)	10 14	6, 13, 28, 38	4 (0%)

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	414	PHE	7.0
1	A	414	PHE	6.7
1	B	488	ALA	5.7
1	B	399	ILE	5.7
1	A	400	THR	5.6
1	A	509	PHE	5.6
1	A	399	ILE	5.6
1	B	413	GLY	5.5
1	B	452	THR	5.3
1	B	510	PHE	5.3
1	A	452	THR	5.1
1	B	400	THR	4.5
1	A	515	VAL	4.2
1	B	487	GLY	4.1
1	B	486	ASN	4.0
1	B	489	SER	4.0
1	B	516	VAL	3.9
1	A	453	VAL	3.9
1	B	383	SER	3.8
1	B	514	LYS	3.8
1	B	451	PRO	3.7
1	B	485	PRO	3.7
1	B	415	GLY	3.6
1	A	487	GLY	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	397	SER	3.5
1	B	453	VAL	3.5
1	A	465	ASP	3.4
1	B	490	SER	3.4
1	A	451	PRO	3.2
1	B	467	GLY	3.2
1	A	397	SER	3.1
1	B	491	GLY	3.1
1	A	488	ALA	3.0
1	B	450	ALA	3.0
1	A	486	ASN	3.0
1	B	410	TYR	2.9
1	A	464	PRO	2.8
1	B	454	GLY	2.8
1	B	511	TYR	2.8
1	B	509	ARG	2.8
1	A	489	SER	2.8
1	B	464	PRO	2.8
1	B	384	GLY	2.7
1	B	484	VAL	2.7
1	B	494	GLN	2.6
1	A	466	THR	2.6
1	B	515	PRO	2.6
1	B	417	VAL	2.6
1	B	465	ASP	2.5
1	A	259	ASN	2.5
1	A	511	GLN	2.4
1	B	304	LEU	2.4
1	A	398	SER	2.3
1	B	506	TRP	2.3
1	A	296	GLY	2.3
1	A	261	GLN	2.2
1	B	513	LEU	2.2
1	B	398	SER	2.2
1	A	304	LEU	2.1
1	B	468	ARG	2.1
1	B	493	GLN	2.1
1	A	350	ASP	2.1
1	A	484	VAL	2.1
1	B	508	SER	2.0
1	B	466	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GAL	C	1	12/12	0.93	0.08	13,16,18,19	0
2	FUC	C	2	10/11	0.94	0.07	13,15,17,17	0
2	NGA	C	3	14/15	0.95	0.07	11,12,15,16	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CA	A	1	1/1	0.93	0.28	42,42,42,42	0
5	ACT	A	520	4/4	0.93	0.14	17,18,18,19	0
5	ACT	B	521	4/4	0.94	0.11	17,17,17,18	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MG	A	519	1/1	0.98	0.07	18,18,18,18	0
4	MG	B	520	1/1	0.98	0.12	19,19,19,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.