



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 08:15 AM UTC

PDB ID : 4ZLG / pdb_00004zlg
Title : Cellobionic acid phosphorylase - gluconic acid complex
Authors : Nam, Y.W.; Arakawa, T.; Fushinobu, S.
Deposited on : 2015-05-01
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	796	5% 87% 10% ..

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6964 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Putative b-glycan phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	785	6228	3965	1060	1171	32	0	0	0

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	MET	engineered mutation	UNP Q21MB1
A	789	LEU	-	expression tag	UNP Q21MB1
A	790	GLU	-	expression tag	UNP Q21MB1
A	791	HIS	-	expression tag	UNP Q21MB1
A	792	HIS	-	expression tag	UNP Q21MB1
A	793	HIS	-	expression tag	UNP Q21MB1
A	794	HIS	-	expression tag	UNP Q21MB1
A	795	HIS	-	expression tag	UNP Q21MB1
A	796	HIS	-	expression tag	UNP Q21MB1

- Molecule 2 is D-gluconic acid (CCD ID: GCO) (formula: C₆H₁₂O₇).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	6	7		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is D-glucono-1,5-lactone (CCD ID: LGC) (formula: $C_6H_{10}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			12	6	6		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Cl	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	688	Total	O	0	0
			688	688		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative b-glycan phosphorylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	106.99Å 106.99Å 186.54Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	41.66 – 1.75 41.66 – 1.75	Depositor EDS
% Data completeness (in resolution range)	98.0 (41.66-1.75) 98.1 (41.66-1.75)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.70 (at 1.75Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, R_{free}	0.169 , 0.201 0.181 , 0.209	Depositor DCC
R_{free} test set	6113 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	23.1	Xtriage
Anisotropy	0.009	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 37.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.019 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6964	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, LGC, CL, GCO, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.60	42/6393 (0.7%)	1.24	15/8683 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (42) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	349	ARG	CD-NE	-8.23	1.34	1.46
1	A	264	GLU	CA-C	6.97	1.61	1.52
1	A	699	TRP	C-O	6.96	1.32	1.24
1	A	334	PHE	N-CA	-6.91	1.38	1.46
1	A	426	LEU	N-CA	6.78	1.54	1.46
1	A	492	THR	CA-C	6.67	1.61	1.52
1	A	320	GLU	N-CA	6.09	1.53	1.46
1	A	515	PHE	N-CA	6.07	1.54	1.46
1	A	452	TRP	C-O	5.84	1.30	1.24
1	A	386	GLY	CA-C	-5.84	1.47	1.52
1	A	291	ALA	N-CA	5.83	1.53	1.46
1	A	539	GLY	C-O	-5.83	1.17	1.24
1	A	418	LEU	C-O	5.80	1.31	1.24
1	A	493	ALA	N-CA	5.75	1.53	1.46
1	A	675	ARG	CA-C	5.68	1.60	1.52
1	A	678	TYR	N-CA	-5.66	1.39	1.46
1	A	370	PHE	CA-CB	5.65	1.62	1.53
1	A	216	TRP	N-CA	5.61	1.52	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	596	LEU	CA-C	-5.51	1.45	1.52
1	A	127	LEU	CB-CG	-5.42	1.42	1.53
1	A	508	GLN	CD-NE2	5.38	1.44	1.33
1	A	562	GLN	CA-C	5.34	1.59	1.52
1	A	31	MET	SD-CE	-5.25	1.66	1.79
1	A	368	GLN	CA-C	5.24	1.59	1.52
1	A	504	GLU	CA-C	5.22	1.59	1.52
1	A	577	PRO	C-O	5.22	1.30	1.24
1	A	498	LEU	C-O	5.17	1.30	1.24
1	A	550	SER	C-O	-5.16	1.17	1.24
1	A	589	THR	CA-CB	5.15	1.61	1.53
1	A	642	GLU	CA-C	5.15	1.59	1.52
1	A	160	PHE	N-CA	5.15	1.52	1.46
1	A	256	TYR	N-CA	5.09	1.52	1.46
1	A	269	ARG	CA-C	-5.09	1.46	1.52
1	A	207	PHE	N-CA	-5.09	1.40	1.46
1	A	451	ARG	CA-CB	-5.08	1.45	1.53
1	A	564	TRP	N-CA	-5.07	1.39	1.46
1	A	510	SER	CA-C	5.07	1.59	1.52
1	A	602	ALA	CA-C	-5.05	1.46	1.52
1	A	143	LYS	CA-C	-5.05	1.46	1.52
1	A	530	ILE	N-CA	5.03	1.51	1.45
1	A	102	LEU	C-O	-5.02	1.17	1.23
1	A	364	ASN	CG-ND2	5.01	1.43	1.33

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	420	GLU	N-CA-C	8.10	120.18	111.36
1	A	485	SER	CA-C-N	7.63	128.26	119.94
1	A	485	SER	C-N-CA	7.63	128.26	119.94
1	A	431	VAL	N-CA-CB	-7.11	101.26	111.21
1	A	349	ARG	NE-CZ-NH2	-7.00	112.90	119.20
1	A	241	GLY	N-CA-C	6.72	122.06	114.67
1	A	605	ASP	N-CA-C	5.82	120.07	113.15
1	A	361	ILE	CA-C-N	5.79	127.37	122.28
1	A	361	ILE	C-N-CA	5.79	127.37	122.28
1	A	349	ARG	NE-CZ-NH1	5.34	126.84	121.50
1	A	331	ARG	N-CA-CB	-5.33	102.29	110.01
1	A	491	ALA	N-CA-C	5.13	116.96	111.36
1	A	681	HIS	CA-C-N	-5.09	114.42	119.56
1	A	681	HIS	C-N-CA	-5.09	114.42	119.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	47	GLN	N-CA-C	5.00	120.86	109.81

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	349	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6228	0	6003	36	0
2	A	13	0	11	0	0
3	A	12	0	16	0	0
4	A	12	0	10	0	0
5	A	10	0	0	0	0
6	A	1	0	0	0	0
7	A	688	0	0	8	1
All	All	6964	0	6040	36	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:349:ARG:HD3	1:A:405:ASP:OD2	1.39	1.19
1:A:349:ARG:CD	1:A:405:ASP:OD2	1.96	1.13
1:A:349:ARG:HG2	1:A:409:TRP:CD1	2.11	0.85
1:A:349:ARG:HD2	1:A:405:ASP:OD2	1.79	0.82
1:A:736:GLU:CD	7:A:972:HOH:O	2.30	0.74
1:A:353:GLN:HE21	1:A:699:TRP:HE1	1.35	0.73
1:A:32:MET:HE3	1:A:692:PHE:CZ	2.24	0.72
1:A:755:MET:HE3	1:A:757:ILE:HD11	1.74	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:349:ARG:HG3	7:A:1233:HOH:O	1.94	0.67
1:A:62:LYS:HE3	7:A:1461:HOH:O	1.96	0.65
1:A:32:MET:HE3	1:A:692:PHE:HZ	1.63	0.60
1:A:349:ARG:HG2	1:A:409:TRP:NE1	2.14	0.60
1:A:377:GLN:OE1	1:A:445:HIS:HD2	1.84	0.60
1:A:447:HIS:HD2	1:A:499:TRP:HE1	1.49	0.59
1:A:6:ASN:HD22	1:A:10:ARG:HH11	1.51	0.58
1:A:349:ARG:HD3	1:A:405:ASP:CG	2.23	0.58
1:A:64:PHE:CE2	7:A:1459:HOH:O	2.57	0.57
1:A:64:PHE:CD2	7:A:1459:HOH:O	2.53	0.56
1:A:431:VAL:O	1:A:445:HIS:HE1	1.91	0.54
1:A:573:GLU:H	1:A:573:GLU:CD	2.15	0.54
1:A:447:HIS:CD2	1:A:499:TRP:HE1	2.26	0.53
1:A:353:GLN:HE21	1:A:699:TRP:NE1	2.06	0.52
1:A:377:GLN:OE1	1:A:445:HIS:CD2	2.63	0.51
1:A:227:GLY:H	1:A:233:ASN:HD22	1.59	0.50
1:A:86:GLU:HG3	7:A:1519:HOH:O	2.12	0.50
1:A:32:MET:SD	1:A:342:LEU:HD12	2.53	0.49
1:A:6:ASN:ND2	1:A:10:ARG:HH11	2.12	0.48
1:A:32:MET:HE2	1:A:64:PHE:CE2	2.51	0.45
1:A:32:MET:HB2	7:A:1432:HOH:O	2.17	0.44
1:A:438:LYS:HE2	1:A:440:GLU:OE2	2.18	0.43
1:A:36:ASN:ND2	1:A:38:ARG:H	2.17	0.43
1:A:736:GLU:HG3	7:A:972:HOH:O	2.17	0.43
1:A:648:LEU:HD11	1:A:700:VAL:HG13	2.01	0.42
1:A:2:LEU:HA	1:A:12:GLN:O	2.20	0.41
1:A:34:GLN:HE22	1:A:692:PHE:HD2	1.68	0.41
1:A:532:ASP:O	1:A:532:ASP:CG	2.62	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:1278:HOH:O	7:A:1281:HOH:O[6_445]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	783/796 (98%)	762 (97%)	20 (3%)	1 (0%)	48	32

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	95	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	655/665 (98%)	645 (98%)	10 (2%)	57	41

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	GLN
1	A	52	LYS
1	A	361	ILE
1	A	364	ASN
1	A	408	VAL
1	A	438	LYS
1	A	455	GLN
1	A	459	GLU
1	A	750	SER
1	A	753	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	ASN
1	A	34	GLN
1	A	36	ASN
1	A	66	GLN
1	A	103	ASN
1	A	233	ASN
1	A	313	GLN
1	A	340	ASN
1	A	353	GLN
1	A	355	ASN
1	A	368	GLN
1	A	445	HIS
1	A	447	HIS
1	A	529	HIS
1	A	560	ASN
1	A	574	GLN
1	A	586	GLN
1	A	693	ASN
1	A	723	GLN
1	A	729	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	LGC	A	804	-	12,12,12	1.59	2 (16%)	15,17,17	1.37	3 (20%)
5	SO4	A	805	-	4,4,4	0.58	0	6,6,6	0.86	0
3	GOL	A	802	-	5,5,5	0.48	0	5,5,5	0.68	0
2	GCO	A	801	-	12,12,12	1.90	3 (25%)	15,16,16	1.98	4 (26%)
3	GOL	A	803	-	5,5,5	1.09	0	5,5,5	0.46	0
5	SO4	A	806	-	4,4,4	0.45	0	6,6,6	0.88	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	LGC	A	804	-	-	0/2/22/22	0/1/1/1
3	GOL	A	802	-	-	0/4/4/4	-
2	GCO	A	801	-	-	0/18/18/18	-
3	GOL	A	803	-	-	3/4/4/4	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	GCO	C2-C1	-3.57	1.47	1.52
4	A	804	LGC	O5-C1	3.23	1.39	1.34
2	A	801	GCO	O3-C3	3.05	1.50	1.43
4	A	804	LGC	C3-C2	2.98	1.57	1.53
2	A	801	GCO	O1B-C1	-2.31	1.23	1.30

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	GCO	C4-C3-C2	-5.44	103.99	113.62
4	A	804	LGC	O5-C5-C6	-2.94	103.10	106.05
2	A	801	GCO	O3-C3-C2	2.66	114.06	109.29
2	A	801	GCO	O2-C2-C3	-2.50	104.95	110.38
4	A	804	LGC	C6-C5-C4	2.14	118.27	113.02
4	A	804	LGC	O5-C1-C2	2.10	122.30	119.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	GCO	O6-C6-C5	-2.05	106.85	111.16

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	803	GOL	O1-C1-C2-O2
3	A	803	GOL	O2-C2-C3-O3
3	A	803	GOL	O1-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	785/796 (98%)	0.33	38 (4%) 35 41	18, 25, 43, 81	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	785	PRO	4.8
1	A	63	THR	4.0
1	A	64	PHE	3.8
1	A	750	SER	3.8
1	A	66	GLN	3.7
1	A	752	ILE	3.3
1	A	86	GLU	3.3
1	A	764	ILE	3.2
1	A	692	PHE	3.0
1	A	763	VAL	2.9
1	A	543	ASP	2.9
1	A	510	SER	2.8
1	A	65	MET	2.8
1	A	784	LEU	2.7
1	A	303	THR	2.7
1	A	551	LYS	2.6
1	A	749	SER	2.6
1	A	390	LEU	2.5
1	A	134	ASP	2.5
1	A	574	GLN	2.5
1	A	1	GLY	2.5
1	A	437	GLU	2.4
1	A	751	ASP	2.4
1	A	511	CYS	2.4
1	A	783	LEU	2.4
1	A	149	ALA	2.3
1	A	508	GLN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	447	HIS	2.2
1	A	127	LEU	2.2
1	A	199	LYS	2.2
1	A	581	ASP	2.2
1	A	392	GLY	2.2
1	A	299	SER	2.2
1	A	401	ILE	2.1
1	A	451	ARG	2.1
1	A	7	ASN	2.1
1	A	761	GLU	2.1
1	A	438	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GCO	A	801	13/13	0.88	0.16	27,35,54,57	0
3	GOL	A	803	6/6	0.94	0.10	33,41,45,47	0
3	GOL	A	802	6/6	0.97	0.07	26,29,31,35	0
5	SO4	A	806	5/5	0.97	0.08	34,37,39,42	0
4	LGC	A	804	12/12	0.98	0.06	20,21,25,25	0
5	SO4	A	805	5/5	0.99	0.06	24,25,26,30	0
6	CL	A	807	1/1	0.99	0.03	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.