



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 03:51 PM UTC

PDB ID : 4ZLO / pdb_00004zlo
Title : Serine/threonine-protein kinase PAK1 complexed with a dibenzodiazepine:
identification of an allosteric site on PAK1
Authors : Bellamacina, C.R.; Bussiere, D.E.
Deposited on : 2015-05-01
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	297	5% 84% 9% 6%
1	B	297	10% 79% 9% 11%

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 4239 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase PAK 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	278	Total	C	N	O	P	S	0	0	0
			2095	1329	354	396	1	15			
1	B	264	Total	C	N	O	P	S	0	1	0
			1993	1269	330	379	1	14			

- Molecule 2 is 2,8-difluoro-11-(4-methylpiperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (CCD ID: 4PV) (formula: C₁₈H₁₈F₂N₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	F	N	0	0
			24	18	2	4		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	83	Total	O	0	0
			83	83		
4	B	38	Total	O	0	0
			38	38		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine/threonine-protein kinase PAK 1



- Molecule 1: Serine/threonine-protein kinase PAK 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.51Å 79.34Å 65.79Å 90.00° 108.11° 90.00°	Depositor
Resolution (Å)	52.31 – 2.50 52.31 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.8 (52.31-2.50) 98.8 (52.31-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.03 (at 2.51Å)	Xtriage
Refinement program	BUSTER 2.11.4	Depositor
R, R_{free}	0.195 , 0.222 0.200 , 0.234	Depositor DCC
R_{free} test set	1081 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	50.3	Xtriage
Anisotropy	0.208	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 72.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.024 for l,-k,h	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4239	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, 4PV, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	1/2115 (0.0%)	1.30	5/2863 (0.2%)
1	B	0.85	0/2013	1.33	3/2728 (0.1%)
All	All	0.85	1/4128 (0.0%)	1.32	8/5591 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	288	MET	SD-CE	-8.61	1.58	1.79

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	435	VAL	CA-C-N	6.93	129.29	120.56
1	A	435	VAL	C-N-CA	6.93	129.29	120.56
1	B	427	THR	CB-CA-C	6.89	117.82	110.17
1	B	257	LEU	N-CA-C	-6.46	105.53	113.41
1	B	468	ASN	CA-CB-CG	5.46	118.06	112.60
1	A	428	PRO	CA-C-N	5.31	128.13	120.38
1	A	428	PRO	C-N-CA	5.31	128.13	120.38
1	A	362	ASP	CA-CB-CG	5.05	117.65	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2095	0	2060	12	0
1	B	1993	0	1949	12	0
2	A	24	0	18	1	0
3	A	6	0	8	1	0
4	A	83	0	0	0	0
4	B	38	0	0	0	0
All	All	4239	0	4035	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:433:PRO:HG2	1:B:504:MET:SD	2.41	0.60
1:A:424:MET:HE3	1:A:469:PRO:HB2	1.83	0.59
1:B:462:PRO:HD2	1:B:465:LEU:HD22	1.87	0.56
1:A:408:PHE:H	3:A:602:GOL:H31	1.72	0.54
1:A:316:ILE:HD11	1:A:340:LEU:HD21	1.91	0.52
1:B:255:GLU:C	1:B:257:LEU:H	2.18	0.50
1:A:324:ASN:HB3	1:A:327:ILE:HG12	1.93	0.49
1:A:432:ALA:HB3	1:A:435:VAL:HG23	1.97	0.47
1:A:489:LYS:HA	1:B:489:LYS:HG2	1.96	0.47
1:B:261:VAL:HG13	1:B:334:TYR:HA	1.96	0.47
1:A:429:TYR:CD1	1:A:462:PRO:HB2	2.50	0.46
1:B:428:PRO:HG2	1:B:469:PRO:HB3	1.98	0.46
1:B:324:ASN:HB3	1:B:327:ILE:HD12	1.98	0.46
1:A:271:THR:HG23	1:A:272:ARG:HG3	2.00	0.44
1:A:385:VAL:HG21	2:A:601:4PV:H9	1.99	0.43
1:B:454:ALA:HA	1:B:457:MET:HE3	2.00	0.42
1:B:453:MET:O	1:B:456:GLU:HB2	2.20	0.42
1:A:314:ASN:O	1:A:318:VAL:HG23	2.20	0.42
1:A:344:MET:HE3	1:A:344:MET:HB2	1.82	0.41
1:B:346:TYR:CE2	1:B:348:ALA:HB2	2.56	0.41
1:B:455:ILE:HA	1:B:458:ILE:HD12	2.03	0.41
1:A:454:ALA:HA	1:A:457:MET:HE3	2.03	0.41
1:B:315:GLU:HG2	1:B:319:MET:HE2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	269/297 (91%)	263 (98%)	6 (2%)	0	100	100
1	B	252/297 (85%)	241 (96%)	11 (4%)	0	100	100
All	All	521/594 (88%)	504 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	216/257 (84%)	212 (98%)	4 (2%)	50	76
1	B	207/257 (80%)	202 (98%)	5 (2%)	43	70
All	All	423/514 (82%)	414 (98%)	9 (2%)	47	74

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	263	VAL
1	A	281	SER
1	A	301	MET
1	A	523	ILE
1	B	261	VAL
1	B	333	SER
1	B	419	SER
1	B	424	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	468	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	326	ASN
1	A	486	ASN
1	B	326	ASN
1	B	517	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	TPO	B	423	1	8,10,11	1.12	1 (12%)	10,14,16	1.36	2 (20%)
1	TPO	A	423	1	8,10,11	0.86	0	10,14,16	2.07	3 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	B	423	1	-	0/9/11/13	-
1	TPO	A	423	1	-	2/9/11/13	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	423	TPO	P-OG1	-2.42	1.55	1.59

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	423	TPO	P-OG1-CB	-4.17	111.99	123.33
1	A	423	TPO	CG2-CB-CA	3.43	119.95	113.26
1	B	423	TPO	O3P-P-OG1	2.59	115.95	105.85
1	B	423	TPO	O-C-CA	-2.03	119.56	124.77
1	A	423	TPO	O3P-P-OG1	2.01	113.68	105.85

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	423	TPO	O-C-CA-CB
1	A	423	TPO	C-CA-CB-CG2

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	4PV	A	601	-	27,27,27	0.39	0	39,39,39	0.49	1 (2%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	602	-	5,5,5	0.08	0	5,5,5	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	4PV	A	601	-	-	0/4/26/26	0/4/4/4
3	GOL	A	602	-	-	0/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	4PV	C3-N1-C5	2.10	125.72	119.38

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	4PV	1	0
3	A	602	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	277/297 (93%)	0.25	15 (5%) 31 27	34, 54, 94, 119	0
1	B	263/297 (88%)	0.51	31 (11%) 9 7	34, 63, 121, 138	1 (0%)
All	All	540/594 (90%)	0.38	46 (8%) 16 14	34, 58, 106, 138	1 (0%)

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	254	LEU	4.3
1	B	263	VAL	4.0
1	B	253	ILE	3.6
1	B	293	GLY	3.6
1	B	288	MET	3.2
1	A	250	ASP	3.2
1	B	285	TYR	3.2
1	A	543	ASN	3.2
1	B	287	ALA	3.1
1	A	253	ILE	2.9
1	B	278	GLN	2.9
1	B	261	VAL	2.8
1	B	277	GLY	2.7
1	B	310	GLU	2.7
1	B	296	VAL	2.6
1	B	286	THR	2.6
1	B	271	THR	2.6
1	A	302	ASN	2.5
1	A	312	ILE	2.5
1	A	481	THR	2.5
1	B	255	GLU	2.5
1	B	260	ILE	2.4
1	B	284	VAL	2.4
1	B	477	ALA	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	267	LYS	2.4
1	B	298	ILE	2.4
1	B	492	ALA	2.4
1	B	301	MET	2.3
1	A	311	LEU	2.3
1	B	543	ASN	2.3
1	B	256	LYS	2.3
1	B	360	CYS	2.3
1	A	360	CYS	2.2
1	B	270	TYR	2.2
1	A	488	GLU	2.2
1	A	336	VAL	2.2
1	A	480	GLY	2.2
1	B	273	PHE	2.2
1	A	313	ILE	2.1
1	A	466	ASN	2.1
1	B	258	ARG	2.1
1	B	274	GLU	2.1
1	B	342	VAL	2.1
1	A	258	ARG	2.1
1	B	275	LYS	2.0
1	A	427	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	TPO	A	423	11/12	0.87	0.13	76,81,89,89	0
1	TPO	B	423	11/12	0.95	0.07	67,68,74,77	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	602	6/6	0.73	0.22	67,75,78,80	0
2	4PV	A	601	24/24	0.94	0.08	47,54,66,67	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 4PV A 601:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.