



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 04:39 PM UTC

PDB ID : 3ZMF / pdb_00003zmf
Title : Salmonella enterica SadA 303-358 fused to GCN4 adaptors (SadAK2)
Authors : Hartmann, M.D.; Hernandez Alvarez, B.; Albrecht, R.; Lupas, A.N.
Deposited on : 2013-02-08
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	113	32% 74% 22% ..
1	B	113	21% 75% 20% ..
1	C	113	32% 80% 17% ..

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2550 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GENERAL CONTROL PROTEIN GCN4, PUTATIVE INNER MEMBRANE PROTEIN, GENERAL CONTROL PROTEIN GCN4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	111	Total	C	N	O	S	0	0	0
			807	508	134	164	1			
1	B	111	Total	C	N	O	S	0	0	0
			812	513	131	167	1			
1	C	111	Total	C	N	O	S	0	1	0
			808	509	131	166	2			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	278	ILE	LEU	engineered mutation	UNP P03069
A	282	ILE	VAL	engineered mutation	UNP P03069
A	285	ILE	LEU	engineered mutation	UNP P03069
A	289	ILE	ASN	engineered mutation	UNP P03069
A	292	ILE	LEU	engineered mutation	UNP P03069
A	296	ILE	VAL	engineered mutation	UNP P03069
A	299	ILE	LEU	engineered mutation	UNP P03069
A	303	ILE	VAL	engineered mutation	UNP P03069
A	362	ILE	LEU	engineered mutation	UNP P03069
A	366	ILE	VAL	engineered mutation	UNP P03069
A	369	ILE	LEU	engineered mutation	UNP P03069
A	373	ILE	ASN	engineered mutation	UNP P03069
A	376	ILE	LEU	engineered mutation	UNP P03069
A	380	ILE	VAL	engineered mutation	UNP P03069
A	383	ILE	LEU	engineered mutation	UNP P03069
A	387	ILE	VAL	engineered mutation	UNP P03069
B	278	ILE	LEU	engineered mutation	UNP P03069
B	282	ILE	VAL	engineered mutation	UNP P03069
B	285	ILE	LEU	engineered mutation	UNP P03069
B	289	ILE	ASN	engineered mutation	UNP P03069
B	292	ILE	LEU	engineered mutation	UNP P03069
B	296	ILE	VAL	engineered mutation	UNP P03069

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	299	ILE	LEU	engineered mutation	UNP P03069
B	303	ILE	VAL	engineered mutation	UNP P03069
B	362	ILE	LEU	engineered mutation	UNP P03069
B	366	ILE	VAL	engineered mutation	UNP P03069
B	369	ILE	LEU	engineered mutation	UNP P03069
B	373	ILE	ASN	engineered mutation	UNP P03069
B	376	ILE	LEU	engineered mutation	UNP P03069
B	380	ILE	VAL	engineered mutation	UNP P03069
B	383	ILE	LEU	engineered mutation	UNP P03069
B	387	ILE	VAL	engineered mutation	UNP P03069
C	278	ILE	LEU	engineered mutation	UNP P03069
C	282	ILE	VAL	engineered mutation	UNP P03069
C	285	ILE	LEU	engineered mutation	UNP P03069
C	289	ILE	ASN	engineered mutation	UNP P03069
C	292	ILE	LEU	engineered mutation	UNP P03069
C	296	ILE	VAL	engineered mutation	UNP P03069
C	299	ILE	LEU	engineered mutation	UNP P03069
C	303	ILE	VAL	engineered mutation	UNP P03069
C	362	ILE	LEU	engineered mutation	UNP P03069
C	366	ILE	VAL	engineered mutation	UNP P03069
C	369	ILE	LEU	engineered mutation	UNP P03069
C	373	ILE	ASN	engineered mutation	UNP P03069
C	376	ILE	LEU	engineered mutation	UNP P03069
C	380	ILE	VAL	engineered mutation	UNP P03069
C	383	ILE	LEU	engineered mutation	UNP P03069
C	387	ILE	VAL	engineered mutation	UNP P03069


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	39	Total O 39 39	0	0
2	B	45	Total O 45 45	0	0
2	C	39	Total O 39 39	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GENERAL CONTROL PROTEIN GCN4, PUTATIVE INNER MEMBRANE PROTEIN, GENERAL CONTROL PROTEIN GCN4



- Molecule 1: GENERAL CONTROL PROTEIN GCN4, PUTATIVE INNER MEMBRANE PROTEIN, GENERAL CONTROL PROTEIN GCN4



- Molecule 1: GENERAL CONTROL PROTEIN GCN4, PUTATIVE INNER MEMBRANE PROTEIN, GENERAL CONTROL PROTEIN GCN4



4 Data and refinement statistics

Property	Value	Source
Space group	P 3	Depositor
Cell constants a, b, c, α , β , γ	46.59Å 46.59Å 128.98Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.51 – 1.85 38.51 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.1 (38.51-1.85) 99.9 (38.51-1.85)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.36 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.181 , 0.221 (Not available) , 0.245	Depositor DCC
R_{free} test set	1381 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å ²)	23.7	Xtriage
Anisotropy	0.677	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 78.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.54$, $\langle L^2 \rangle = 0.39$	Xtriage
Estimated twinning fraction	0.001 for -h,-k,l 0.080 for h,-h-k,-l 0.022 for -k,-h,-l	Xtriage
Reported twinning fraction	0.726 for H, K, L 0.274 for K, H, -L	Depositor
Outliers	0 of 26768 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2550	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.59	6/814 (0.7%)	1.43	9/1104 (0.8%)
1	B	1.70	14/819 (1.7%)	1.50	7/1110 (0.6%)
1	C	1.64	10/817 (1.2%)	1.33	4/1108 (0.4%)
All	All	1.65	30/2450 (1.2%)	1.42	20/3322 (0.6%)

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	328	SER	CA-C	8.56	1.63	1.52
1	C	369	ILE	CA-C	8.55	1.63	1.52
1	C	370	LEU	N-CA	7.76	1.55	1.46
1	C	355	ASN	N-CA	6.66	1.54	1.45
1	B	328	SER	N-CA	6.59	1.54	1.46
1	A	350	SER	C-O	6.13	1.31	1.24
1	B	342	ALA	CA-C	6.05	1.60	1.52
1	C	353	ALA	N-CA	6.04	1.53	1.45
1	A	371	SER	CA-C	6.04	1.60	1.52
1	B	339	THR	N-CA	-6.03	1.37	1.46
1	B	353	ALA	N-CA	5.99	1.53	1.46
1	B	319	LEU	N-CA	5.94	1.53	1.45
1	A	351	THR	CA-C	5.83	1.60	1.53
1	C	370	LEU	C-N	5.77	1.41	1.33
1	B	345	THR	CB-CG2	-5.73	1.33	1.52
1	B	332	THR	CA-CB	5.67	1.62	1.53
1	B	334	ASN	C-O	5.62	1.30	1.24
1	B	339	THR	CB-CG2	-5.62	1.34	1.52
1	A	345	THR	CA-CB	5.60	1.62	1.53
1	B	332	THR	CA-C	5.56	1.60	1.53
1	B	355	ASN	N-CA	5.50	1.52	1.45
1	A	334	ASN	CA-C	5.46	1.59	1.52
1	C	363	GLU	CA-C	-5.40	1.46	1.52
1	B	324	ILE	CA-C	5.32	1.58	1.52
1	C	338	ILE	N-CA	-5.28	1.40	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	373	ILE	CA-CB	-5.22	1.48	1.54
1	B	349	ASP	CA-C	5.20	1.60	1.52
1	A	351	THR	N-CA	-5.15	1.40	1.46
1	C	364	ASP	CA-C	-5.05	1.46	1.52
1	C	321	ASP	CA-C	-5.04	1.46	1.52

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	348	ALA	CA-C-N	-7.47	110.63	123.25
1	C	348	ALA	C-N-CA	-7.47	110.63	123.25
1	A	328	SER	N-CA-C	7.39	120.57	108.52
1	A	356	GLY	N-CA-C	6.99	121.11	112.73
1	A	338	ILE	N-CA-C	-6.83	98.62	108.53
1	B	343	ALA	N-CA-C	-6.76	99.82	109.96
1	A	352	ASP	N-CA-CB	6.36	120.77	110.41
1	A	337	LYS	N-CA-C	6.30	119.38	110.50
1	B	342	ALA	CA-C-N	6.15	129.56	120.95
1	B	342	ALA	C-N-CA	6.15	129.56	120.95
1	A	378	ASN	N-CA-C	5.92	121.14	111.37
1	B	355	ASN	CA-C-N	5.88	127.49	120.14
1	B	355	ASN	C-N-CA	5.88	127.49	120.14
1	A	359	MET	CG-SD-CE	-5.75	88.25	100.90
1	B	311	ALA	N-CA-C	5.70	117.95	111.11
1	C	289	ILE	N-CA-C	5.63	116.40	110.72
1	B	344	GLY	N-CA-C	5.46	119.19	112.14
1	C	366	ILE	N-CA-C	5.25	115.88	110.36
1	A	355	ASN	N-CA-C	5.12	117.66	110.14
1	A	278	ILE	N-CA-C	5.01	119.12	112.76

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	807	0	780	6	0
1	B	812	0	791	6	0
1	C	808	0	779	4	0
2	A	39	0	0	0	0
2	B	45	0	0	2	0
2	C	39	0	0	2	0
All	All	2550	0	2350	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:331:HIS:O	1:A:332:THR:HG22	1.98	0.62
1:C:374:TYR:OH	2:C:2037:HOH:O	2.11	0.61
1:A:332:THR:HG23	1:A:332:THR:O	2.02	0.58
1:B:365:LYS:NZ	2:B:2045:HOH:O	2.39	0.56
1:B:358:GLN:HA	1:B:361:GLN:NE2	2.28	0.48
1:C:278:ILE:C	1:C:278:ILE:HD12	2.40	0.47
1:A:332:THR:HG23	1:A:334:ASN:HD22	1.80	0.46
1:B:278:ILE:O	1:B:282:ILE:HG12	2.16	0.45
1:A:349:ASP:OD1	1:A:349:ASP:N	2.50	0.44
1:C:334:ASN:ND2	2:C:2018:HOH:O	2.49	0.44
1:C:298:ARG:O	1:C:302:LEU:HB2	2.18	0.43
1:B:332:THR:OG1	2:B:2027:HOH:O	2.04	0.43
1:A:308:ALA:O	1:A:311:ALA:N	2.54	0.41
1:B:281:LYS:HA	1:B:284:GLU:OE2	2.21	0.41
1:A:329:ALA:O	1:A:336:SER:HB2	2.21	0.41
1:B:339:THR:O	1:B:340:ASN:HB2	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	109/113 (96%)	107 (98%)	2 (2%)	0	100	100
1	B	109/113 (96%)	108 (99%)	1 (1%)	0	100	100
1	C	110/113 (97%)	110 (100%)	0	0	100	100
All	All	328/339 (97%)	325 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	79/95 (83%)	72 (91%)	7 (9%)	9	1
1	B	81/95 (85%)	76 (94%)	5 (6%)	16	5
1	C	79/95 (83%)	73 (92%)	6 (8%)	12	3
All	All	239/285 (84%)	221 (92%)	18 (8%)	12	3

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	278	ILE
1	A	299	ILE
1	A	302	LEU
1	A	304	THR
1	A	305	ASN
1	A	309	SER
1	A	325	SER
1	B	279	GLU
1	B	302	LEU
1	B	365	LYS
1	B	372	LYS
1	B	373	ILE
1	C	278	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	286	LEU
1	C	299	ILE
1	C	313	LEU
1	C	334	ASN
1	C	362	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	334	ASN
1	A	378	ASN
1	B	361	GLN
1	C	334	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	111/113 (98%)	1.52	36 (32%) 1 1	18, 38, 103, 111	0
1	B	111/113 (98%)	1.09	24 (21%) 2 2	14, 33, 82, 102	0
1	C	111/113 (98%)	1.37	36 (32%) 1 1	18, 36, 81, 93	1 (0%)
All	All	333/339 (98%)	1.33	96 (28%) 1 1	14, 36, 94, 111	1 (0%)

All (96) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	277	GLN	5.5
1	A	278	ILE	5.0
1	C	302	LEU	4.9
1	A	286	LEU	4.8
1	A	386	LEU	4.8
1	A	302	LEU	4.7
1	A	280	ASP	4.6
1	A	298	ARG	4.4
1	A	290	TYR	4.3
1	B	297	ALA	4.1
1	A	297	ALA	4.1
1	A	292	ILE	4.1
1	A	304	THR	4.0
1	C	298	ARG	3.9
1	B	300	LYS	3.9
1	C	285	ILE	3.7
1	A	287	SER	3.7
1	B	304	THR	3.6
1	B	387	ILE	3.5
1	B	381	ALA	3.5
1	C	291	HIS	3.5
1	B	374	TYR	3.5
1	C	277	GLN	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	290	TYR	3.5
1	C	292	ILE	3.5
1	B	294	ASN	3.4
1	B	375	HIS	3.4
1	C	284	GLU	3.4
1	A	299	ILE	3.4
1	C	308	ALA	3.3
1	B	302	LEU	3.3
1	C	386	LEU	3.3
1	C	297	ALA	3.2
1	C	295	GLU	3.2
1	A	295	GLU	3.2
1	C	305	ASN	3.1
1	C	387	ILE	3.1
1	C	301	LYS	3.1
1	C	307	GLU	3.1
1	B	277	GLN	3.1
1	A	296	ILE	3.0
1	A	284	GLU	3.0
1	C	296	ILE	2.9
1	A	309	SER	2.9
1	B	386	LEU	2.9
1	A	314	ALA	2.9
1	B	290	TYR	2.8
1	B	291	HIS	2.8
1	C	348	ALA	2.8
1	A	374	TYR	2.8
1	A	300	LYS	2.8
1	A	285	ILE	2.8
1	C	278	ILE	2.8
1	A	313	LEU	2.7
1	B	371	SER	2.7
1	A	283	GLU	2.7
1	B	278	ILE	2.6
1	A	308	ALA	2.6
1	C	279	GLU	2.6
1	C	282	ILE	2.6
1	C	312	GLY	2.6
1	A	291	HIS	2.6
1	A	311	ALA	2.6
1	C	311	ALA	2.6
1	B	298	ARG	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	287	SER	2.5
1	C	361	GLN	2.5
1	A	289	ILE	2.5
1	A	387	ILE	2.5
1	B	293	GLU	2.4
1	A	282	ILE	2.4
1	B	282	ILE	2.4
1	B	296	ILE	2.4
1	C	299	ILE	2.4
1	C	303	ILE	2.4
1	B	384	LYS	2.3
1	B	287	SER	2.3
1	A	382	ARG	2.3
1	C	313	LEU	2.3
1	C	314	ALA	2.3
1	A	333	GLY	2.3
1	A	334	ASN	2.3
1	C	286	LEU	2.3
1	C	306	THR	2.3
1	A	307	GLU	2.2
1	A	303	ILE	2.2
1	A	281	LYS	2.2
1	B	373	ILE	2.1
1	A	288	LYS	2.1
1	C	293	GLU	2.1
1	C	332	THR	2.1
1	C	283	GLU	2.1
1	B	305	ASN	2.1
1	B	286	LEU	2.1
1	C	294	ASN	2.0
1	C	289	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.