



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 01:56 AM UTC

PDB ID : 3ZMN / pdb_00003zmn
Title : VP17, a capsid protein of bacteriophage P23-77
Authors : Rissanen, I.; Grimes, J.M.; Pawlowski, A.; Mantynen, S.; Harlos, K.; Bamford, J.K.H.; Stuart, D.I.
Deposited on : 2013-02-11
Resolution : 2.26 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	291	7% 71% 7% 22%
1	B	291	7% 70% 6% 23%
2	C	8	75% 25%
2	D	8	75% 25%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4047 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called VP17.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	228	Total	C	N	O	S	0	2	0
			1791	1137	319	328	7			
1	B	224	Total	C	N	O	S	0	0	0
			1752	1114	310	322	6			

- Molecule 2 is a protein called VP17.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	8	Total	C	N	O	0	0	0
			40	24	8	8			
2	D	8	Total	C	N	O	0	0	0
			40	24	8	8			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	206	Total	O	0	0
			206	206		
3	B	209	Total	O	0	0
			209	209		
3	C	8	Total	O	0	0
			8	8		
3	D	1	Total	O	0	0
			1	1		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: VP17




• Molecule 1: VP17



• Molecule 2: VP17



• Molecule 2: VP17



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	107.21Å 107.21Å 233.78Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.94 – 2.26 43.94 – 2.27	Depositor EDS
% Data completeness (in resolution range)	99.4 (43.94-2.26) 100.0 (43.94-2.27)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.02 (at 2.27Å)	Xtriage
Refinement program	BUSTER 2.11.2	Depositor
R, R_{free}	0.185 , 0.213 0.187 , 0.220	Depositor DCC
R_{free} test set	1891 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	42.1	Xtriage
Anisotropy	0.142	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 71.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4047	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	1/1844 (0.1%)	1.14	10/2516 (0.4%)
1	B	0.80	0/1799	1.12	4/2457 (0.2%)
All	All	0.81	1/3643 (0.0%)	1.13	14/4973 (0.3%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	219	ASP	CA-C	5.60	1.60	1.52

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	224	GLU	CA-C-N	9.31	138.46	121.70
1	B	224	GLU	C-N-CA	9.31	138.46	121.70
1	A	193	VAL	N-CA-CB	8.61	122.31	111.46
1	A	219	ASP	CA-C-N	7.41	135.04	121.70
1	A	219	ASP	C-N-CA	7.41	135.04	121.70
1	A	221	ASN	CA-C-N	6.01	132.52	121.70
1	A	221	ASN	C-N-CA	6.01	132.52	121.70
1	A	258	SER	N-CA-C	5.87	118.46	111.71
1	A	230	PRO	N-CA-C	5.56	121.43	113.47
1	A	148	THR	N-CA-C	5.51	117.94	109.07
1	B	258	SER	N-CA-C	5.32	117.82	111.71
1	A	209	ASN	CA-CB-CG	5.15	117.75	112.60
1	B	218	VAL	N-CA-C	5.07	119.89	109.34
1	A	53	ILE	N-CA-C	-5.01	99.92	107.73

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1791	0	1726	7	0
1	B	1752	0	1680	4	0
2	C	40	0	11	1	0
2	D	40	0	10	1	0
3	A	206	0	0	0	0
3	B	209	0	0	0	0
3	C	8	0	0	0	0
3	D	1	0	0	0	0
All	All	4047	0	3427	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:3:UNK:HA	2:D:4:UNK:CB	2.18	0.74
1:A:219:ASP:HB2	1:A:220:GLN:HA	1.81	0.62
1:A:219:ASP:HB2	1:A:220:GLN:CA	2.30	0.61
1:B:214:THR:HG22	1:B:227:ILE:HD11	1.86	0.57
2:C:7:UNK:N	2:C:8:UNK:HA	2.21	0.55
1:B:84:PHE:HB2	1:B:235:LEU:HB2	1.90	0.53
1:A:219:ASP:HB2	1:A:220:GLN:C	2.35	0.52
1:A:179:GLU:OE1	1:A:257:ARG:NH1	2.44	0.49
1:B:75:GLN:HG3	1:B:242:VAL:HG22	1.94	0.49
1:A:124:THR:CG2	1:A:183:VAL:HG11	2.48	0.43
1:A:133:SER:OG	1:A:135:TYR:O	2.32	0.43
1:B:45:VAL:HG12	1:B:271:ILE:HG12	2.01	0.42
1:A:125:PRO:HG2	1:A:219:ASP:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	226/291 (78%)	218 (96%)	4 (2%)	4 (2%)	6	3
1	B	220/291 (76%)	208 (94%)	9 (4%)	3 (1%)	9	5
All	All	446/582 (77%)	426 (96%)	13 (3%)	7 (2%)	7	4

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	219	ASP
1	A	223	VAL
1	A	224	GLU
1	A	226	MET
1	B	218	VAL
1	B	219	ASP
1	B	221	ASN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	186/234 (80%)	181 (97%)	5 (3%)	39	49
1	B	182/234 (78%)	176 (97%)	6 (3%)	33	42
All	All	368/468 (79%)	357 (97%)	11 (3%)	36	45

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	80	GLU
1	A	122	MET
1	A	193	VAL
1	A	219	ASP
1	A	231	GLN
1	B	43	VAL
1	B	80	GLU
1	B	195	ARG
1	B	196	ASP
1	B	231	GLN
1	B	267	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	150	GLN
1	B	44	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	228/291 (78%)	0.07	20 (8%) 15 14	22, 40, 85, 122	2 (0%)
1	B	224/291 (76%)	0.32	19 (8%) 16 15	30, 46, 92, 138	0
2	C	0/8	-	-	-	-
2	D	0/8	-	-	-	-
All	All	452/598 (75%)	0.20	39 (8%) 16 15	22, 43, 91, 138	2 (0%)

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	273	VAL	6.7
1	B	43	VAL	6.6
1	A	220	GLN	5.6
1	A	273	VAL	5.3
1	B	222	ASN	5.2
1	B	221	ASN	5.1
1	A	42	ALA	4.6
1	A	221	ASN	4.6
1	B	225	THR	4.4
1	B	223	VAL	4.1
1	A	225	THR	4.1
1	A	196	ASP	3.9
1	A	203	SER	3.8
1	A	41	ALA	3.7
1	B	224	GLU	3.6
1	A	222	ASN	3.6
1	B	226	MET	3.5
1	A	219	ASP	3.4
1	A	223	VAL	3.3
1	A	202	ASP	3.2
1	B	204	ALA	3.1
1	B	218	VAL	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	271	ILE	3.1
1	A	135	TYR	3.0
1	A	218	VAL	3.0
1	B	196	ASP	2.7
1	A	224	GLU	2.5
1	A	43	VAL	2.5
1	B	267	MET	2.5
1	B	53	ILE	2.5
1	B	93	LEU	2.3
1	A	272	GLU	2.3
1	B	219	ASP	2.2
1	A	226	MET	2.2
1	B	227	ILE	2.2
1	A	147	SER	2.1
1	B	270	ARG	2.0
1	A	148	THR	2.0
1	B	237	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.