



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 01:02 PM UTC

PDB ID : 3ZMR / pdb_00003zmr
Title : Bacteroides ovatus GH5 xyloglucanase in complex with a XXXG heptasaccharide
Authors : Larsbrink, J.; Rogers, T.E.; Hemsworth, G.R.; McKee, L.S.; Spadiut, O.; Klintner, S.; Pudlo, N.A.; Urs, K.; Kelly, A.G.; Cederholm, S.N.; Davies, G.J.; Martens, E.C.; Brumer, H.
Deposited on : 2013-02-12
Resolution : 1.43 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3234 (1.46-1.42)
Clashscore	190562	3289 (1.46-1.42)
Ramachandran outliers	187476	3248 (1.46-1.42)
Sidechain outliers	187428	3248 (1.46-1.42)
RSRZ outliers	180081	3234 (1.46-1.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	475	2% 97% ..
1	B	475	% 92% 5% .
2	C	7	57% 29% 14%
3	D	7	57% 29% 14%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GLO	D	1	-	X	-	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 8939 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CELLULASE (GLYCOSYL HYDROLASE FAMILY 5).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	469	Total	C	N	O	S	0	7	0
			3697	2325	638	720	14			
1	B	462	Total	C	N	O	S	0	14	0
			3677	2324	622	716	15			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	36	MET	-	expression tag	UNP A7LXT7
A	503	LEU	-	expression tag	UNP A7LXT7
A	504	GLU	-	expression tag	UNP A7LXT7
A	505	HIS	-	expression tag	UNP A7LXT7
A	506	HIS	-	expression tag	UNP A7LXT7
A	507	HIS	-	expression tag	UNP A7LXT7
A	508	HIS	-	expression tag	UNP A7LXT7
A	509	HIS	-	expression tag	UNP A7LXT7
A	510	HIS	-	expression tag	UNP A7LXT7
B	36	MET	-	expression tag	UNP A7LXT7
B	503	LEU	-	expression tag	UNP A7LXT7
B	504	GLU	-	expression tag	UNP A7LXT7
B	505	HIS	-	expression tag	UNP A7LXT7
B	506	HIS	-	expression tag	UNP A7LXT7
B	507	HIS	-	expression tag	UNP A7LXT7
B	508	HIS	-	expression tag	UNP A7LXT7
B	509	HIS	-	expression tag	UNP A7LXT7
B	510	HIS	-	expression tag	UNP A7LXT7

- Molecule 2 is an oligosaccharide called alpha-D-xylopyranose-(1-6)-beta-D-glucopyranose-(1-4)-[alpha-D-xylopyranose-(1-6)]beta-D-glucopyranose-(1-4)-[alpha-D-xylopyranose-(1-6)]beta-D-glucopyranose-(1-4)-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	7	Total	C	O	0	1	0
			71	39	32			

- Molecule 3 is an oligosaccharide called alpha-D-xylopyranose-(1-6)-beta-D-glucopyranose-(1-4)-[alpha-D-xylopyranose-(1-6)]beta-D-glucopyranose-(1-4)-[alpha-D-xylopyranose-(1-6)]beta-D-glucopyranose-(1-4)-D-glucose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	D	7	Total	C	O	0	0	0
			72	39	33			

- Molecule 4 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN E-1,3-DIOL (CCD ID: BTB) (formula: C₈H₁₉NO₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is alpha-D-glucopyranose (CCD ID: GLC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	1
			11	6	5		


- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	724	Total	O	0	0
			724	724		
7	B	617	Total	O	0	0
			617	617		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CELLULASE (GLYCOSYL HYDROLASE FAMILY 5)

Chain A: 



- Molecule 1: CELLULASE (GLYCOSYL HYDROLASE FAMILY 5)

Chain B: 



- Molecule 2: alpha-D-xylopyranose-(1-6)-beta-D-glucopyranose-(1-4)-[alpha-D-xylopyranose-(1-6)]beta-D-glucopyranose-(1-4)-[alpha-D-xylopyranose-(1-6)]beta-D-glucopyranose-(1-4)-beta-D-glucopyranose

Chain C: 



- Molecule 3: alpha-D-xylopyranose-(1-6)-beta-D-glucopyranose-(1-4)-[alpha-D-xylopyranose-(1-6)]beta-D-glucopyranose-(1-4)-[alpha-D-xylopyranose-(1-6)]beta-D-glucopyranose-(1-4)-D-glucose

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	46.83Å 147.24Å 84.14Å 90.00° 92.84° 90.00°	Depositor
Resolution (Å)	72.99 – 1.43 72.99 – 1.43	Depositor EDS
% Data completeness (in resolution range)	99.5 (72.99-1.43) 99.5 (72.99-1.43)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.44 (at 1.43Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.122 , 0.157 0.130 , 0.163	Depositor DCC
R_{free} test set	10458 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	13.3	Xtriage
Anisotropy	0.256	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 45.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.029 for h,-k,-l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	8939	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BTB, BGC, EDO, GLC, XYS, GLO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	0/3801	0.81	0/5173
1	B	0.72	1/3800 (0.0%)	0.82	2/5174 (0.0%)
All	All	0.72	1/7601 (0.0%)	0.82	2/10347 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	302	ASP	N-CA	5.33	1.52	1.46

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	307	GLN	CA-C-N	-6.69	112.87	119.76
1	B	307	GLN	C-N-CA	-6.69	112.87	119.76

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	502	ASN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3697	0	3534	6	0
1	B	3677	0	3550	21	0
2	C	71	0	53	1	0
3	D	72	0	60	3	0
4	A	14	0	19	2	0
5	A	28	0	42	0	0
5	B	28	0	42	8	0
6	A	11	0	3	0	0
7	A	724	0	0	1	0
7	B	617	0	0	3	0
All	All	8939	0	7303	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:347[B]:ASP:OD1	5:B:1505:EDO:H12	1.51	1.09
1:B:347[B]:ASP:OD1	5:B:1505:EDO:C1	2.13	0.96
1:B:271:ARG:NH1	5:B:1510:EDO:O1	2.12	0.82
4:A:1511:BTB:H31	4:A:1511:BTB:O8	1.81	0.80
1:B:365[B]:ARG:NH1	7:B:2431:HOH:O	2.20	0.73
1:B:69:ALA:O	1:B:88:GLU:HG2	1.89	0.72
1:B:229[A]:MET:HE3	1:B:229[A]:MET:HA	1.74	0.70
1:A:289:ARG:NH1	7:A:2416:HOH:O	2.25	0.69
1:A:297:GLU:OE2	2:C:1[B]:BGC:C1	2.41	0.68
1:B:336:TYR:CD1	5:B:1507:EDO:H21	2.30	0.66
1:B:105:GLU:OE1	5:B:1509:EDO:H21	2.01	0.59
1:B:251:HIS:CE1	3:D:1:GLO:H1	2.38	0.59
1:A:159:ALA:HB2	1:A:501:LEU:HD21	1.87	0.55
1:B:502:ASN:OD1	1:B:502:ASN:O	2.28	0.52
1:B:231:LYS:HE2	7:B:2235:HOH:O	2.10	0.50
1:B:160:GLY:HA3	1:B:205[B]:VAL:HG13	1.96	0.47
1:A:357[B]:MET:SD	1:A:358:PRO:HD2	2.55	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:347[B]:ASP:OD1	5:B:1505:EDO:O1	2.34	0.45
1:B:501:LEU:C	1:B:503:LEU:H	2.25	0.44
5:B:1509:EDO:H22	7:B:2121:HOH:O	2.18	0.44
1:B:377:ASP:HA	1:B:381[B]:MET:HB2	2.00	0.44
4:A:1511:BTB:O8	4:A:1511:BTB:C3	2.61	0.43
1:B:215:GLN:HG3	1:B:228:TRP:CD2	2.54	0.43
1:A:163:VAL:HG23	1:A:206:VAL:CG2	2.49	0.42
1:B:229[A]:MET:HE3	1:B:229[A]:MET:CA	2.42	0.42
1:B:251:HIS:NE2	3:D:1:GLO:H1	2.35	0.41
1:B:296:ASN:HD21	3:D:1:GLO:C1	2.34	0.41
1:B:125:LYS:HD3	5:B:1509:EDO:H11	2.04	0.40
1:B:212:TYR:CE2	1:B:229[B]:MET:HE1	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	474/475 (100%)	457 (96%)	17 (4%)	0	100	100
1	B	474/475 (100%)	460 (97%)	14 (3%)	0	100	100
All	All	948/950 (100%)	917 (97%)	31 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	391/394 (99%)	390 (100%)	1 (0%)	86	74
1	B	393/394 (100%)	392 (100%)	1 (0%)	86	74
All	All	784/788 (100%)	782 (100%)	2 (0%)	86	74

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60	ARG
1	B	103	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	103	GLN
1	A	123	ASN
1	A	165	ASN
1	A	299	ASN
1	A	304	ASN
1	A	364	ASN
1	B	61	GLN
1	B	123	ASN
1	B	264	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	C	1[B]	2	11,11,12	0.67	0	15,15,17	1.91	4 (26%)
2	BGC	C	2	2,6	11,11,12	0.63	0	15,15,17	1.05	0
2	BGC	C	3	2	11,11,12	0.54	0	15,15,17	0.82	0
2	BGC	C	4	2	11,11,12	0.45	0	15,15,17	1.00	0
2	XYS	C	5	2	9,9,10	0.64	0	10,12,14	0.66	0
2	XYS	C	6	2	9,9,10	0.47	0	10,12,14	1.39	1 (10%)
2	XYS	C	7	2	9,9,10	0.83	0	10,12,14	1.64	3 (30%)
3	GLO	D	1	3	10,11,11	1.50	2 (20%)	13,14,14	4.19	8 (61%)
3	BGC	D	2	3	11,11,12	1.42	2 (18%)	15,15,17	1.55	3 (20%)
3	BGC	D	3	3	11,11,12	0.49	0	15,15,17	0.98	0
3	BGC	D	4	3	11,11,12	0.63	0	15,15,17	0.67	0
3	XYS	D	5	3	9,9,10	0.83	0	10,12,14	0.72	0
3	XYS	D	6	3	9,9,10	0.66	0	10,12,14	0.97	0
3	XYS	D	7	3	9,9,10	0.90	0	10,12,14	1.22	2 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	C	1[B]	2	-	0/2/19/22	0/1/1/1
2	BGC	C	2	2,6	-	1/2/19/22	0/1/1/1
2	BGC	C	3	2	-	0/2/19/22	0/1/1/1
2	BGC	C	4	2	-	0/2/19/22	0/1/1/1
2	XYS	C	5	2	-	-	0/1/1/1
2	XYS	C	6	2	-	-	0/1/1/1
2	XYS	C	7	2	-	-	0/1/1/1
3	GLO	D	1	3	-	14/15/16/16	-
3	BGC	D	2	3	-	2/2/19/22	0/1/1/1
3	BGC	D	3	3	-	0/2/19/22	0/1/1/1
3	BGC	D	4	3	-	0/2/19/22	0/1/1/1
3	XYS	D	5	3	-	-	0/1/1/1
3	XYS	D	6	3	-	-	0/1/1/1
3	XYS	D	7	3	-	-	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	2	BGC	C2-C3	2.86	1.56	1.52
3	D	2	BGC	C1-C2	2.84	1.59	1.52
3	D	1	GLO	O4-C4	2.63	1.49	1.43
3	D	1	GLO	C3-C2	-2.07	1.50	1.53

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1	GLO	C5-C4-C3	-10.16	96.88	112.48
3	D	1	GLO	O4-C4-C5	6.80	124.38	108.93
3	D	1	GLO	O4-C4-C3	-5.01	97.77	109.46
2	C	1[B]	BGC	C1-C2-C3	4.93	116.82	109.64
3	D	1	GLO	O3-C3-C2	4.01	116.59	109.13
2	C	6	XYS	C5-O5-C1	3.45	116.98	111.42
2	C	1[B]	BGC	C1-O5-C5	3.16	116.43	112.19
3	D	1	GLO	O5-C5-C6	-3.08	102.02	109.03
2	C	1[B]	BGC	C3-C4-C5	-3.07	104.67	110.23
3	D	1	GLO	C4-C3-C2	-2.90	108.41	113.49
3	D	2	BGC	O2-C2-C1	2.81	115.66	109.22
2	C	7	XYS	C5-O5-C1	2.73	115.82	111.42
3	D	1	GLO	C3-C2-C1	-2.63	103.03	111.03
2	C	7	XYS	O2-C2-C1	2.55	115.06	109.22
3	D	2	BGC	O6-C6-C5	-2.52	102.75	111.33
3	D	1	GLO	C6-C5-C4	2.44	117.14	112.17
3	D	7	XYS	C5-O5-C1	2.37	115.24	111.42
3	D	7	XYS	C5-C4-C3	2.34	113.06	109.64
2	C	1[B]	BGC	O4-C4-C5	2.34	115.09	109.32
2	C	7	XYS	C5-C4-C3	2.28	112.96	109.64
3	D	2	BGC	O3-C3-C2	2.26	114.68	110.05

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	1	GLO	C1-C2-C3-C4
3	D	1	GLO	C2-C3-C4-C5
3	D	1	GLO	C2-C3-C4-O4
3	D	1	GLO	O3-C3-C4-C5
3	D	1	GLO	O4-C4-C5-C6
3	D	1	GLO	O4-C4-C5-O5
3	D	1	GLO	O5-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	D	1	GLO	C4-C5-C6-O6
3	D	1	GLO	O3-C3-C4-O4
3	D	2	BGC	C4-C5-C6-O6
3	D	1	GLO	O2-C2-C3-O3
2	C	2	BGC	C4-C5-C6-O6
3	D	1	GLO	O2-C2-C3-C4
3	D	1	GLO	C1-C2-C3-O3
3	D	2	BGC	O5-C5-C6-O6
3	D	1	GLO	C3-C4-C5-C6
3	D	1	GLO	C3-C4-C5-O5


There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1[B]	BGC	1	0
3	D	1	GLO	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	BTB	A	1511	-	13,13,13	0.82	0	7,16,16	1.03	0
5	EDO	A	1517	-	3,3,3	0.34	0	2,2,2	0.20	0
5	EDO	A	1515	-	3,3,3	0.39	0	2,2,2	0.59	0
6	GLC	A	1519[A]	2	11,11,12	0.87	0	15,15,17	1.25	1 (6%)
5	EDO	A	1514	-	3,3,3	0.38	0	2,2,2	0.46	0
5	EDO	A	1513	-	3,3,3	0.43	0	2,2,2	0.20	0
5	EDO	B	1504	-	3,3,3	0.51	0	2,2,2	0.32	0
5	EDO	B	1505	-	3,3,3	0.43	0	2,2,2	0.33	0
5	EDO	A	1512	-	3,3,3	0.49	0	2,2,2	0.45	0
5	EDO	B	1509	-	3,3,3	0.53	0	2,2,2	0.26	0
5	EDO	B	1507	-	3,3,3	0.35	0	2,2,2	0.35	0
5	EDO	B	1510	-	3,3,3	0.44	0	2,2,2	0.18	0
5	EDO	A	1516	-	3,3,3	0.57	0	2,2,2	0.26	0
5	EDO	B	1506	-	3,3,3	0.45	0	2,2,2	0.29	0
5	EDO	A	1600	-	3,3,3	0.45	0	2,2,2	0.28	0
5	EDO	B	1508	-	3,3,3	0.40	0	2,2,2	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BTB	A	1511	-	-	1/21/21/21	-
5	EDO	A	1517	-	-	1/1/1/1	-
5	EDO	A	1515	-	-	0/1/1/1	-
6	GLC	A	1519[A]	2	-	2/2/19/22	0/1/1/1
5	EDO	A	1514	-	-	0/1/1/1	-
5	EDO	A	1513	-	-	1/1/1/1	-
5	EDO	B	1504	-	-	0/1/1/1	-
5	EDO	B	1505	-	-	1/1/1/1	-
5	EDO	A	1512	-	-	0/1/1/1	-
5	EDO	B	1509	-	-	1/1/1/1	-
5	EDO	B	1507	-	-	1/1/1/1	-
5	EDO	B	1510	-	-	1/1/1/1	-
5	EDO	A	1516	-	-	0/1/1/1	-
5	EDO	B	1506	-	-	0/1/1/1	-
5	EDO	A	1600	-	-	0/1/1/1	-
5	EDO	B	1508	-	-	0/1/1/1	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	1519[A]	GLC	C2-C3-C4	-2.10	107.17	110.86

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	1519[A]	GLC	O5-C5-C6-O6
6	A	1519[A]	GLC	C4-C5-C6-O6
5	A	1513	EDO	O1-C1-C2-O2
5	B	1505	EDO	O1-C1-C2-O2
4	A	1511	BTB	C6-C5-N-C7
5	B	1507	EDO	O1-C1-C2-O2
5	B	1510	EDO	O1-C1-C2-O2
5	A	1517	EDO	O1-C1-C2-O2
5	B	1509	EDO	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1511	BTB	2	0
5	B	1505	EDO	3	0
5	B	1509	EDO	3	0
5	B	1507	EDO	1	0
5	B	1510	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	469/475 (98%)	-0.44	10 (2%) 63 67	7, 13, 44, 111	7 (1%)
1	B	462/475 (97%)	-0.45	5 (1%) 78 82	6, 14, 45, 95	14 (3%)
All	All	931/950 (98%)	-0.44	15 (1%) 70 75	6, 14, 45, 111	21 (2%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	503	LEU	5.3
1	A	506	HIS	4.8
1	A	503	LEU	4.7
1	A	116	GLY	4.4
1	A	507	HIS	3.9
1	A	505	HIS	3.8
1	A	509	HIS	3.8
1	B	69	ALA	3.5
1	A	510	HIS	3.4
1	A	504	GLU	2.9
1	B	86	ILE	2.5
1	A	508	HIS	2.4
1	B	303	ALA	2.3
1	A	42	SER	2.2
1	B	89	GLY	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BGC	D	2	11/12	0.85	0.12	19,27,48,68	0
3	XYS	D	7	9/10	0.87	0.13	26,33,44,104	0
2	XYS	C	6	9/10	0.88	0.17	37,42,162,212	0
3	GLO	D	1	12/12	0.88	0.16	17,41,76,136	0
3	XYS	D	6	9/10	0.89	0.13	28,43,90,152	0
2	BGC	C	1[B]	11/12	0.89	0.12	15,25,41,61	11
3	XYS	D	5	9/10	0.93	0.08	28,35,42,45	0
3	BGC	D	4	11/12	0.94	0.08	29,33,58,67	0
2	XYS	C	7	9/10	0.95	0.09	19,24,32,48	0
2	BGC	C	4	11/12	0.96	0.07	19,23,32,35	0
3	BGC	D	3	11/12	0.96	0.06	20,25,36,37	0
2	BGC	C	2	11/12	0.96	0.08	17,21,28,30	0
2	XYS	C	5	9/10	0.97	0.06	16,18,24,26	0
2	BGC	C	3	11/12	0.97	0.07	17,21,24,29	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	EDO	B	1509	4/4	0.75	0.16	36,40,48,58	0
5	EDO	B	1510	4/4	0.83	0.17	21,39,42,56	0
5	EDO	A	1600	4/4	0.88	0.11	35,36,41,77	0
5	EDO	B	1507	4/4	0.91	0.19	18,40,51,96	0
5	EDO	B	1504	4/4	0.91	0.14	31,46,50,57	0
5	EDO	B	1505	4/4	0.91	0.14	23,44,52,79	0
4	BTB	A	1511	14/14	0.93	0.12	17,29,70,97	0
5	EDO	A	1513	4/4	0.93	0.11	24,30,35,64	0
5	EDO	A	1515	4/4	0.93	0.11	29,31,34,44	0
6	GLC	A	1519[A]	11/12	0.93	0.13	7,18,70,90	11
5	EDO	B	1506	4/4	0.94	0.08	26,30,32,39	0
5	EDO	A	1517	4/4	0.95	0.07	28,29,30,42	0
5	EDO	B	1508	4/4	0.96	0.08	26,26,36,47	0
5	EDO	A	1516	4/4	0.96	0.09	21,28,30,53	0
5	EDO	A	1514	4/4	0.97	0.09	19,23,36,59	0
5	EDO	A	1512	4/4	0.98	0.06	20,21,21,22	0

6.5 Other polymers

There are no such residues in this entry.