



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 12:53 PM UTC

PDB ID : 5ZMQ / pdb_00005zmq
Title : Crystal structure of Zika NS3 protease with phenylacetyl-Lys-Lys-Arg-COO
H inhibitor
Authors : Phoo, W.W.; Wirawan, M.
Deposited on : 2018-04-05
Resolution : 1.99 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1506 (1.98-1.98)
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	53	
1	C	53	
1	E	53	
1	G	53	
2	B	178	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	178	76% 10% 13%
2	F	178	70% 13% 16%
2	H	178	77% 10% 13%
3	I	4	75% 25%
3	K	4	25% 75%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6187 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine protease subunit NS2B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	38	Total	C	N	O	S	0	0	0
			288	178	46	63	1			
1	C	37	Total	C	N	O	S	0	0	0
			280	175	46	58	1			
1	E	38	Total	C	N	O	S	0	0	0
			285	177	46	61	1			
1	G	39	Total	C	N	O	S	0	0	0
			293	181	47	64	1			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	44	MET	-	expression tag	UNP Q32ZE1
A	45	THR	-	expression tag	UNP Q32ZE1
C	44	MET	-	expression tag	UNP Q32ZE1
C	45	THR	-	expression tag	UNP Q32ZE1
E	44	MET	-	expression tag	UNP Q32ZE1
E	45	THR	-	expression tag	UNP Q32ZE1
G	44	MET	-	expression tag	UNP Q32ZE1
G	45	THR	-	expression tag	UNP Q32ZE1

- Molecule 2 is a protein called Serine protease NS3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	153	Total	C	N	O	S	0	0	0
			1117	706	192	215	4			
2	D	154	Total	C	N	O	S	0	0	0
			1118	709	191	214	4			
2	F	150	Total	C	N	O	S	0	0	0
			1098	698	187	209	4			
2	H	154	Total	C	N	O	S	0	0	0
			1131	718	195	214	4			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	GLY	-	expression tag	UNP H8XX12
B	143	SER	CYS	engineered mutation	UNP H8XX12
D	0	GLY	-	expression tag	UNP H8XX12
D	143	SER	CYS	engineered mutation	UNP H8XX12
F	0	GLY	-	expression tag	UNP H8XX12
F	143	SER	CYS	engineered mutation	UNP H8XX12
H	0	GLY	-	expression tag	UNP H8XX12
H	143	SER	CYS	engineered mutation	UNP H8XX12

- Molecule 3 is a protein (with D amino acids) called peptide PAC-DLY-DLY-DAR.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	I	4	Total	C	N	O	0	0	0
			39	26	8	5			
3	K	4	Total	C	N	O	0	0	0
			39	26	8	5			

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	37	Total	O	0	0
			37	37		
4	B	92	Total	O	0	0
			92	92		
4	C	37	Total	O	0	0
			37	37		
4	D	80	Total	O	0	0
			80	80		
4	E	33	Total	O	0	0
			33	33		
4	F	87	Total	O	0	0
			87	87		
4	G	32	Total	O	0	0
			32	32		
4	H	94	Total	O	0	0
			94	94		
4	I	6	Total	O	0	0
			6	6		
4	K	1	Total	O	0	0
			1	1		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine protease subunit NS2B

Chain A: 



- Molecule 1: Serine protease subunit NS2B

Chain C: 



- Molecule 1: Serine protease subunit NS2B

Chain E: 




- Molecule 1: Serine protease subunit NS2B

Chain G: 




- Molecule 2: Serine protease NS3

Chain B: 




- Molecule 2: Serine protease NS3

Chain D:  76% 10% 13%




- Molecule 2: Serine protease NS3

Chain F:  70% 13% 16%



- Molecule 2: Serine protease NS3

Chain H:  77% 10% 13%



- Molecule 3: peptide PAC-DLY-DLY-DAR

Chain I:  75% 25%



- Molecule 3: peptide PAC-DLY-DLY-DAR

Chain K:  25% 75%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.15Å 60.33Å 214.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	57.91 – 1.99 57.91 – 1.99	Depositor EDS
% Data completeness (in resolution range)	98.3 (57.91-1.99) 97.7 (57.91-1.99)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.59 (at 1.98Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.171 , 0.201 0.170 , 0.201	Depositor DCC
R_{free} test set	2609 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å ²)	20.8	Xtriage
Anisotropy	0.542	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 32.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.39$, $\langle L^2 \rangle = 0.22$	Xtriage
Estimated twinning fraction	0.417 for k,h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6187	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DAR, PAC, DLY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/292	0.65	0/397
1	C	0.23	0/284	0.61	0/385
1	E	0.23	0/289	0.62	0/393
1	G	0.22	0/297	0.61	0/404
2	B	0.28	0/1140	0.66	0/1550
2	D	0.26	0/1141	0.68	0/1552
2	F	0.26	0/1121	0.64	0/1526
2	H	0.26	0/1153	0.66	0/1564
All	All	0.26	0/5717	0.65	0/7771

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	288	0	258	4	0
1	C	280	0	261	3	0
1	E	285	0	256	3	0
1	G	293	0	260	3	0
2	B	1117	0	1082	25	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1118	0	1087	14	0
2	F	1098	0	1074	20	0
2	H	1131	0	1111	12	0
3	I	39	0	42	3	0
3	K	39	0	41	3	0
4	A	37	0	0	0	0
4	B	92	0	0	5	0
4	C	37	0	0	0	0
4	D	80	0	0	1	0
4	E	33	0	0	0	0
4	F	87	0	0	4	0
4	G	32	0	0	0	0
4	H	94	0	0	0	0
4	I	6	0	0	0	0
4	K	1	0	0	0	0
All	All	6187	0	5472	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 70 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:156:ILE:O	4:B:301:HOH:O	2.11	0.69
1:E:51:MET:HG2	2:F:27:THR:HG22	1.75	0.67
2:D:64:ARG:NH1	2:D:66:ASP:OD1	2.24	0.66
1:G:51:MET:HE1	2:H:53:THR:HA	1.77	0.64
2:B:101:PRO:HA	2:B:134:THR:HG22	1.78	0.63

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	36/53 (68%)	35 (97%)	1 (3%)	0	100	100
1	C	35/53 (66%)	35 (100%)	0	0	100	100
1	E	36/53 (68%)	33 (92%)	3 (8%)	0	100	100
1	G	37/53 (70%)	37 (100%)	0	0	100	100
2	B	149/178 (84%)	143 (96%)	6 (4%)	0	100	100
2	D	150/178 (84%)	145 (97%)	4 (3%)	1 (1%)	18	9
2	F	146/178 (82%)	142 (97%)	4 (3%)	0	100	100
2	H	148/178 (83%)	145 (98%)	3 (2%)	0	100	100
All	All	737/924 (80%)	715 (97%)	21 (3%)	1 (0%)	48	41

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	14	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	30/45 (67%)	30 (100%)	0	100	100
1	C	29/45 (64%)	29 (100%)	0	100	100
1	E	29/45 (64%)	29 (100%)	0	100	100
1	G	30/45 (67%)	30 (100%)	0	100	100
2	B	113/140 (81%)	112 (99%)	1 (1%)	70	67
2	D	113/140 (81%)	113 (100%)	0	100	100
2	F	112/140 (80%)	110 (98%)	2 (2%)	51	46
2	H	115/140 (82%)	113 (98%)	2 (2%)	53	48
All	All	571/740 (77%)	566 (99%)	5 (1%)	70	67

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	155	VAL
2	F	29	ARG
2	F	123	ILE
2	H	31	LEU
2	H	135	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	42	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	38/53 (71%)	-1.28	0 100 100	16, 21, 30, 38	0
1	C	37/53 (69%)	-1.24	0 100 100	20, 23, 28, 31	0
1	E	38/53 (71%)	-1.19	0 100 100	17, 24, 29, 33	0
1	G	39/53 (73%)	-1.19	0 100 100	18, 22, 34, 41	0
2	B	153/178 (85%)	-1.31	0 100 100	13, 19, 32, 43	0
2	D	154/178 (86%)	-1.26	0 100 100	15, 20, 34, 45	0
2	F	150/178 (84%)	-1.28	0 100 100	14, 20, 32, 41	0
2	H	154/178 (86%)	-1.31	0 100 100	12, 18, 29, 35	0
3	I	0/4	-	-	-	-
3	K	0/4	-	-	-	-
All	All	763/932 (81%)	-1.28	0 100 100	12, 20, 32, 45	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	DLY	I	3	9/10	0.98	0.04	17,23,25,29	0
3	DAR	K	4	12/12	0.98	0.04	19,25,31,32	0
3	DLY	I	2	9/10	0.99	0.04	22,23,25,26	0
3	DLY	K	3	9/10	0.99	0.03	11,17,23,25	0
3	DAR	I	4	12/12	0.99	0.04	12,19,31,31	0
3	DLY	K	2	9/10	0.99	0.04	17,20,28,29	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.