



wwPDB EM Validation Summary Report ⓘ

Mar 9, 2026 – 10:17 PM UTC

PDB ID : 3ZN8 / pdb_00003zn8
EMDB ID : EMD-2316
Title : Structural Basis of Signal Sequence Surveillance and Selection by the SRP-SR Complex
Authors : von Loeffelholz, O.; Knoop, K.; Ariosa, A.; Zhang, X.; Karuppasamy, M.; Huard, K.; Schoehn, G.; Berger, I.; Shan, S.O.; Schaffitzel, C.
Deposited on : 2013-02-13
Resolution : 12.00 Å (reported)
Based on initial models : 3KL4, 1FFH, 1FTS, 2XXA

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMD archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 12.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
RNA backbone	8273	3508	-
Q-score	-	25397	93 (11.50 - 12.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	294	46% 59% 38% .
2	D	295	47% 55% 44% .
3	G	88	8% 42% 55% .

Continued on next page...

Mol	Chain	Length	Quality of chain
4	M	125	42% 30% 50% 15%
5	S	14	57% 64% 29% 7%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7453 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SIGNAL RECOGNITION PARTICLE PROTEIN.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	288	Total	C	N	O	S	0	1
			2215	1391	406	413	5		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	48	THR	ALA	conflict	UNP O07347

- Molecule 2 is a protein called SIGNAL RECOGNITION PARTICLE RECEPTOR FTSY.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	D	295	Total	C	N	O	S	0	0
			2261	1430	394	431	6		

- Molecule 3 is a RNA chain called 4.5 S RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	G	88	Total	C	N	O	P	0	0
			1886	840	346	613	87		

- Molecule 4 is a protein called SIGNAL RECOGNITION PARTICLE 54 KDA PROTEIN.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	M	106	Total	C	N	O	S	0	0
			861	547	153	155	6		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	307	LEU	-	expression tag	UNP Q97ZE7

- Molecule 5 is a protein called DIPEPTIDYL AMINOPEPTIDASE B.

Mol	Chain	Residues	Atoms				AltConf	Trace
5	S	14	Total	C	N	O	0	0
			108	77	15	16		


- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
6	A	1	Total	Mg	0
			1	1	


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		AltConf
7	A	121	Total	O	0
			121	121	


- Molecule 3: 4.5 S RNA



- Molecule 4: SIGNAL RECOGNITION PARTICLE 54 KDA PROTEIN



- Molecule 5: DIPEPTIDYL AMINOPEPTIDASE B



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	46945	Depositor
Resolution determination method	Not provided	
CTF correction method	MICROGRAPH	Depositor
Microscope	FEI POLARA 300	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	15	Depositor
Minimum defocus (nm)	700	Depositor
Maximum defocus (nm)	5700	Depositor
Magnification	76000	Depositor
Image detector	GATAN ULTRASCAN 4000 (4k x 4k)	Depositor
Maximum map value	0.001	Depositor
Minimum map value	-0.001	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.000	Depositor
Recommended contour level	0.00015	Depositor
Map size ($\text{\AA}$)	562.5, 562.5, 562.5	wwPDB
Map dimensions	150, 150, 150	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	3.75, 3.75, 3.75	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.30	1/2238 (0.0%)	0.69	0/3013
2	D	0.25	0/2287	0.66	0/3078
3	G	0.25	0/2109	0.48	0/3290
4	M	0.27	0/872	0.69	0/1170
5	S	0.38	0/109	0.68	0/148
All	All	0.27	1/7615 (0.0%)	0.62	0/10699

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	271	GLY	C-N	-5.72	1.25	1.33

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2215	0	2313	93	0
2	D	2261	0	2333	124	0
3	G	1886	0	956	40	0
4	M	861	0	907	98	0
5	S	108	0	128	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	1	0	0	0	0
7	A	121	0	0	0	0
All	All	7453	0	6637	344	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 344 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:G:45:G:H1	3:G:54:C:H42	1.09	0.94
2:D:388:GLN:H	2:D:388:GLN:NE2	1.68	0.92
2:D:400:ILE:HG22	2:D:404:MET:HE2	1.52	0.91
1:A:28:LYS:HB3	1:A:32:ARG:HH12	1.38	0.88
4:M:343:GLY:HA3	4:M:344:PRO:C	1.99	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	284/294 (97%)	264 (93%)	18 (6%)	2 (1%)	18	56
2	D	293/295 (99%)	256 (87%)	36 (12%)	1 (0%)	36	72
4	M	103/125 (82%)	78 (76%)	21 (20%)	4 (4%)	2	19
5	S	12/14 (86%)	10 (83%)	2 (17%)	0	100	100
All	All	692/728 (95%)	608 (88%)	77 (11%)	7 (1%)	15	49

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	M	344	PRO
1	A	262	VAL
4	M	342	MET
1	A	97	ASP
4	M	345	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/233 (97%)	227 (100%)	0	100	100
2	D	238/238 (100%)	233 (98%)	5 (2%)	47	65
4	M	95/112 (85%)	91 (96%)	4 (4%)	26	48
5	S	12/12 (100%)	11 (92%)	1 (8%)	10	30
All	All	572/595 (96%)	562 (98%)	10 (2%)	52	69

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	M	344	PRO
4	M	364	GLU
5	S	457	TRP
2	D	345	GLN
2	D	388	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	388	GLN
4	M	423	ASN
4	M	382	ASN
2	D	319	GLN
2	D	355	HIS

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
3	G	87/88 (98%)	8 (9%)	0

5 of 8 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
3	G	13	C
3	G	37	U
3	G	49	G
3	G	72	C
3	G	73	U

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-2316. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 75



Y Index: 75




Z Index: 75


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 68



Y Index: 72




Z Index: 70


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.00015. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 3114 nm^3 ; this corresponds to an approximate mass of 2813 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)



*Reported resolution corresponds to spatial frequency of 0.083 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-2316 and PDB model 3ZN8. Per-residue inclusion information can be found in section [3](#) on page [6](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.00015 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.00015).

9.4 Atom inclusion ⓘ



At the recommended contour level, 56% of all backbone atoms, 59% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.00015) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.5880	0.0420
A	0.4780	0.0250
D	0.4830	0.0540
G	0.8970	0.0540
M	0.5130	0.0270
S	0.3080	0.0290

