



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 01:49 PM UTC

PDB ID : 3ZNR / pdb_00003znr
Title : HDAC7 bound with inhibitor TMP269
Authors : Lobera, m.; madauss, k.; pohlhaus, d.; trump, r.; nolan, m.
Deposited on : 2013-02-15
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	423	79% 8% 13%
1	B	423	3% 82% 7% 10%
1	C	423	76% 9% 15%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8454 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HISTONE DEACETYLASE 7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	368	Total	C	N	O	S	0	1	0
			2778	1744	499	516	19			
1	B	381	Total	C	N	O	S	0	0	0
			2872	1801	515	537	19			
1	C	359	Total	C	N	O	S	0	1	0
			2712	1696	493	505	18			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	481	GLY	-	expression tag	UNP Q8WUI4
B	481	GLY	-	expression tag	UNP Q8WUI4
C	481	GLY	-	expression tag	UNP Q8WUI4

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		
2	C	2	Total	Zn	0	0
			2	2		

- Molecule 3 is POTASSIUM ION (CCD ID: K) (formula: K).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	K	0	0
			2	2		
3	B	2	Total	K	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	2	Total K 2 2	0	0

- Molecule 4 is N-{{[4-(4-phenyl-1,3-thiazol-2-yl)tetrahydro-2H-pyran-4-yl]methyl}}-3-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide (CCD ID: NU9) (formula: C₂₅H₂₁F₃N₄O₃S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C F N O S 36 25 3 4 3 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	16	Total O 16 16	0	0
5	B	21	Total O 21 21	0	0
5	C	7	Total O 7 7	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: HISTONE DEACETYLASE 7



• Molecule 1: HISTONE DEACETYLASE 7



• Molecule 1: HISTONE DEACETYLASE 7



A831	
L837	
V856	
L859	
W872	
L878	
I888	
H891	
M898	
Q899	
ARG	
LEU	
ALA	
SER	

4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	81.39Å 81.39Å 149.26Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	70.49 – 2.40 70.49 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.3 (70.49-2.40) 99.3 (70.49-2.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.212 , 0.241 0.255 , 0.276	Depositor DCC
R_{free} test set	2178 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	45.6	Xtriage
Anisotropy	0.006	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 14.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.007 for -h,-k,l 0.043 for h,-h-k,-l 0.025 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8454	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NU9, K, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/2849	0.67	0/3869
1	B	0.40	0/2939	0.66	0/3988
1	C	0.40	0/2781	0.67	0/3778
All	All	0.40	0/8569	0.66	0/11635

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2778	0	2657	18	0
1	B	2872	0	2753	16	0
1	C	2712	0	2583	17	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
4	A	36	0	21	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	16	0	0	0	0
5	B	21	0	0	0	0
5	C	7	0	0	0	0
All	All	8454	0	8014	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:523:ILE:HG21	1:C:644:SER:HB3	1.66	0.77
1:C:623:VAL:HG22	1:C:627:THR:HB	1.74	0.69
1:C:888:ILE:HA	1:C:898:MET:HE3	1.75	0.67
1:B:623:VAL:HG22	1:B:627:THR:HB	1.77	0.66
1:A:623:VAL:HG22	1:A:627:THR:HB	1.77	0.65
1:B:523:ILE:HG21	1:B:644:SER:HB3	1.79	0.64
1:A:523:ILE:HG21	1:A:644:SER:HB3	1.80	0.64
1:C:661:PHE:CE2	1:C:663:VAL:HG22	2.33	0.63
1:C:692:LEU:HD21	1:C:793:LEU:HD13	1.82	0.61
1:C:555:LEU:HD23	1:C:560:LEU:HD12	1.84	0.59
1:A:744:VAL:HG12	1:A:890:VAL:HG21	1.86	0.58
1:A:520:THR:HG23	1:A:860:LEU:HD23	1.86	0.57
1:B:520:THR:HG22	1:B:659:ASN:OD1	2.06	0.56
1:B:808:ALA:HB3	1:B:809:PRO:HD3	1.87	0.55
1:A:519:THR:HG22	1:A:520:THR:H	1.71	0.55
1:A:824:THR:HG21	1:A:859:LEU:HD13	1.89	0.53
1:B:548:ILE:HD13	1:B:665:ARG:HG2	1.91	0.51
1:C:567:LEU:HD11	1:C:651:LYS:HE3	1.93	0.50
1:C:548:ILE:HD13	1:C:665:ARG:HG2	1.94	0.50
1:A:765:LEU:HD11	1:A:808:ALA:HB2	1.93	0.50
1:C:757:VAL:HG22	1:C:891:HIS:CE1	2.47	0.50
1:A:795:LEU:HD23	1:A:836:VAL:HB	1.95	0.49
1:C:529:LEU:HD23	1:C:548:ILE:HD11	1.95	0.48
1:C:708:VAL:HG12	1:C:801:ASP:CG	2.39	0.48
1:A:650:PHE:CE1	1:A:692:LEU:HD12	2.49	0.47
1:B:649:ALA:HB1	1:B:692:LEU:HD21	1.97	0.47
1:B:628:ILE:C	1:B:628:ILE:HD12	2.41	0.46
1:C:796:VAL:HB	1:C:837:LEU:HD23	1.97	0.46
1:B:708:VAL:HG12	1:B:801:ASP:CG	2.41	0.45
1:C:555:LEU:HD21	1:C:856:VAL:HG21	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:577:LEU:HD11	1:A:639:ARG:HG2	1.97	0.45
1:A:520:THR:HG22	1:A:659:ASN:OD1	2.15	0.45
1:C:541:HIS:CE1	1:C:628:ILE:HD11	2.52	0.45
1:A:548:ILE:HD13	1:A:665:ARG:HG2	1.97	0.45
1:B:577:LEU:HD23	1:B:687:ILE:HD13	1.98	0.45
1:B:796:VAL:HB	1:B:837:LEU:HD22	1.99	0.44
1:A:725:LEU:HD21	1:A:785:ILE:HG22	2.00	0.44
1:B:659:ASN:HB2	1:B:859:LEU:HD22	2.01	0.43
1:B:548:ILE:HB	1:B:663:VAL:HG12	2.01	0.42
1:C:820:PHE:HA	1:C:823:MET:HE3	2.01	0.42
1:C:792:ASP:O	1:C:831:ALA:HB1	2.20	0.42
1:C:775:LEU:HD11	1:C:872:TRP:CD2	2.55	0.41
1:A:688:ALA:O	1:A:692:LEU:HD13	2.21	0.41
1:A:672:ASP:OD1	1:A:672:ASP:C	2.63	0.41
1:B:616:LEU:HD22	1:B:625:THR:HA	2.01	0.41
1:A:574:LEU:HD21	1:A:591:THR:HG22	2.03	0.40
1:A:708:VAL:HG12	1:A:801:ASP:CG	2.46	0.40
1:B:688:ALA:O	1:B:692:LEU:HD13	2.21	0.40
1:B:761:TRP:CE3	1:B:769:MET:CE	3.04	0.40
1:B:593:PRO:HA	1:B:594:LEU:HA	1.80	0.40
1:A:818:LYS:HE2	1:A:864:VAL:HG13	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	365/423 (86%)	354 (97%)	10 (3%)	1 (0%)	36	50
1	B	377/423 (89%)	365 (97%)	11 (3%)	1 (0%)	36	50
1	C	356/423 (84%)	343 (96%)	12 (3%)	1 (0%)	36	50
All	All	1098/1269 (86%)	1062 (97%)	33 (3%)	3 (0%)	36	50

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	811	GLY
1	B	811	GLY
1	C	811	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	289/336 (86%)	284 (98%)	5 (2%)	53	74
1	B	298/336 (89%)	292 (98%)	6 (2%)	48	70
1	C	281/336 (84%)	276 (98%)	5 (2%)	51	73
All	All	868/1008 (86%)	852 (98%)	16 (2%)	51	73

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	519	THR
1	A	623	VAL
1	A	655	ARG
1	A	697	LYS
1	A	859	LEU
1	B	523	ILE
1	B	647	ASP
1	B	765	LEU
1	B	769	MET
1	B	839	LEU
1	B	889	ARG
1	C	554	ARG
1	C	567	LEU
1	C	765	LEU
1	C	859	LEU
1	C	878	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (24)

such sidechains are listed below:

Mol	Chain	Res	Type
1	A	636	ASN
1	A	673	HIS
1	A	691	GLN
1	A	732	HIS
1	A	736	ASN
1	A	756	ASN
1	A	758	ASN
1	A	843	HIS
1	A	879	ASN
1	A	891	HIS
1	B	610	GLN
1	B	636	ASN
1	B	691	GLN
1	B	826	GLN
1	B	874	GLN
1	B	879	ASN
1	B	899	GLN
1	C	636	ASN
1	C	673	HIS
1	C	695	GLN
1	C	736	ASN
1	C	758	ASN
1	C	826	GLN
1	C	879	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 13 ligands modelled in this entry, 12 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NU9	A	1000	2	39,40,40	4.04	11 (28%)	53,58,58	3.67	18 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NU9	A	1000	2	-	11/30/40/40	1/5/5/5

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1000	NU9	C28-N27	13.97	1.63	1.39
4	A	1000	NU9	C26-N27	10.34	1.48	1.30
4	A	1000	NU9	C26-S30	-9.47	1.54	1.73
4	A	1000	NU9	C29-C28	7.05	1.44	1.36
4	A	1000	NU9	C05-N06	6.96	1.35	1.28
4	A	1000	NU9	C31-C28	-5.34	1.39	1.47
4	A	1000	NU9	C10-C07	-5.03	1.39	1.47
4	A	1000	NU9	O09-C05	4.26	1.38	1.33
4	A	1000	NU9	C16-N18	4.21	1.43	1.33
4	A	1000	NU9	C29-S30	-4.06	1.61	1.71
4	A	1000	NU9	C07-N08	3.70	1.36	1.30

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1000	NU9	O09-C05-N06	-11.03	108.16	115.25
4	A	1000	NU9	N06-C07-N08	-9.50	107.54	114.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1000	NU9	O09-C05-C02	9.16	126.07	116.75
4	A	1000	NU9	O09-N08-C07	8.56	108.29	103.28
4	A	1000	NU9	C21-C20-C26	8.25	118.20	110.08
4	A	1000	NU9	C05-N06-C07	7.00	108.30	101.53
4	A	1000	NU9	C29-C28-N27	-6.50	107.06	114.53
4	A	1000	NU9	C25-C20-C26	6.16	116.14	110.08
4	A	1000	NU9	C05-O09-N08	5.57	107.65	105.34
4	A	1000	NU9	C10-C07-N08	5.17	126.63	122.22
4	A	1000	NU9	C25-C20-C19	-3.32	102.65	109.39
4	A	1000	NU9	C19-C20-C26	-3.11	101.54	106.80
4	A	1000	NU9	C31-C28-N27	2.94	124.45	119.95
4	A	1000	NU9	C21-C20-C19	-2.89	103.52	109.39
4	A	1000	NU9	C25-C20-C21	2.87	111.87	108.16
4	A	1000	NU9	C28-C29-S30	-2.39	108.27	111.16
4	A	1000	NU9	O23-C22-C21	2.27	113.77	111.50
4	A	1000	NU9	C20-C19-N18	2.23	117.09	114.17

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1000	NU9	C19-C20-C26-S30
4	A	1000	NU9	C21-C20-C26-S30
4	A	1000	NU9	C13-C14-C16-N18
4	A	1000	NU9	C13-C14-C16-O17
4	A	1000	NU9	C15-C14-C16-N18
4	A	1000	NU9	C15-C14-C16-O17
4	A	1000	NU9	C19-C20-C26-N27
4	A	1000	NU9	C21-C20-C26-N27
4	A	1000	NU9	C25-C20-C26-N27
4	A	1000	NU9	C25-C20-C26-S30
4	A	1000	NU9	C20-C19-N18-C16


All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1000	NU9	C20-C21-C22-C24-C25-O23

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	368/423 (86%)	0.20	4 (1%) 78 74	14, 22, 27, 31	1 (0%)
1	B	381/423 (90%)	0.33	11 (2%) 53 50	16, 22, 28, 35	0
1	C	359/423 (84%)	0.23	1 (0%) 90 88	11, 23, 26, 28	1 (0%)
All	All	1108/1269 (87%)	0.25	16 (1%) 73 69	11, 23, 27, 35	2 (0%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	594	LEU	4.6
1	B	593	PRO	4.2
1	B	596	ARG	4.0
1	B	560	LEU	3.2
1	A	808	ALA	2.6
1	A	809	PRO	2.6
1	B	734	ASP	2.5
1	B	517	PRO	2.4
1	B	870	GLU	2.3
1	B	592	ASN	2.3
1	C	564	CYS	2.2
1	B	768	PRO	2.1
1	A	614	VAL	2.1
1	B	597	LEU	2.1
1	A	870	GLU	2.0
1	B	695	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates


There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	K	A	202	1/1	0.35	0.26	80,80,80,80	0
3	K	C	202	1/1	0.52	0.17	73,73,73,73	0
4	NU9	A	1000	36/36	0.84	0.17	67,68,69,69	0
3	K	C	201	1/1	0.87	0.08	45,45,45,45	0
3	K	B	202	1/1	0.88	0.09	42,42,42,42	0
3	K	A	201	1/1	0.92	0.06	36,36,36,36	0
2	ZN	C	102	1/1	0.92	0.15	21,21,21,21	0
2	ZN	C	101	1/1	0.97	0.03	27,27,27,27	0
2	ZN	A	102	1/1	0.98	0.06	24,24,24,24	0
2	ZN	B	101	1/1	0.98	0.11	32,32,32,32	0
2	ZN	A	101	1/1	0.98	0.03	28,28,28,28	0
3	K	B	201	1/1	0.98	0.07	38,38,38,38	0
2	ZN	B	102	1/1	0.99	0.04	19,19,19,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.