



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 10:20 PM UTC

PDB ID : 6ZNJ / pdb_00006znj
Title : MaeB full-length enzyme apoprotein form
Authors : Lovering, A.L.; Harding, C.J.
Deposited on : 2020-07-06
Resolution : 3.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1131 (3.80-3.60)
Clashscore	190562	1171 (3.80-3.60)
Ramachandran outliers	187476	1129 (3.80-3.60)
Sidechain outliers	187428	1126 (3.80-3.60)
RSRZ outliers	180081	1130 (3.80-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	791	% 62% 32% • •
1	B	791	% 64% 29% • •
1	C	791	% 65% 29% • •

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 17400 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NADP-dependent malate dehydrogenase, Malate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	761	Total	C	N	O	S	0	2	0
			5800	3684	992	1096	28			
1	C	761	Total	C	N	O	S	0	2	0
			5800	3684	992	1096	28			
1	A	761	Total	C	N	O	S	0	2	0
			5800	3684	992	1096	28			


There are 45 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-10	MET	-	initiating methionine	UNP Q6MM14
B	-9	GLY	-	expression tag	UNP Q6MM14
B	-8	SER	-	expression tag	UNP Q6MM14
B	-7	SER	-	expression tag	UNP Q6MM14
B	-6	HIS	-	expression tag	UNP Q6MM14
B	-5	HIS	-	expression tag	UNP Q6MM14
B	-4	HIS	-	expression tag	UNP Q6MM14
B	-3	HIS	-	expression tag	UNP Q6MM14
B	-2	HIS	-	expression tag	UNP Q6MM14
B	-1	HIS	-	expression tag	UNP Q6MM14
B	0	SER	-	expression tag	UNP Q6MM14
B	121	ASP	-	linker	UNP Q6MM14
B	122	ILE	-	linker	UNP Q6MM14
B	123	GLU	-	linker	UNP Q6MM14
B	124	VAL	-	linker	UNP Q6MM14
C	-10	MET	-	initiating methionine	UNP Q6MM14
C	-9	GLY	-	expression tag	UNP Q6MM14
C	-8	SER	-	expression tag	UNP Q6MM14
C	-7	SER	-	expression tag	UNP Q6MM14
C	-6	HIS	-	expression tag	UNP Q6MM14
C	-5	HIS	-	expression tag	UNP Q6MM14
C	-4	HIS	-	expression tag	UNP Q6MM14

Continued on next page...


Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	-3	HIS	-	expression tag	UNP Q6MM14
C	-2	HIS	-	expression tag	UNP Q6MM14
C	-1	HIS	-	expression tag	UNP Q6MM14
C	0	SER	-	expression tag	UNP Q6MM14
C	121	ASP	-	linker	UNP Q6MM14
C	122	ILE	-	linker	UNP Q6MM14
C	123	GLU	-	linker	UNP Q6MM14
C	124	VAL	-	linker	UNP Q6MM14
A	-10	MET	-	initiating methionine	UNP Q6MM14
A	-9	GLY	-	expression tag	UNP Q6MM14
A	-8	SER	-	expression tag	UNP Q6MM14
A	-7	SER	-	expression tag	UNP Q6MM14
A	-6	HIS	-	expression tag	UNP Q6MM14
A	-5	HIS	-	expression tag	UNP Q6MM14
A	-4	HIS	-	expression tag	UNP Q6MM14
A	-3	HIS	-	expression tag	UNP Q6MM14
A	-2	HIS	-	expression tag	UNP Q6MM14
A	-1	HIS	-	expression tag	UNP Q6MM14
A	0	SER	-	expression tag	UNP Q6MM14
A	121	ASP	-	linker	UNP Q6MM14
A	122	ILE	-	linker	UNP Q6MM14
A	123	GLU	-	linker	UNP Q6MM14
A	124	VAL	-	linker	UNP Q6MM14



• Molecule 1: NADP-dependent malate dehydrogenase, Malate dehydrogenase

Chain A: 62% 32%





4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	81.70Å 273.58Å 308.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	66.77 – 3.70 66.77 – 3.70	Depositor EDS
% Data completeness (in resolution range)	96.5 (66.77-3.70) 96.6 (66.77-3.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.08 (at 3.68Å)	Xtriage
Refinement program	PHENIX v1.0	Depositor
R, R_{free}	0.210 , 0.270 0.226 , 0.287	Depositor DCC
R_{free} test set	1791 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å ²)	101.5	Xtriage
Anisotropy	0.946	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 50.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	17400	wwPDB-VP
Average B, all atoms (Å ²)	103.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.85	0/5909	1.22	28/8004 (0.3%)
1	B	0.82	2/5909 (0.0%)	1.19	27/8004 (0.3%)
1	C	0.80	1/5909 (0.0%)	1.15	18/8004 (0.2%)
All	All	0.83	3/17727 (0.0%)	1.19	73/24012 (0.3%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	456	ALA	CA-CB	-5.47	1.45	1.53
1	C	171	VAL	CA-CB	-5.44	1.47	1.54
1	B	610	LEU	CA-C	5.16	1.58	1.52

The worst 5 of 73 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	436	GLN	N-CA-C	-10.77	91.26	109.24
1	A	436	GLN	CB-CA-C	10.43	130.00	109.35
1	B	689	ASN	CA-C-N	9.12	128.85	119.64
1	B	689	ASN	C-N-CA	9.12	128.85	119.64
1	C	146	ILE	N-CA-C	7.91	119.48	107.77

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5800	0	5906	172	0
1	B	5800	0	5906	171	0
1	C	5800	0	5906	169	0
All	All	17400	0	17718	493	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 493 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:607:PHE:HE2	1:B:640:ILE:HD12	1.32	0.94
1:C:443:ILE:HD11	1:C:758:ASN:HB3	1.49	0.94
1:C:607:PHE:HE2	1:C:640:ILE:HD12	1.34	0.91
1:A:226:ARG:HH21	1:A:253:PHE:HD1	1.20	0.88
1:A:307:PRO:HA	1:A:319:ILE:HD13	1.55	0.87

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	761/791 (96%)	731 (96%)	30 (4%)	0	100	100
1	B	761/791 (96%)	731 (96%)	30 (4%)	0	100	100
1	C	761/791 (96%)	738 (97%)	23 (3%)	0	100	100
All	All	2283/2373 (96%)	2200 (96%)	83 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	617/642 (96%)	603 (98%)	14 (2%)	44	62
1	B	617/642 (96%)	595 (96%)	22 (4%)	31	54
1	C	617/642 (96%)	607 (98%)	10 (2%)	55	68
All	All	1851/1926 (96%)	1805 (98%)	46 (2%)	43	61

5 of 46 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	709	LEU
1	A	130	ASP
1	C	721	TYR
1	A	34	ILE
1	A	426	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	328	ASN
1	A	453	ASN
1	A	758	ASN
1	A	630	GLN
1	C	511	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	761/791 (96%)	-0.15	10 (1%) 75 50	62, 92, 121, 142	2 (0%)
1	B	761/791 (96%)	-0.13	5 (0%) 84 63	63, 100, 132, 164	2 (0%)
1	C	761/791 (96%)	-0.14	6 (0%) 82 60	68, 109, 143, 163	2 (0%)
All	All	2283/2373 (96%)	-0.14	21 (0%) 81 58	62, 100, 136, 164	6 (0%)

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	143	PHE	3.1
1	A	487	CYS	3.1
1	A	114	PHE	3.0
1	A	483	GLU	2.9
1	A	80	VAL	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.