



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 26, 2026 – 10:32 AM EDT

PDB ID : 7ZN6 / pdb_00007zn6
Title : Crystal structure of laccase-like multicopper oxidase (LMCO) from *Thermothelomyces thermophilus*
Authors : Kosinas, C.; Dimarogona, M.; Zerva, A.; Topakas, E.
Deposited on : 2022-04-20
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	654	
2	B	2	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 9432 atoms, of which 4455 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Laccase-like multicopper oxidase 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	576	Total	C	H	N	O	S	149	4	0
			8920	2885	4368	787	859	21			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLU	-	expression tag	UNP G2QFD0
A	2	PHE	-	expression tag	UNP G2QFD0
A	633	LEU	-	expression tag	UNP G2QFD0
A	634	GLU	-	expression tag	UNP G2QFD0
A	635	GLN	-	expression tag	UNP G2QFD0
A	636	LYS	-	expression tag	UNP G2QFD0
A	637	LEU	-	expression tag	UNP G2QFD0
A	638	ILE	-	expression tag	UNP G2QFD0
A	639	SER	-	expression tag	UNP G2QFD0
A	640	GLU	-	expression tag	UNP G2QFD0
A	641	GLU	-	expression tag	UNP G2QFD0
A	642	ASP	-	expression tag	UNP G2QFD0
A	643	LEU	-	expression tag	UNP G2QFD0
A	644	ASN	-	expression tag	UNP G2QFD0
A	645	SER	-	expression tag	UNP G2QFD0
A	646	ALA	-	expression tag	UNP G2QFD0
A	647	VAL	-	expression tag	UNP G2QFD0
A	648	ASP	-	expression tag	UNP G2QFD0
A	649	HIS	-	expression tag	UNP G2QFD0
A	650	HIS	-	expression tag	UNP G2QFD0
A	651	HIS	-	expression tag	UNP G2QFD0
A	652	HIS	-	expression tag	UNP G2QFD0
A	653	HIS	-	expression tag	UNP G2QFD0
A	654	HIS	-	expression tag	UNP G2QFD0

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	2	Total	C	H	N	O	5	0	0
			55	16	27	2	10			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	3	0
			28	8	14	1	5		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	1	0
			10	2	6	2		
4	A	1	Total	C	H	O	1	0
			10	2	6	2		
4	A	1	Total	C	H	O	1	0
			10	2	6	2		
4	A	1	Total	C	H	O	1	0
			10	2	6	2		
4	A	1	Total	C	H	O	1	0
			10	2	6	2		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C H O 14 3 8 3	2	0
5	A	1	Total C H O 14 3 8 3	2	0

- Molecule 6 is OXYGEN MOLECULE (CCD ID: OXY) (formula: O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total O 2 2	0	0

- Molecule 7 is COPPER (II) ION (CCD ID: CU) (formula: Cu).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	4	Total Cu 4 4	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	345	Total O 345 345	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Laccase-like multicopper oxidase 1



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	74.86Å 74.86Å 118.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	74.97 – 1.90 74.86 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (74.97-1.90) 100.0 (74.86-1.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.13 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.8.0267, REFMAC 5.8.0267	Depositor
R, R_{free}	0.152 , 0.185 0.152 , 0.186	Depositor DCC
R_{free} test set	2567 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	29.1	Xtriage
Anisotropy	0.551	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 39.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.043 for h,-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9432	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CU, GOL, NAG, OXY, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.08	6/4693 (0.1%)	1.26	6/6403 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	585	PRO	C-O	-6.60	1.15	1.23
1	A	491	HIS	CE1-NE2	6.36	1.39	1.32
1	A	483	HIS	CE1-NE2	6.22	1.38	1.32
1	A	332	PRO	C-O	-5.57	1.17	1.23
1	A	13	PRO	C-O	-5.25	1.17	1.23
1	A	120	MET	CG-SD	-5.04	1.68	1.80

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	270	ASN	CA-CB-CG	-6.70	105.90	112.60
1	A	330	PRO	N-CA-C	6.48	123.21	113.81
1	A	287	THR	CA-CB-OG1	-5.23	101.75	109.60
1	A	113	PHE	CA-CB-CG	5.21	119.01	113.80
1	A	281	GLU	CA-C-O	-5.20	115.51	120.92
1	A	281	GLU	CB-CG-CD	5.01	121.12	112.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	280	SER	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4552	4368	4343	20	0
2	B	28	27	25	0	0
3	A	14	14	13	0	0
4	A	20	30	30	3	0
5	A	12	16	16	1	0
6	A	2	0	0	0	0
7	A	4	0	0	0	0
8	A	345	0	0	4	0
All	All	4977	4455	4427	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:268:ASP:OD2	8:A:801:HOH:O	2.13	0.66
1:A:195:ARG:NH1	8:A:804:HOH:O	2.34	0.58
1:A:23:GLN:HB3	1:A:30[B]:ASN:OD1	2.04	0.58
1:A:432[B]:GLU:HG3	8:A:1056:HOH:O	2.04	0.56
1:A:46:HIS:ND1	1:A:132:GLU:OE2	2.37	0.54
1:A:334:ARG:HD2	4:A:705:EDO:H11	1.89	0.54
1:A:276:GLN:CD	1:A:279:GLY:O	2.52	0.52
1:A:388:ARG:HD2	1:A:442:LEU:HD22	1.95	0.49
1:A:281:GLU:HG2	1:A:519:ILE:HB	1.94	0.48
1:A:508:ASN:C	4:A:709:EDO:H21	2.39	0.48
1:A:276:GLN:HB3	1:A:300:ASP:HB2	1.97	0.47
1:A:148:LEU:HD23	1:A:148:LEU:C	2.41	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:586:LEU:HD22	1:A:588:VAL:HG22	1.98	0.45
1:A:410:ALA:HA	8:A:986:HOH:O	2.15	0.45
1:A:74:SER:HB3	5:A:706:GOL:H12	1.99	0.44
1:A:248:THR:OG1	1:A:304:ARG:HD2	2.18	0.43
1:A:436:PRO:HG2	1:A:455:PRO:HB3	2.00	0.43
1:A:509:ALA:N	4:A:709:EDO:H21	2.35	0.42
1:A:92:ILE:HG12	1:A:98:PHE:HB2	2.01	0.42
1:A:82:ASP:HB3	1:A:88:THR:HG21	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	576/654 (88%)	555 (96%)	21 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	488/549 (89%)	478 (98%)	10 (2%)	48	46

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	136	SER
1	A	195	ARG
1	A	231	GLU
1	A	235	ASP
1	A	287	THR
1	A	413	THR
1	A	432[A]	GLU
1	A	432[B]	GLU
1	A	531	GLU
1	A	586	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	292	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	2,1	14,14,15	0.53	0	17,19,21	1.29	3 (17%)
2	NAG	B	2	2	14,14,15	0.70	0	17,19,21	1.60	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	NAG	C4-C3-C2	3.65	116.36	111.02
2	B	2	NAG	O3-C3-C4	-3.21	102.81	110.38
2	B	1	NAG	C2-N2-C7	-2.48	119.57	122.90
2	B	2	NAG	O4-C4-C5	2.25	114.86	109.32
2	B	1	NAG	O5-C5-C4	-2.23	105.39	110.83
2	B	2	NAG	C1-C2-N2	-2.14	107.06	110.43
2	B	1	NAG	O5-C1-C2	-2.06	108.10	111.29


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 4 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	701	1	14,14,15	1.07	0	17,19,21	2.00	4 (23%)
4	EDO	A	709	-	3,3,3	0.27	0	2,2,2	0.26	0
5	GOL	A	704	-	5,5,5	0.23	0	5,5,5	0.49	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	A	702	-	3,3,3	0.31	0	2,2,2	0.26	0
4	EDO	A	703	-	3,3,3	0.42	0	2,2,2	0.34	0
4	EDO	A	705	-	3,3,3	0.38	0	2,2,2	0.60	0
5	GOL	A	706	-	5,5,5	0.16	0	5,5,5	0.40	0
4	EDO	A	708	-	3,3,3	0.07	0	2,2,2	0.31	0
6	OXY	A	707	7	1,1,1	0.36	0	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsions and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	701	1	-	0/6/23/26	0/1/1/1
4	EDO	A	709	-	-	0/1/1/1	-
5	GOL	A	704	-	-	2/4/4/4	-
4	EDO	A	702	-	-	1/1/1/1	-
4	EDO	A	703	-	-	1/1/1/1	-
4	EDO	A	705	-	-	1/1/1/1	-
5	GOL	A	706	-	-	4/4/4/4	-
4	EDO	A	708	-	-	1/1/1/1	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	701	NAG	C8-C7-N2	3.91	122.60	116.12
3	A	701	NAG	O7-C7-C8	-3.19	116.38	122.05
3	A	701	NAG	O5-C5-C6	3.15	113.79	107.66
3	A	701	NAG	C1-O5-C5	2.57	115.63	112.19

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	704	GOL	C1-C2-C3-O3
5	A	704	GOL	O2-C2-C3-O3
5	A	706	GOL	O2-C2-C3-O3
5	A	706	GOL	O1-C1-C2-C3
5	A	706	GOL	C1-C2-C3-O3
4	A	705	EDO	O1-C1-C2-O2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	A	706	GOL	O1-C1-C2-O2
4	A	702	EDO	O1-C1-C2-O2
4	A	708	EDO	O1-C1-C2-O2
4	A	703	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	709	EDO	2	0
4	A	705	EDO	1	0
5	A	706	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	576/654 (88%)	-0.29	6 (1%) 79 82	13, 35, 58, 87	4 (0%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	415	THR	3.4
1	A	416	ASP	2.4
1	A	602	ASN	2.3
1	A	134	CYS	2.2
1	A	135	GLY	2.2
1	A	417	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	B	2	14/15	0.88	0.10	30,62,71,73	3
2	NAG	B	1	14/15	0.95	0.06	30,49,54,55	2

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	GOL	A	704	6/6	0.77	0.17	30,54,62,66	2
4	EDO	A	705	4/4	0.83	0.15	30,49,55,57	1
4	EDO	A	708	4/4	0.84	0.15	30,66,69,69	1
4	EDO	A	702	4/4	0.84	0.16	30,54,60,62	1
4	EDO	A	709	4/4	0.85	0.15	30,66,72,73	1
4	EDO	A	703	4/4	0.91	0.12	30,54,59,60	1
5	GOL	A	706	6/6	0.92	0.12	30,63,66,67	2
3	NAG	A	701	14/15	0.93	0.08	30,40,47,48	3
6	OXY	A	707	2/2	0.98	0.09	32,32,32,38	0
7	CU	A	710	1/1	0.99	0.03	28,28,28,28	0
7	CU	A	711	1/1	0.99	0.02	31,31,31,31	0
7	CU	A	712	1/1	0.99	0.03	29,29,29,29	0
7	CU	A	713	1/1	0.99	0.02	28,28,28,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.