



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 12:02 PM UTC

PDB ID : 4ZO0 / pdb_00004zo0
Title : X-ray Structure of AAV-2 Origin Binding Domain
Authors : Musayev, F.N.; Zarate-Perez, F.; Escalante, C.R.
Deposited on : 2015-05-05
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	209	3% 75% 21% ••
1	B	209	27% 57% 31% 6% 5%
1	C	209	10% 72% 18% • 9%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5028 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Protein Rep68.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	202	Total	C	N	O	S	0	0	0
			1628	1049	275	299	5			
1	B	198	Total	C	N	O	S	0	0	0
			1549	1000	260	284	5			
1	C	191	Total	C	N	O	S	0	0	0
			1529	990	255	279	5			

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP P03132
A	-1	SER	-	expression tag	UNP P03132
A	0	HIS	-	expression tag	UNP P03132
A	17	GLU	GLY	engineered mutation	UNP P03132
A	151	SER	CYS	engineered mutation	UNP P03132
B	-2	GLY	-	expression tag	UNP P03132
B	-1	SER	-	expression tag	UNP P03132
B	0	HIS	-	expression tag	UNP P03132
B	17	GLU	GLY	engineered mutation	UNP P03132
B	151	SER	CYS	engineered mutation	UNP P03132
C	-2	GLY	-	expression tag	UNP P03132
C	-1	SER	-	expression tag	UNP P03132
C	0	HIS	-	expression tag	UNP P03132
C	17	GLU	GLY	engineered mutation	UNP P03132
C	151	SER	CYS	engineered mutation	UNP P03132

- Molecule 2 is ISOPROPYL ALCOHOL (CCD ID: IPA) (formula: C₃H₈O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	3	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		
3	C	1	Total	Mg	0	0
			1	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	187	Total	O	0	0
			187	187		
4	B	29	Total	O	0	0
			29	29		
4	C	99	Total	O	0	0
			99	99		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Protein Rep68

Chain A: 




• Molecule 1: Protein Rep68

Chain B: 



• Molecule 1: Protein Rep68

Chain C: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	186.36Å 154.40Å 38.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.72 – 2.30 29.72 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.8 (29.72-2.30) 99.7 (29.72-2.30)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.65 (at 2.31Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.225 , 0.242 0.230 , 0.242	Depositor DCC
R_{free} test set	2546 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	41.7	Xtriage
Anisotropy	0.376	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 45.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5028	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, IPA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.77	15/1670 (0.9%)	1.06	12/2274 (0.5%)
1	B	1.38	2/1591 (0.1%)	1.10	12/2177 (0.6%)
1	C	1.57	4/1569 (0.3%)	1.05	7/2137 (0.3%)
All	All	1.59	21/4830 (0.4%)	1.07	31/6588 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	51	ALA	C-O	-5.99	1.18	1.24
1	C	51	ALA	C-O	-5.96	1.18	1.24
1	C	120	ILE	C-O	-5.64	1.17	1.24
1	A	61	ARG	C-O	-5.54	1.17	1.24
1	A	18	HIS	C-O	-5.46	1.17	1.24
1	C	31	ALA	C-O	-5.44	1.17	1.24
1	A	11	VAL	C-O	-5.42	1.18	1.23
1	A	8	VAL	C-O	-5.32	1.18	1.24
1	A	117	ILE	C-O	-5.31	1.18	1.24
1	A	158	LEU	C-O	-5.26	1.19	1.24
1	C	25	SER	C-O	-5.24	1.18	1.24
1	B	74	PRO	N-CD	5.23	1.55	1.47
1	B	73	ALA	C-N	5.22	1.40	1.34
1	A	105	LEU	C-O	-5.19	1.18	1.24
1	A	135	THR	C-O	-5.15	1.17	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	147	VAL	C-O	-5.12	1.18	1.24
1	A	22	ILE	C-O	-5.11	1.18	1.24
1	A	102	SER	C-O	-5.02	1.18	1.24
1	A	7	ILE	C-O	-5.02	1.18	1.24
1	A	53	LEU	C-O	-5.02	1.18	1.24
1	A	114	GLU	C-O	-5.01	1.18	1.24

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	157	LEU	N-CA-C	11.37	123.42	111.14
1	A	119	ARG	N-CA-C	8.78	120.63	111.14
1	C	157	LEU	N-CA-C	8.52	120.34	111.14
1	C	15	LEU	N-CA-C	8.37	120.09	110.97
1	B	170	THR	N-CA-C	7.71	122.17	109.06
1	C	69	ARG	N-CA-C	7.47	119.21	111.14
1	A	18	HIS	N-CA-C	6.90	118.88	111.36
1	B	119	ARG	N-CA-C	6.73	118.27	111.07
1	C	50	GLN	N-CA-C	6.72	118.61	111.28
1	A	131	TRP	N-CA-C	6.70	119.41	111.71
1	B	104	VAL	N-CA-C	6.66	121.42	113.22
1	B	155	ASN	N-CA-C	6.59	118.54	111.36
1	A	155	ASN	N-CA-C	6.39	118.32	111.36
1	A	133	ALA	N-CA-C	6.20	118.56	108.26
1	A	104	VAL	N-CA-C	6.07	118.84	112.83
1	B	107	ARG	N-CA-C	-6.05	104.69	111.28
1	A	50	GLN	N-CA-C	5.88	117.69	111.28
1	B	15	LEU	N-CA-C	5.81	117.61	111.28
1	A	62	ASP	N-CA-C	5.67	117.54	111.36
1	B	171	ASN	N-CA-C	5.58	117.45	111.36
1	C	133	ALA	N-CA-C	5.53	117.42	108.34
1	A	70	VAL	N-CA-C	5.34	116.12	110.72
1	C	11	VAL	N-CA-C	5.34	113.97	107.61
1	A	17	GLU	N-CA-C	5.32	116.94	111.03
1	A	205	GLU	N-CA-C	-5.31	103.88	110.41
1	A	16	ASP	N-CA-C	5.14	116.97	111.36
1	C	131	TRP	N-CA-C	5.14	118.81	112.54
1	B	1	MET	CA-C-N	-5.10	113.47	119.84
1	B	1	MET	C-N-CA	-5.10	113.47	119.84
1	B	2	PRO	CA-N-CD	-5.09	104.87	112.00
1	B	11	VAL	N-CA-C	5.02	113.59	107.61

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	69	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1628	0	1586	10	0
1	B	1549	0	1462	71	0
1	C	1529	0	1492	16	0
2	A	4	0	8	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	A	187	0	0	0	0
4	B	29	0	0	1	0
4	C	99	0	0	1	0
All	All	5028	0	4548	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (97) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:18:HIS:O	1:B:19:LEU:HD23	1.49	1.12
1:B:137:THR:HG22	1:B:144:GLY:O	1.54	1.06
1:B:104:VAL:HG13	1:B:108:PHE:CE2	2.06	0.91
1:B:61:ARG:HH11	1:B:61:ARG:HG2	1.35	0.89
1:B:19:LEU:O	1:B:22:ILE:HG22	1.79	0.83
1:B:100:VAL:HG11	1:B:105:LEU:CD1	2.09	0.83
1:B:104:VAL:CG1	1:B:108:PHE:CE2	2.62	0.82
1:B:117:ILE:HD11	1:B:131:TRP:HZ2	1.45	0.82
1:B:18:HIS:C	1:B:19:LEU:HD23	2.09	0.76
1:B:51:ALA:HB3	1:B:52:PRO:HD3	1.68	0.75

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:137:THR:HG22	1:B:144:GLY:C	2.12	0.75
1:B:100:VAL:HG11	1:B:105:LEU:HD13	1.68	0.74
1:B:170:THR:HG23	1:B:176:LEU:HD12	1.69	0.73
1:B:104:VAL:HG13	1:B:108:PHE:CD2	2.24	0.72
1:C:158:LEU:N	1:C:159:PRO:HD2	2.04	0.72
1:B:71:SER:O	1:B:72:LYS:CB	2.39	0.71
1:B:58:LYS:HD3	1:B:61:ARG:HH22	1.54	0.71
1:A:51:ALA:HB3	1:A:52:PRO:HD3	1.74	0.69
1:B:172:MET:O	1:B:176:LEU:HD13	1.92	0.69
1:B:74:PRO:HG2	1:B:75:GLU:HG2	1.76	0.68
1:B:6:GLU:HG3	1:B:94:LEU:CD2	2.24	0.68
1:B:100:VAL:HG11	1:B:105:LEU:HD12	1.77	0.67
1:B:137:THR:HA	1:B:146:LYS:HB2	1.76	0.67
1:C:153:ILE:HB	1:C:154:PRO:CD	2.26	0.66
1:B:117:ILE:HD11	1:B:131:TRP:CZ2	2.32	0.65
1:B:61:ARG:HG2	1:B:61:ARG:NH1	2.03	0.65
1:C:185:ARG:O	1:C:189:VAL:HG23	1.97	0.64
1:B:77:LEU:HA	1:B:171:ASN:HD22	1.62	0.64
1:B:24:ASP:O	1:B:28:ASN:ND2	2.31	0.64
1:B:6:GLU:HG3	1:B:94:LEU:HD21	1.79	0.63
1:B:22:ILE:HG13	1:B:23:SER:N	2.13	0.63
1:B:105:LEU:O	1:B:108:PHE:HB2	1.99	0.63
1:B:104:VAL:HG11	1:B:108:PHE:CE2	2.36	0.60
1:B:58:LYS:CD	1:B:61:ARG:HH22	2.15	0.60
1:B:58:LYS:HD3	1:B:61:ARG:NH2	2.16	0.60
1:C:153:ILE:HB	1:C:154:PRO:HD2	1.85	0.59
1:B:5:TYR:CD2	1:B:97:THR:HG22	2.38	0.59
1:C:68:ARG:NH2	4:C:405:HOH:O	2.35	0.58
1:B:74:PRO:HB2	1:B:75:GLU:OE2	2.04	0.56
1:B:100:VAL:HG13	1:B:108:PHE:CD2	2.40	0.56
1:B:4:PHE:HB2	1:B:148:VAL:HG22	1.88	0.56
1:B:115:LYS:O	1:B:118:GLN:HG3	2.06	0.55
1:B:62:ASP:O	1:B:65:THR:OG1	2.24	0.54
1:B:40:ASP:OD1	1:B:40:ASP:N	2.24	0.54
1:B:6:GLU:HB3	1:B:135:THR:OG1	2.07	0.54
1:B:170:THR:CG2	1:B:176:LEU:HD12	2.38	0.53
1:B:67:TRP:O	1:B:70:VAL:HG12	2.09	0.53
1:A:196:VAL:O	1:A:200:GLN:HG3	2.09	0.53
1:B:76:ALA:O	1:B:171:ASN:ND2	2.42	0.52
1:A:1:MET:N	1:A:2:PRO:HD2	2.24	0.52
1:B:100:VAL:CG1	1:B:105:LEU:HD13	2.38	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1:MET:N	1:B:2:PRO:CD	2.73	0.52
1:B:137:THR:HG22	1:B:144:GLY:H	1.75	0.52
1:C:158:LEU:N	1:C:159:PRO:CD	2.73	0.52
1:C:79:PHE:CD2	1:C:170:THR:HB	2.45	0.52
1:B:109:LEU:O	1:B:112:ILE:N	2.43	0.51
1:B:18:HIS:O	1:B:127:THR:HG23	2.10	0.51
1:B:6:GLU:HG3	1:B:94:LEU:HD23	1.93	0.51
1:B:15:LEU:C	1:B:15:LEU:HD23	2.36	0.50
1:B:1:MET:H2	1:B:2:PRO:CD	2.25	0.50
1:B:113:ARG:O	1:B:116:LEU:N	2.44	0.50
1:A:160:LYS:NZ	1:A:164:GLU:OE2	2.45	0.49
1:B:61:ARG:NH1	1:B:61:ARG:CG	2.73	0.49
1:C:150:GLU:OE2	1:C:175:TYR:OH	2.23	0.48
1:C:3:GLY:O	1:C:97:THR:HG23	2.13	0.48
1:A:26:PHE:C	1:A:26:PHE:CD1	2.92	0.48
1:B:105:LEU:HD23	1:B:145:ASN:ND2	2.28	0.47
1:B:100:VAL:HG13	1:B:108:PHE:CE2	2.50	0.47
1:B:138:ARG:NH1	1:B:143:GLY:HA2	2.30	0.46
1:B:137:THR:HG22	1:B:144:GLY:N	2.30	0.46
1:B:67:TRP:O	1:B:70:VAL:CG1	2.64	0.46
1:A:158:LEU:N	1:A:159:PRO:CD	2.79	0.46
1:B:80:VAL:HG22	1:B:93:VAL:HG13	1.98	0.45
1:B:124:ILE:HA	4:B:408:HOH:O	2.15	0.45
1:B:124:ILE:O	1:B:126:PRO:HD3	2.16	0.45
1:A:71:SER:O	1:A:72:LYS:HB2	2.16	0.45
1:B:154:PRO:HA	1:B:158:LEU:HD12	1.99	0.44
1:A:58:LYS:HE3	1:A:120:ILE:HG13	1.99	0.44
1:C:51:ALA:HB3	1:C:52:PRO:HD3	1.99	0.44
1:B:113:ARG:O	1:B:116:LEU:HB2	2.17	0.44
1:C:44:ASP:HB3	1:C:47:LEU:HD12	1.99	0.44
1:B:104:VAL:CG1	1:B:108:PHE:HE2	2.23	0.43
1:B:51:ALA:N	1:B:52:PRO:CD	2.80	0.43
1:C:78:PHE:CD1	1:C:78:PHE:C	2.96	0.43
1:A:150:GLU:HG2	1:A:150:GLU:O	2.17	0.43
1:B:1:MET:N	1:B:2:PRO:HD3	2.32	0.43
1:B:78:PHE:H	1:B:171:ASN:HB2	1.83	0.43
1:C:41:SER:OG	1:C:43:MET:HB2	2.19	0.43
1:B:63:PHE:CD1	1:B:63:PHE:C	2.97	0.42
1:C:153:ILE:CB	1:C:154:PRO:CD	2.94	0.42
1:B:146:LYS:HG3	1:B:147:VAL:N	2.35	0.42
1:C:187:ARG:HG2	1:C:188:LEU:HD12	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:64:LEU:O	1:B:68:ARG:HG3	2.20	0.41
1:B:64:LEU:O	1:B:67:TRP:N	2.54	0.41
1:C:1:MET:N	1:C:2:PRO:CD	2.83	0.41
1:A:125:GLU:HA	1:A:126:PRO:HD2	1.91	0.41
1:B:158:LEU:N	1:B:159:PRO:CD	2.84	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	198/209 (95%)	197 (100%)	1 (0%)	0	100	100
1	B	194/209 (93%)	190 (98%)	4 (2%)	0	100	100
1	C	187/209 (90%)	186 (100%)	1 (0%)	0	100	100
All	All	579/627 (92%)	573 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	176/188 (94%)	172 (98%)	4 (2%)	44	63
1	B	161/188 (86%)	149 (92%)	12 (8%)	12	17

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	163/188 (87%)	156 (96%)	7 (4%)	26	39
All	All	500/564 (89%)	477 (95%)	23 (5%)	24	36

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	46	ASN
1	A	49	GLU
1	A	91	MET
1	A	114	GLU
1	B	22	ILE
1	B	40	ASP
1	B	87	SER
1	B	91	MET
1	B	95	VAL
1	B	113	ARG
1	B	120	ILE
1	B	125	GLU
1	B	146	LYS
1	B	148	VAL
1	B	174	GLN
1	B	176	LEU
1	C	72	LYS
1	C	86	GLU
1	C	107	ARG
1	C	125	GLU
1	C	137	THR
1	C	149	ASP
1	C	150	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	145	ASN
1	B	90	HIS
1	B	92	HIS
1	B	145	ASN
1	B	171	ASN
1	B	200	GLN
1	C	90	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	IPA	A	401	-	3,3,3	0.57	0	3,3,3	0.31	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	202/209 (96%)	-0.02	6 (2%) 52 54	26, 36, 57, 75	0
1	B	198/209 (94%)	1.44	57 (28%) 1 1	42, 64, 85, 91	0
1	C	191/209 (91%)	0.52	20 (10%) 11 13	29, 46, 74, 85	0
All	All	591/627 (94%)	0.64	83 (14%) 6 7	26, 48, 81, 91	0

All (83) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	143	GLY	6.1
1	B	104	VAL	5.2
1	B	112	ILE	4.9
1	C	142	GLY	4.9
1	B	1	MET	4.9
1	B	108	PHE	4.8
1	A	138	ARG	4.7
1	B	201	GLU	4.7
1	A	206	ASN	4.4
1	C	141	ALA	4.2
1	B	64	LEU	4.2
1	C	1	MET	4.2
1	C	139	ASN	4.1
1	B	139	ASN	4.1
1	B	65	THR	4.0
1	B	111	GLN	4.0
1	B	70	VAL	3.9
1	A	1	MET	3.9
1	B	77	LEU	3.9
1	B	67	TRP	3.8
1	B	71	SER	3.7
1	B	2	PRO	3.7
1	B	114	GLU	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	98	THR	3.6
1	A	143	GLY	3.6
1	B	66	GLU	3.6
1	B	118	GLN	3.5
1	B	117	ILE	3.4
1	B	109	LEU	3.3
1	B	116	LEU	3.3
1	B	124	ILE	3.3
1	B	32	GLU	3.2
1	B	102	SER	3.2
1	B	96	GLU	3.0
1	B	110	SER	3.0
1	C	138	ARG	3.0
1	C	2	PRO	3.0
1	C	0	HIS	3.0
1	C	155	ASN	3.0
1	B	119	ARG	2.9
1	B	103	MET	2.9
1	B	107	ARG	2.9
1	C	183	THR	2.9
1	B	100	VAL	2.9
1	A	2	PRO	2.8
1	B	115	LYS	2.7
1	B	105	LEU	2.7
1	B	106	GLY	2.6
1	B	5	TYR	2.6
1	C	151	SER	2.5
1	C	191	GLN	2.5
1	B	21	GLY	2.5
1	B	69	ARG	2.4
1	B	101	LYS	2.4
1	B	78	PHE	2.4
1	C	40	ASP	2.3
1	C	175	TYR	2.3
1	B	58	LYS	2.3
1	B	22	ILE	2.3
1	B	3	GLY	2.3
1	B	120	ILE	2.3
1	B	137	THR	2.2
1	B	145	ASN	2.2
1	B	127	THR	2.2
1	B	97	THR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	138	ARG	2.1
1	C	69	ARG	2.1
1	B	144	GLY	2.1
1	B	19	LEU	2.1
1	B	136	LYS	2.1
1	B	38	PRO	2.1
1	B	39	PRO	2.1
1	B	15	LEU	2.1
1	C	176	LEU	2.1
1	B	20	PRO	2.1
1	B	68	ARG	2.1
1	C	190	ALA	2.1
1	C	182	LEU	2.1
1	B	171	ASN	2.1
1	B	95	VAL	2.1
1	C	144	GLY	2.0
1	A	114	GLU	2.0
1	C	177	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MG	C	301	1/1	0.88	0.20	70,70,70,70	0
2	IPA	A	401	4/4	0.89	0.14	34,36,47,50	0
3	MG	B	301	1/1	0.90	0.14	72,72,72,72	0
3	MG	A	402	1/1	0.90	0.08	53,53,53,53	0

6.5 Other polymers [i](#)

There are no such residues in this entry.