



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 04:38 PM UTC

PDB ID : 4ZOD / pdb_00004zod
Title : Crystal Structure of beta-glucosidase from *Listeria innocua* in complex with glucose
Authors : Nakajima, M.; Yoshida, R.; Miyanaga, A.; Abe, K.; Takahashi, Y.; Sugimoto, N.; Toyozumi, H.; Nakai, H.; Kitaoka, M.; Taguchi, H.
Deposited on : 2015-05-06
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	731	 85% 12% ..
1	B	731	 87% 11% ..

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 11754 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lin1840 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	720	Total	C	N	O	S	0	2	0
			5578	3533	923	1097	25			
1	B	721	Total	C	N	O	S	0	1	0
			5578	3531	923	1099	25			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	GLU	LYS	engineered mutation	UNP Q92AS9
A	724	LEU	-	expression tag	UNP Q92AS9
A	725	GLU	-	expression tag	UNP Q92AS9
A	726	HIS	-	expression tag	UNP Q92AS9
A	727	HIS	-	expression tag	UNP Q92AS9
A	728	HIS	-	expression tag	UNP Q92AS9
A	729	HIS	-	expression tag	UNP Q92AS9
A	730	HIS	-	expression tag	UNP Q92AS9
A	731	HIS	-	expression tag	UNP Q92AS9
B	2	GLU	LYS	engineered mutation	UNP Q92AS9
B	724	LEU	-	expression tag	UNP Q92AS9
B	725	GLU	-	expression tag	UNP Q92AS9
B	726	HIS	-	expression tag	UNP Q92AS9
B	727	HIS	-	expression tag	UNP Q92AS9
B	728	HIS	-	expression tag	UNP Q92AS9
B	729	HIS	-	expression tag	UNP Q92AS9
B	730	HIS	-	expression tag	UNP Q92AS9
B	731	HIS	-	expression tag	UNP Q92AS9

- Molecule 2 is beta-D-glucopyranose (CCD ID: BGC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	1
			12	6	6		
4	A	1	Total	C	O	0	1
			12	6	6		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	1
			12	6	6		
4	B	1	Total	C	O	0	1
			12	6	6		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		
5	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	242	Total 242	O 242	0	0
7	B	233	Total 233	O 233	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	89.79Å 95.41Å 213.23Å 90.00° 97.27° 90.00°	Depositor
Resolution (Å)	37.10 – 2.10 37.10 – 2.10	Depositor EDS
% Data completeness (in resolution range)	97.5 (37.10-2.10) 97.5 (37.10-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.10 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.167 , 0.233 (Not available) , 0.203	Depositor DCC
R_{free} test set	5084 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	25.0	Xtriage
Anisotropy	0.062	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 47.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11754	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, GOL, SO4, MG, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.14	5/5679 (0.1%)	1.13	18/7696 (0.2%)
1	B	1.10	7/5679 (0.1%)	1.12	14/7695 (0.2%)
All	All	1.12	12/11358 (0.1%)	1.13	32/15391 (0.2%)

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	212	THR	C-O	-6.52	1.16	1.24
1	A	594	PRO	C-O	-5.88	1.16	1.23
1	A	501	VAL	C-O	-5.84	1.17	1.24
1	B	84	ILE	N-CA	5.74	1.50	1.46
1	A	274	VAL	CA-CB	5.65	1.61	1.54

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	482	LEU	CA-C-N	-7.52	112.23	119.90
1	B	482	LEU	C-N-CA	-7.52	112.23	119.90
1	A	672	GLU	N-CA-C	7.20	120.14	110.35
1	B	501	VAL	CB-CA-C	-6.94	100.49	110.83
1	A	575	THR	CA-C-N	-6.87	112.62	119.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5578	0	5534	51	0
1	B	5578	0	5527	44	0
2	A	12	0	12	1	0
2	B	24	0	24	2	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	42	0	56	5	0
4	B	24	0	32	2	0
5	A	7	0	10	1	0
5	B	7	0	10	1	0
6	A	5	0	0	0	0
7	A	242	0	0	2	0
7	B	233	0	0	3	0
All	All	11754	0	11205	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 94 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:210[A]:MET:HE1	1:B:219:TYR:CD1	2.06	0.91
1:A:210[A]:MET:HE1	1:A:219:TYR:CD1	2.08	0.88
1:B:299:ASP:OD1	1:B:331:ARG:NH1	2.08	0.86
1:A:244[A]:ILE:HD11	1:B:653:ILE:HD13	1.66	0.75
1:A:299:ASP:OD1	1:A:331:ARG:NH1	2.20	0.74

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	718/731 (98%)	696 (97%)	22 (3%)	0	100	100
1	B	718/731 (98%)	694 (97%)	24 (3%)	0	100	100
All	All	1436/1462 (98%)	1390 (97%)	46 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	603/612 (98%)	586 (97%)	17 (3%)	38	43
1	B	603/612 (98%)	585 (97%)	18 (3%)	36	41
All	All	1206/1224 (98%)	1171 (97%)	35 (3%)	37	42

5 of 35 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	455	GLN
1	B	577	GLU
1	B	664	GLU
1	A	606	GLU
1	A	579	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	3	GLN
1	B	182	ASN
1	B	185	GLN
1	B	468	ASN
1	B	625	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 19 ligands modelled in this entry, 2 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BGC	B	803	-	12,12,12	0.99	0	17,17,17	2.66	8 (47%)
4	GOL	A	808[B]	-	5,5,5	0.54	0	5,5,5	0.88	0
2	BGC	B	802	-	12,12,12	0.69	0	17,17,17	1.37	3 (17%)
4	GOL	B	805[B]	-	5,5,5	0.49	0	5,5,5	0.67	0
4	GOL	A	804[B]	-	5,5,5	0.39	0	5,5,5	0.34	0
4	GOL	A	808[A]	-	5,5,5	0.46	0	5,5,5	1.39	1 (20%)
5	PEG	A	806	-	6,6,6	0.48	0	5,5,5	1.28	1 (20%)
2	BGC	A	801	-	12,12,12	0.94	0	17,17,17	1.54	2 (11%)
4	GOL	A	805	-	5,5,5	0.45	0	5,5,5	0.85	0
4	GOL	A	803[B]	-	5,5,5	0.77	0	5,5,5	0.96	0
4	GOL	B	805[A]	-	5,5,5	0.51	0	5,5,5	0.53	0
4	GOL	B	806	-	5,5,5	0.23	0	5,5,5	0.71	0
4	GOL	A	803[A]	-	5,5,5	0.74	0	5,5,5	1.45	1 (20%)
6	SO4	A	807	-	4,4,4	0.34	0	6,6,6	0.26	0
4	GOL	A	804[A]	-	5,5,5	0.70	0	5,5,5	0.75	0
5	PEG	B	801	-	6,6,6	0.66	0	5,5,5	0.24	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	B	807	-	5,5,5	0.51	0	5,5,5	1.30	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	B	803	-	-	2/2/22/22	0/1/1/1
4	GOL	A	808[B]	-	-	4/4/4/4	-
2	BGC	B	802	-	-	0/2/22/22	0/1/1/1
4	GOL	B	805[B]	-	-	2/4/4/4	-
4	GOL	A	804[B]	-	-	0/4/4/4	-
4	GOL	A	808[A]	-	-	4/4/4/4	-
5	PEG	A	806	-	-	2/4/4/4	-
2	BGC	A	801	-	-	0/2/22/22	0/1/1/1
4	GOL	A	805	-	-	2/4/4/4	-
4	GOL	A	803[B]	-	-	0/4/4/4	-
4	GOL	B	805[A]	-	-	2/4/4/4	-
4	GOL	B	806	-	-	0/4/4/4	-
4	GOL	A	803[A]	-	-	3/4/4/4	-
4	GOL	A	804[A]	-	-	2/4/4/4	-
5	PEG	B	801	-	-	2/4/4/4	-
4	GOL	B	807	-	-	2/4/4/4	-

There are no bond length outliers.

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	803	BGC	C6-C5-C4	5.32	126.08	113.02
2	A	801	BGC	O5-C5-C4	-4.02	102.45	109.70
2	B	803	BGC	O1-C1-C2	3.91	120.33	108.98
2	B	803	BGC	C3-C4-C5	-3.79	103.36	110.23
2	B	803	BGC	O2-C2-C3	-3.43	102.28	110.38

There are no chirality outliers.

5 of 27 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	804[A]	GOL	O1-C1-C2-C3
4	A	808[A]	GOL	C1-C2-C3-O3
4	A	808[B]	GOL	C1-C2-C3-O3
4	B	805[B]	GOL	C1-C2-C3-O3
2	B	803	BGC	C4-C5-C6-O6

There are no ring outliers.

9 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	802	BGC	2	0
4	B	805[B]	GOL	1	0
5	A	806	PEG	1	0
2	A	801	BGC	1	0
4	A	805	GOL	1	0
4	A	803[B]	GOL	1	0
4	A	804[A]	GOL	3	0
5	B	801	PEG	1	0
4	B	807	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	720/731 (98%)	-0.41	1 (0%) 92 92	8, 26, 50, 73	2 (0%)
1	B	721/731 (98%)	-0.38	3 (0%) 88 90	10, 27, 50, 70	1 (0%)
All	All	1441/1462 (98%)	-0.39	4 (0%) 90 91	8, 27, 50, 73	3 (0%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	67	LEU	2.8
1	A	579	LYS	2.6
1	B	579	LYS	2.3
1	B	39	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	A	803[A]	6/6	0.85	0.13	18,23,24,25	6

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	A	803[B]	6/6	0.85	0.13	19,20,21,22	6
4	GOL	A	808[A]	6/6	0.88	0.11	9,18,23,25	6
4	GOL	A	808[B]	6/6	0.88	0.11	22,23,24,26	6
4	GOL	A	804[A]	6/6	0.89	0.10	10,14,14,17	6
4	GOL	A	804[B]	6/6	0.89	0.10	18,26,28,36	6
5	PEG	B	801	7/7	0.89	0.13	39,44,50,53	0
4	GOL	B	806	6/6	0.93	0.10	40,47,50,50	0
5	PEG	A	806	7/7	0.93	0.09	32,36,39,43	0
4	GOL	A	805	6/6	0.93	0.10	39,44,47,50	0
2	BGC	B	803	12/12	0.94	0.07	33,41,55,57	0
4	GOL	B	807	6/6	0.94	0.08	32,39,44,48	0
4	GOL	B	805[A]	6/6	0.96	0.07	22,23,23,23	6
4	GOL	B	805[B]	6/6	0.96	0.07	17,22,24,26	6
6	SO4	A	807	5/5	0.96	0.12	16,17,18,18	5
2	BGC	B	802	12/12	0.97	0.04	20,24,26,28	0
2	BGC	A	801	12/12	0.97	0.05	15,19,20,22	0
3	MG	A	802	1/1	0.99	0.01	24,24,24,24	0
3	MG	B	804	1/1	1.00	0.01	24,24,24,24	0

6.5 Other polymers

There are no such residues in this entry.