



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:01 PM UTC

PDB ID : 4ZOP / pdb_00004zop
Title : Co-crystal Structure of Lipid Kinase PI3K alpha with a selective phosphatidylinositol-3 kinase alpha inhibitor
Authors : Knapp, M.S.; Elling, R.A.
Deposited on : 2015-05-06
Resolution : 2.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4951 (2.64-2.60)
Clashscore	190562	5303 (2.64-2.60)
Ramachandran outliers	187476	5217 (2.64-2.60)
Sidechain outliers	187428	5217 (2.64-2.60)
RSRZ outliers	180081	4950 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1074	2% 84% 8% 7%
2	B	290	4% 76% 5% 17%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10227 atoms, of which 26 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1000	Total	C	N	O	S	0	1	0
			8080	5169	1378	1468	65			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	232	LYS	MET	engineered mutation	UNP P42336
A	233	LYS	LEU	engineered mutation	UNP P42336
A	1069	HIS	-	expression tag	UNP P42336
A	1070	HIS	-	expression tag	UNP P42336
A	1071	HIS	-	expression tag	UNP P42336
A	1072	HIS	-	expression tag	UNP P42336
A	1073	HIS	-	expression tag	UNP P42336
A	1074	HIS	-	expression tag	UNP P42336


- Molecule 2 is a protein called Phosphatidylinositol 3-kinase regulatory subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	240	Total	C	N	O	S	0	0	0
			1965	1231	349	381	4			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	306	TYR	THR	conflict	UNP P27986

- Molecule 3 is (2S,3R)-N 1 -(8-tert-butyl-4,5-dihydro[1,3]thiazolo[4,5-h]quinazolin-2-yl)-3-methylpyrrolidine-1,2-dicarboxamide (CCD ID: 4Q2) (formula: C₂₀H₂₆N₆O₂S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	S	0	0
			55	20	26	6	2	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	111	Total	O	0	0
			111	111		
4	B	16	Total	O	0	0
			16	16		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform



- Molecule 2: Phosphatidylinositol 3-kinase regulatory subunit alpha



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	104.81Å 106.56Å 134.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.49 – 2.62 47.49 – 2.62	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.49-2.62) 99.9 (47.49-2.62)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 2.61Å)	Xtriage
Refinement program	BUSTER-TNT	Depositor
R, R_{free}	0.193 , 0.258 0.195 , 0.260	Depositor DCC
R_{free} test set	2200 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å ²)	59.5	Xtriage
Anisotropy	0.073	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 47.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.016 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10227	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 4Q2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.89	3/8264 (0.0%)	1.33	40/11188 (0.4%)
2	B	0.90	5/1994 (0.3%)	1.51	10/2678 (0.4%)
All	All	0.89	8/10258 (0.1%)	1.36	50/13866 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1004	MET	SD-CE	-8.26	1.58	1.79
2	B	395	PRO	N-CD	7.49	1.58	1.47
2	B	398	PHE	CA-C	7.12	1.62	1.52
2	B	397	THR	C-N	5.36	1.41	1.34
1	A	633	ILE	CG1-CD1	5.16	1.71	1.51
1	A	216	VAL	CA-C	5.14	1.58	1.52
2	B	397	THR	CA-C	5.09	1.59	1.52
2	B	563	MET	SD-CE	-5.03	1.67	1.79

All (50) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	396	LEU	CA-C-N	11.44	143.39	121.54
2	B	396	LEU	C-N-CA	11.44	143.39	121.54
2	B	394	ASP	CA-CB-CG	9.24	121.84	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	937	PHE	CA-CB-CG	7.97	121.78	113.80
2	B	440	ASP	CA-CB-CG	7.58	120.18	112.60
1	A	603	ASP	CA-CB-CG	6.96	119.56	112.60
2	B	397	THR	N-CA-CB	6.59	121.62	110.49
2	B	433	GLN	N-CA-C	-6.27	103.97	111.69
1	A	377	PRO	CA-C-N	6.24	128.93	120.38
1	A	377	PRO	C-N-CA	6.24	128.93	120.38
1	A	873	ASN	CA-CB-CG	6.06	118.66	112.60
1	A	415	ALA	N-CA-C	-6.05	105.21	112.59
1	A	133	ASP	CA-CB-CG	5.81	118.41	112.60
1	A	478	ASP	CA-CB-CG	5.80	118.40	112.60
1	A	120	ALA	CA-C-N	5.71	130.37	121.35
1	A	120	ALA	C-N-CA	5.71	130.37	121.35
1	A	900	SER	CA-C-N	5.68	128.16	120.38
1	A	900	SER	C-N-CA	5.68	128.16	120.38
1	A	64	ASP	CA-CB-CG	5.64	118.24	112.60
1	A	310	ARG	CA-C-N	5.62	130.61	121.62
1	A	310	ARG	C-N-CA	5.62	130.61	121.62
1	A	843	ASP	CA-CB-CG	5.60	118.20	112.60
1	A	883	ASP	CA-CB-CG	5.58	118.18	112.60
1	A	897	PHE	CA-C-N	5.40	127.46	120.44
1	A	897	PHE	C-N-CA	5.40	127.46	120.44
1	A	369	ASP	CA-CB-CG	5.39	117.99	112.60
1	A	684	PHE	CA-C-N	5.36	125.79	119.94
1	A	684	PHE	C-N-CA	5.36	125.79	119.94
1	A	482	SER	CA-C-N	5.33	129.04	122.37
1	A	482	SER	C-N-CA	5.33	129.04	122.37
2	B	396	LEU	CB-CA-C	-5.33	99.81	110.42
1	A	958	GLN	CA-C-N	5.33	127.68	120.38
1	A	958	GLN	C-N-CA	5.33	127.68	120.38
1	A	1037	GLU	CA-C-N	5.31	127.34	120.44
1	A	1037	GLU	C-N-CA	5.31	127.34	120.44
2	B	395	PRO	CA-C-N	5.30	131.66	121.54
2	B	395	PRO	C-N-CA	5.30	131.66	121.54
1	A	976	GLU	CB-CG-CD	5.26	121.55	112.60
1	A	145	ASN	CA-C-N	5.25	127.66	120.46
1	A	145	ASN	C-N-CA	5.25	127.66	120.46
1	A	488	ASP	CA-CB-CG	5.25	117.85	112.60
1	A	548	LYS	CA-C-N	5.24	127.56	120.38
1	A	548	LYS	C-N-CA	5.24	127.56	120.38
1	A	983	MET	CA-C-N	5.11	127.09	120.44
1	A	983	MET	C-N-CA	5.11	127.09	120.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	352	ASP	CA-CB-CG	5.04	117.64	112.60
1	A	214	ASP	CA-CB-CG	5.04	117.64	112.60
1	A	372	ASN	CA-CB-CG	5.04	117.64	112.60
1	A	258	ASP	CA-CB-CG	5.03	117.63	112.60
1	A	998	PHE	CA-CB-CG	-5.02	108.78	113.80

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	398	PHE	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8080	0	7981	36	0
2	B	1965	0	1870	20	0
3	A	29	26	0	1	0
4	A	111	0	0	0	0
4	B	16	0	0	0	0
All	All	10201	26	9851	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:396:LEU:HD22	2:B:398:PHE:CE2	1.52	1.43
2:B:396:LEU:CD2	2:B:398:PHE:HE2	1.70	1.04
2:B:396:LEU:CD2	2:B:398:PHE:CE2	2.48	0.90
1:A:897:PHE:O	1:A:901:CYS:HB2	1.83	0.78
2:B:396:LEU:HD12	2:B:397:THR:N	2.01	0.76
1:A:121:ILE:HG12	1:A:688:LEU:HB3	1.71	0.71

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:494:PHE:HB3	2:B:535:ILE:HG12	1.73	0.70
2:B:396:LEU:HD12	2:B:396:LEU:C	2.18	0.68
2:B:394:ASP:HB2	2:B:395:PRO:HA	1.77	0.65
2:B:396:LEU:CD1	2:B:397:THR:N	2.63	0.62
1:A:712:LEU:HD21	1:A:748:LEU:HD21	1.83	0.61
1:A:697:MET:SD	1:A:700:LYS:HD2	2.42	0.59
1:A:599:MET:HE1	1:A:631:TYR:HB3	1.86	0.57
2:B:334:TYR:O	2:B:430:LYS:HE2	2.04	0.56
1:A:640:LYS:HG2	1:A:680:VAL:HG11	1.87	0.55
2:B:577:ARG:HH11	2:B:577:ARG:HG2	1.72	0.55
1:A:25:LEU:HD13	2:B:497:GLN:HG3	1.90	0.54
1:A:937:PHE:HD1	1:A:938:LEU:HG	1.74	0.53
1:A:42:LEU:HD21	1:A:92:LEU:HD11	1.92	0.52
1:A:572:VAL:CG2	1:A:583:MET:HE3	2.39	0.52
2:B:396:LEU:HD12	2:B:397:THR:CA	2.40	0.51
1:A:279:LEU:HB2	1:A:281:ARG:HG2	1.92	0.51
1:A:572:VAL:HG21	1:A:583:MET:HG2	1.92	0.51
2:B:396:LEU:HD22	2:B:398:PHE:CD2	2.32	0.51
2:B:396:LEU:HB2	2:B:398:PHE:HD2	1.76	0.50
2:B:396:LEU:HD12	2:B:397:THR:HA	1.92	0.50
1:A:328:TRP:HA	1:A:394:PRO:HB3	1.94	0.50
1:A:25:LEU:HD13	2:B:497:GLN:CG	2.41	0.50
1:A:279:LEU:CB	1:A:281:ARG:HG2	2.41	0.49
2:B:396:LEU:HD22	2:B:398:PHE:HE2	0.83	0.49
2:B:396:LEU:HB2	2:B:398:PHE:CD2	2.48	0.48
1:A:363:GLY:N	1:A:607:PRO:HG3	2.28	0.48
1:A:311:ILE:H	1:A:311:ILE:HD12	1.80	0.47
1:A:267:LEU:HG	1:A:273:ILE:HG13	1.95	0.47
2:B:396:LEU:C	2:B:396:LEU:CD1	2.84	0.47
1:A:336:ILE:HD13	1:A:402:LEU:HD22	1.95	0.47
3:A:1101:4Q2:S5	3:A:1101:4Q2:O21	2.73	0.47
1:A:197:ILE:HD11	1:A:789:MET:HE2	1.97	0.45
1:A:83:PHE:CE1	1:A:108:ARG:HG3	2.52	0.45
1:A:937:PHE:CD1	1:A:938:LEU:HG	2.52	0.43
1:A:344:VAL:HG23	1:A:346:VAL:HG22	2.00	0.43
1:A:705:GLN:O	1:A:709:MET:HG2	2.18	0.43
1:A:69:ILE:HG23	1:A:104:PRO:HG3	2.01	0.42
1:A:408:SER:HB3	1:A:422:LEU:HD21	2.02	0.42
1:A:719:LEU:HA	1:A:723:LYS:HB2	2.02	0.42
1:A:808:ARG:HB3	1:A:1010:MET:HE2	2.01	0.42
2:B:396:LEU:CD1	2:B:397:THR:HA	2.50	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:712:LEU:CD2	1:A:748:LEU:HD21	2.50	0.41
1:A:552:TRP:HZ3	1:A:583:MET:HE2	1.86	0.41
1:A:587:VAL:HA	1:A:590:TRP:HB2	2.02	0.41
1:A:647:ASN:O	1:A:651:ARG:HG3	2.21	0.41
1:A:777:ARG:N	1:A:778:PRO:HD3	2.36	0.40
1:A:599:MET:HE1	1:A:631:TYR:CB	2.49	0.40
1:A:638:VAL:HG23	1:A:649:LEU:HD21	2.03	0.40
1:A:159:PRO:HG2	1:A:294:TYR:CD2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	991/1074 (92%)	945 (95%)	44 (4%)	2 (0%)	43	64
2	B	232/290 (80%)	220 (95%)	7 (3%)	5 (2%)	5	9
All	All	1223/1364 (90%)	1165 (95%)	51 (4%)	7 (1%)	21	39

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	396	LEU
2	B	397	THR
2	B	399	SER
1	A	131	VAL
2	B	425	LEU
2	B	395	PRO
1	A	935	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	890/980 (91%)	881 (99%)	9 (1%)	68	85
2	B	203/269 (76%)	200 (98%)	3 (2%)	57	79
All	All	1093/1249 (88%)	1081 (99%)	12 (1%)	65	83

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	115	ARG
1	A	121	ILE
1	A	251	ILE
1	A	628	LEU
1	A	720	LYS
1	A	748	LEU
1	A	834	LEU
1	A	841	ILE
1	A	913	ILE
2	B	328	LEU
2	B	442	ILE
2	B	579	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	212	ASN
1	A	219	GLN
1	A	347	ASN
1	A	380	ASN
1	A	714	ASN
1	A	809	GLN
2	B	441	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	4Q2	A	1101	-	30,32,32	0.30	0	41,49,49	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	4Q2	A	1101	-	-	3/18/40/40	0/4/4/4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1101	4Q2	N16-C17-C27-O28
3	A	1101	4Q2	N16-C17-C27-N29
3	A	1101	4Q2	C18-C17-C27-O28

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1101	4Q2	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1000/1074 (93%)	-0.07	18 (1%) 67 63	29, 63, 99, 150	1 (0%)
2	B	240/290 (82%)	0.25	11 (4%) 37 32	48, 77, 127, 143	0
All	All	1240/1364 (90%)	-0.01	29 (2%) 61 56	29, 66, 108, 150	1 (0%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	864	GLY	5.0
1	A	1061	THR	4.2
1	A	246	TYR	4.0
2	B	361	SER	3.7
1	A	496	ALA	3.7
2	B	398	PHE	3.4
2	B	437	VAL	3.3
1	A	3	PRO	3.2
1	A	311	ILE	3.1
2	B	589	VAL	2.9
2	B	391	GLY	2.8
1	A	1055	MET	2.8
1	A	970	GLU	2.6
2	B	588	GLY	2.6
2	B	328	LEU	2.6
1	A	313	THR	2.6
1	A	312	SER	2.5
1	A	314	ALA	2.5
1	A	247	GLN	2.4
1	A	973	LYS	2.4
2	B	431	TYR	2.3
2	B	435	GLN	2.2
2	B	326	MET	2.2
2	B	436	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	478	ASP	2.1
1	A	1057	TRP	2.1
1	A	378	CYS	2.1
1	A	558	CYS	2.1
1	A	969	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	4Q2	A	1101	29/29	0.96	0.06	35,41,54,56	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 4Q2 A 1101:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.