



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:31 AM UTC

PDB ID : 5ZOG / pdb_00005zog
Title : Crystal Structure of R192F hFen1 in complex with DNA
Authors : Han, W.; Hua, Y.; Zhao, Y.
Deposited on : 2018-04-13
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	344	17% 83% 8% 8%
2	B	7	100%
3	C	19	47% 68% 32%
4	D	13	85% 23% 77%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3359 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Flap endonuclease 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	316	Total	C	N	O	S	0	0	0
			2483	1574	430	464	15			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	192	PHE	ARG	engineered mutation	UNP P39748
A	334	LYS	-	expression tag	UNP P39748
A	335	LEU	-	expression tag	UNP P39748
A	336	ALA	-	expression tag	UNP P39748
A	337	ALA	-	expression tag	UNP P39748
A	338	ALA	-	expression tag	UNP P39748
A	339	LEU	-	expression tag	UNP P39748
A	340	GLU	-	expression tag	UNP P39748
A	341	HIS	-	expression tag	UNP P39748
A	342	HIS	-	expression tag	UNP P39748
A	343	HIS	-	expression tag	UNP P39748
A	344	HIS	-	expression tag	UNP P39748

- Molecule 2 is a DNA chain called DNA (5'-D(*CP*CP*CP*GP*TP*CP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	7	Total	C	N	O	P	0	0	0
			133	65	22	40	6			

- Molecule 3 is a DNA chain called DNA (5'-D(P*TP*CP*CP*TP*CP*TP*GP*CP*CP*TP*CP*AP*AP*GP*AP*CP*GP*GP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	19	Total	C	N	O	P	0	0	0
			386	183	69	115	19			

- Molecule 4 is a DNA chain called DNA (5'-D(*TP*GP*AP*GP*GP*CP*AP*GP*AP*GP*GP*AP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	13	Total 272	C 129	N 57	O 74	P 12	0	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	72	Total 72	O 72	0	0
5	B	5	Total 5	O 5	0	0
5	C	8	Total 8	O 8	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Flap endonuclease 1




- Molecule 2: DNA (5'-D(*CP*CP*CP*GP*TP*CP*C)-3')




There are no outlier residues recorded for this chain.

- Molecule 3: DNA (5'-D(P*TP*CP*CP*TP*CP*TP*GP*CP*CP*TP*CP*AP*AP*GP*AP*C P*GP*GP*G)-3')



- Molecule 4: DNA (5'-D(*TP*GP*AP*GP*GP*CP*AP*GP*AP*GP*GP*AP*T)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	88.33Å 244.80Å 70.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.56 – 2.30 29.56 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.8 (29.56-2.30) 92.9 (29.56-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	-0.04 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.209 , 0.231 0.210 , 0.232	Depositor DCC
R_{free} test set	1742 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	37.0	Xtriage
Anisotropy	0.496	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 38.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3359	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/2531	0.50	0/3410
2	B	0.46	0/147	0.60	0/224
3	C	0.35	0/431	0.55	0/662
4	D	0.21	0/307	0.43	0/474
All	All	0.34	0/3416	0.51	0/4770

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2483	0	2482	20	0
2	B	133	0	77	0	0
3	C	386	0	214	8	0
4	D	272	0	147	12	0
5	A	72	0	0	1	0
5	B	5	0	0	0	0
5	C	8	0	0	0	0
All	All	3359	0	2920	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:160:GLU:OE2	1:A:182:CYS:SG	2.23	0.96
3:C:11:DA:H2''	3:C:12:DA:H5''	1.65	0.77
4:D:11:DG:H2'	4:D:12:DG:C8	2.27	0.69
4:D:10:DA:H2''	4:D:11:DG:H5''	1.74	0.69
4:D:6:DG:H2''	4:D:7:DC:H5''	1.77	0.66
4:D:11:DG:H5''	4:D:11:DG:H8	1.62	0.65
1:A:221:GLU:HG2	1:A:284:LEU:HD11	1.83	0.60
4:D:7:DC:H2''	4:D:8:DA:C8	2.38	0.59
1:A:253:HIS:NE2	1:A:262:ARG:HD3	2.20	0.56
1:A:262:ARG:HB3	1:A:262:ARG:NH1	2.23	0.53
3:C:6:DG:C8	3:C:6:DG:H5'	2.44	0.53
3:C:8:DC:H2''	3:C:9:DT:H5'	1.90	0.53
3:C:1:DC:H6	3:C:1:DC:H5'	1.76	0.51
1:A:260:VAL:HG23	1:A:270:VAL:HG11	1.93	0.49
3:C:1:DC:H5'	3:C:1:DC:C6	2.47	0.49
3:C:12:DA:H5''	3:C:12:DA:C8	2.48	0.49
1:A:27:PHE:CD2	1:A:78:GLY:HA3	2.48	0.49
4:D:3:DG:H4'	4:D:4:DA:OP1	2.14	0.47
1:A:7:ALA:HB2	1:A:180:MET:HE3	1.97	0.47
4:D:10:DA:H2''	4:D:11:DG:C8	2.51	0.45
1:A:92:LEU:HD11	1:A:278:GLU:HB3	2.00	0.43
4:D:6:DG:C2'	4:D:7:DC:H5''	2.46	0.43
1:A:167:VAL:CG1	1:A:188:PRO:HD2	2.49	0.43
1:A:30:LYS:HB2	1:A:30:LYS:HE2	1.76	0.42
4:D:12:DG:H2''	4:D:13:DA:H8	1.82	0.42
1:A:167:VAL:HG12	1:A:188:PRO:HD2	2.01	0.42
1:A:263:LEU:HD22	1:A:268:TYR:HD2	1.85	0.42
4:D:8:DA:H8	4:D:8:DA:O5'	2.03	0.42
4:D:12:DG:H8	4:D:12:DG:OP2	2.03	0.42
1:A:211:ARG:HD3	1:A:211:ARG:HA	1.90	0.41
1:A:284:LEU:HD23	1:A:284:LEU:HA	1.90	0.41
1:A:314:LYS:NZ	5:A:401:HOH:O	2.45	0.41
1:A:67:MET:HE3	1:A:67:MET:HB2	1.95	0.41
3:C:6:DG:H5'	3:C:6:DG:H8	1.85	0.41
4:D:9:DG:H2''	4:D:10:DA:C8	2.55	0.41
1:A:257:GLU:O	1:A:260:VAL:HG12	2.20	0.41
1:A:88:LYS:HA	1:A:88:LYS:HD3	1.82	0.41
1:A:6:LEU:HD23	1:A:180:MET:HB3	2.03	0.41
3:C:12:DA:H5''	3:C:12:DA:H8	1.84	0.41
1:A:92:LEU:HD12	1:A:282:LEU:HD12	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	312/344 (91%)	305 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	268/294 (91%)	262 (98%)	6 (2%)	45	65

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	ILE
1	A	57	GLU
1	A	67	MET
1	A	184	THR
1	A	206	GLU
1	A	270	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	276	HIS
1	A	333	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	316/344 (91%)	0.84	58 (18%) 3 4	27, 48, 107, 124	0
2	B	7/7 (100%)	-0.08	0 100 100	33, 37, 49, 53	0
3	C	19/19 (100%)	1.66	9 (47%) 0 0	34, 96, 124, 130	0
4	D	13/13 (100%)	2.69	11 (84%) 0 0	91, 113, 133, 133	0
All	All	355/383 (92%)	0.93	78 (21%) 2 2	27, 49, 114, 133	0

All (78) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	265	PRO	5.7
1	A	98	ALA	5.3
1	A	100	ARG	5.1
1	A	269	PRO	5.1
3	C	0	DT	4.7
1	A	261	ARG	4.5
1	A	130	LEU	4.4
1	A	275	LEU	4.3
1	A	99	LYS	4.2
1	A	129	ARG	4.0
1	A	268	TYR	4.0
1	A	264	ASP	3.6
1	A	267	LYS	3.6
1	A	262	ARG	3.5
1	A	344	HIS	3.5
3	C	1	DC	3.5
4	D	11	DG	3.5
1	A	257	GLU	3.5
4	D	8	DA	3.4
1	A	277	LYS	3.4
4	D	4	DA	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	246	ALA	3.3
1	A	271	PRO	3.2
1	A	259	ILE	3.2
1	A	260	VAL	3.2
1	A	263	LEU	3.2
1	A	270	VAL	3.2
4	D	14	DT	3.1
1	A	248	ASP	3.1
1	A	240	GLY	3.0
4	D	7	DC	2.9
1	A	245	ARG	2.9
3	C	12	DA	2.9
1	A	97	LEU	2.9
1	A	91	GLN	2.9
1	A	182	CYS	2.9
1	A	241	ILE	2.9
3	C	10	DC	2.9
1	A	88	LYS	2.8
4	D	9	DG	2.8
4	D	12	DG	2.8
1	A	295	GLU	2.8
1	A	128	LYS	2.7
1	A	249	LEU	2.7
3	C	2	DC	2.6
1	A	250	ILE	2.6
1	A	266	ASN	2.6
1	A	276	HIS	2.6
1	A	225	ASP	2.5
4	D	6	DG	2.5
3	C	7	DC	2.5
1	A	252	LYS	2.5
3	C	8	DC	2.5
1	A	22	ASP	2.5
1	A	253	HIS	2.5
1	A	239	ARG	2.4
1	A	341	HIS	2.4
1	A	244	LYS	2.3
4	D	13	DA	2.3
1	A	218	LEU	2.3
3	C	11	DA	2.3
1	A	14	ALA	2.3
1	A	272	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	6	DG	2.2
4	D	5	DG	2.2
1	A	131	VAL	2.2
1	A	41	GLN	2.2
4	D	2	DT	2.2
1	A	304	GLU	2.2
1	A	217	GLY	2.1
1	A	238	ILE	2.1
1	A	282	LEU	2.1
1	A	19	ARG	2.1
1	A	256	ILE	2.1
1	A	258	GLU	2.0
1	A	284	LEU	2.0
1	A	228	ILE	2.0
1	A	273	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.