



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:36 PM UTC

PDB ID : 4ZP2 / pdb_00004zp2
Title : Crystal structure of E. coli multidrug transporter MdfA in complex with n-dodecyl-N,N-dimethylamine-N-oxide
Authors : Zhang, X.C.; Heng, J.; Zhao, Y.; Wang, X.
Deposited on : 2015-05-07
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	392	10% 84% 15% .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3023 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Multidrug transporter MdfA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	392	2973	1974	479	500	20	0	1	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	131	ARG	GLN	engineered mutation	UNP P0AEY8

- Molecule 2 is LAURYL DIMETHYLAMINE-N-OXIDE (CCD ID: LDA) (formula: $C_{14}H_{31}NO$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	16	14	1	1	0	0
2	A	1	16	14	1	1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	18	Total	O	0	0
			18	18		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Multidrug transporter Mdfa



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	94.09Å 64.08Å 101.25Å 90.00° 104.19° 90.00°	Depositor
Resolution (Å)	38.44 – 2.20 38.44 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.1 (38.44-2.20) 99.4 (38.44-2.20)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.36 (at 2.20Å)	Xtriage
Refinement program	PHENIX phenix.refine: 1.8_1069	Depositor
R, R_{free}	0.206 , 0.232 0.216 , 0.218	Depositor DCC
R_{free} test set	1529 reflections (4.70%)	wwPDB-VP
Wilson B-factor (Å ²)	40.2	Xtriage
Anisotropy	0.226	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 42.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3023	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: LDA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/3042	0.82	3/4139 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	331	ASN	N-CA-C	6.21	117.85	111.14
1	A	10	ARG	N-CA-C	5.14	121.33	113.51
1	A	330	ALA	N-CA-C	5.12	118.69	112.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2973	0	3115	30	1
2	A	32	0	62	1	0
3	A	18	0	0	1	1
All	All	3023	0	3177	31	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:220:ARG:NH1	1:A:398:PHE:O	2.25	0.68
1:A:55:PRO:HG2	1:A:369:LYS:HG3	1.77	0.67
1:A:345:SER:OG	3:A:602:HOH:O	2.16	0.62
1:A:82:ARG:NH2	1:A:189:GLN:O	2.35	0.59
1:A:99:ILE:HA	1:A:111:LEU:HD13	1.86	0.57
1:A:127:TYR:HE1	1:A:143:THR:HG22	1.69	0.57
1:A:139:CYS:O	1:A:143:THR:HG23	2.05	0.57
1:A:23:VAL:HG13	1:A:149:VAL:HG21	1.88	0.55
1:A:21:CYS:SG	1:A:187:GLY:HA3	2.47	0.54
1:A:375:GLY:HA3	1:A:379:LEU:HD22	1.91	0.52
1:A:231:VAL:HB	1:A:391:TRP:CD1	2.47	0.49
1:A:142:ILE:HG22	1:A:146:MET:HE3	1.95	0.48
1:A:283:THR:HB	1:A:285:ARG:NH1	2.28	0.48
1:A:282:ARG:HG3	1:A:286:SER:OG	2.13	0.47
1:A:127:TYR:CE1	1:A:143:THR:HG22	2.48	0.47
1:A:284:VAL:HG22	1:A:285:ARG:HH11	1.80	0.46
1:A:282:ARG:HG2	1:A:290:MET:HE1	1.97	0.46
1:A:399:LEU:HA	1:A:400:LYS:HA	1.71	0.45
1:A:74:PRO:O	1:A:78:ARG:HG2	2.17	0.44
1:A:71:LEU:HD21	1:A:75:LEU:HD22	1.99	0.44
1:A:344:MET:HE2	1:A:344:MET:HB3	1.91	0.44
1:A:304:ALA:O	1:A:308:ILE:HG12	2.19	0.43
1:A:121:PHE:O	1:A:125:VAL:HG13	2.19	0.43
1:A:16:LEU:C	1:A:19:PRO:HD2	2.43	0.42
1:A:96:CYS:O	1:A:99:ILE:HG22	2.19	0.42
2:A:500:LDA:HM11	2:A:500:LDA:H22	1.61	0.41
1:A:10:ARG:HA	1:A:14:GLN:HB2	2.01	0.41
1:A:199:ILE:HA	1:A:200:GLY:HA2	1.88	0.41
1:A:25:TYR:OH	1:A:118:SER:HB3	2.21	0.41
1:A:44:VAL:HG22	1:A:49:ALA:HB3	2.02	0.41
1:A:395:MET:HA	1:A:395:MET:HE2	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:281:ARG:NH1	3:A:601:HOH:O[3_555]	2.19	0.01

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	391/392 (100%)	383 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	305/307 (99%)	290 (95%)	15 (5%)	22	29

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	LEU
1	A	44	VAL
1	A	46	GLN
1	A	119	LEU
1	A	125	VAL
1	A	151	LEU
1	A	202	LYS
1	A	252	LEU
1	A	259	LEU
1	A	268	LEU
1	A	282	ARG
1	A	285	ARG
1	A	386	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	396	VAL
1	A	399	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	GLN
1	A	381	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	LDA	A	501	-	13,15,15	2.38	2 (15%)	14,17,17	0.60	0
2	LDA	A	500	-	13,15,15	2.32	2 (15%)	14,17,17	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LDA	A	501	-	-	9/13/13/13	-
2	LDA	A	500	-	-	7/13/13/13	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	LDA	O1-N1	-6.74	1.25	1.42
2	A	500	LDA	O1-N1	-6.65	1.25	1.42
2	A	501	LDA	C1-N1	-5.24	1.46	1.51
2	A	500	LDA	C1-N1	-4.97	1.46	1.51

There are no bond angle outliers.

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	LDA	C2-C1-N1-CM2
2	A	501	LDA	C7-C8-C9-C10
2	A	500	LDA	C4-C5-C6-C7
2	A	501	LDA	C5-C6-C7-C8
2	A	500	LDA	C1-C2-C3-C4
2	A	500	LDA	C5-C6-C7-C8
2	A	501	LDA	C1-C2-C3-C4
2	A	501	LDA	C6-C7-C8-C9
2	A	501	LDA	C11-C10-C9-C8
2	A	501	LDA	C9-C10-C11-C12
2	A	501	LDA	C4-C5-C6-C7
2	A	500	LDA	C7-C8-C9-C10
2	A	500	LDA	C11-C10-C9-C8
2	A	500	LDA	C9-C10-C11-C12
2	A	500	LDA	C3-C4-C5-C6
2	A	501	LDA	C2-C1-N1-CM1

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	LDA	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	392/392 (100%)	0.69	40 (10%) 12 9	32, 48, 75, 119	1 (0%)

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	12	GLY	12.6
1	A	13	ARG	8.7
1	A	9	ALA	7.0
1	A	11	LEU	5.7
1	A	10	ARG	5.2
1	A	340	PHE	4.2
1	A	312	ALA	4.0
1	A	399	LEU	3.6
1	A	15	ALA	3.5
1	A	183	ILE	3.5
1	A	102	ALA	3.5
1	A	343	ASP	3.4
1	A	16	LEU	3.2
1	A	346	LYS	3.0
1	A	191	ALA	2.9
1	A	135	GLU	2.9
1	A	186	PHE	2.9
1	A	197	THR	2.8
1	A	14	GLN	2.8
1	A	400	LYS	2.8
1	A	256	GLU	2.7
1	A	200	GLY	2.7
1	A	190	ARG	2.7
1	A	182	ALA	2.6
1	A	206	LYS	2.5
1	A	30	TYR	2.4
1	A	284	VAL	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	313	TYR	2.4
1	A	211	ASP	2.4
1	A	104	ASN	2.3
1	A	308	ILE	2.3
1	A	188	LEU	2.3
1	A	163	ALA	2.2
1	A	341	ALA	2.2
1	A	196	ALA	2.2
1	A	168	LEU	2.2
1	A	78	ARG	2.2
1	A	133	SER	2.1
1	A	132	GLU	2.1
1	A	46	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	LDA	A	501	16/16	0.78	0.26	53,67,87,91	0
2	LDA	A	500	16/16	0.81	0.26	48,62,73,74	0

6.5 Other polymers [i](#)

There are no such residues in this entry.