



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:06 AM UTC

PDB ID : 6ZPJ / pdb_00006zpj
Title : Crystal structure of the unconventional kinetochore protein Leishmania mexicana KKT4 coiled coil domain
Authors : Ludzia, P.; Lowe, E.D.; Marciano, G.; Mohammed, S.; Redfield, C.; Akiyoshi, B.
Deposited on : 2020-07-08
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	126	10% 75% 12% • 13%
1	B	126	2% 52% 9% 39%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 1816 atoms, of which 38 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Leishmania mexicana KKT4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	110	Total	C	N	O	S	0	3	0
			874	530	168	174	2			
1	B	77	Total	C	N	O	S	0	2	0
			614	368	119	126	1			

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	182	GLY	-	expression tag	UNP E9AN40
A	183	SER	-	expression tag	UNP E9AN40
A	218	GLN	ARG	engineered mutation	UNP E9AN40
A	285	GLU	-	cloning artifact	UNP E9AN40
A	286	PHE	-	cloning artifact	UNP E9AN40
A	287	GLU	-	cloning artifact	UNP E9AN40
A	288	LEU	-	cloning artifact	UNP E9AN40
A	289	GLY	-	cloning artifact	UNP E9AN40
A	290	ALA	-	cloning artifact	UNP E9AN40
A	291	PRO	-	cloning artifact	UNP E9AN40
A	292	ALA	-	cloning artifact	UNP E9AN40
A	293	GLY	-	cloning artifact	UNP E9AN40
A	294	ARG	-	cloning artifact	UNP E9AN40
A	295	GLN	-	cloning artifact	UNP E9AN40
A	296	ALA	-	cloning artifact	UNP E9AN40
A	297	CYS	-	cloning artifact	UNP E9AN40
A	298	GLY	-	cloning artifact	UNP E9AN40
A	299	ARG	-	cloning artifact	UNP E9AN40
A	300	ILE	-	cloning artifact	UNP E9AN40
A	301	MET	-	cloning artifact	UNP E9AN40
A	302	LEU	-	cloning artifact	UNP E9AN40
A	303	LYS	-	cloning artifact	UNP E9AN40
A	304	SER	-	cloning artifact	UNP E9AN40
A	305	ASN	-	cloning artifact	UNP E9AN40
A	306	ARG	-	cloning artifact	UNP E9AN40

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	307	LYS	-	cloning artifact	UNP E9AN40
B	182	GLY	-	expression tag	UNP E9AN40
B	183	SER	-	expression tag	UNP E9AN40
B	218	GLN	ARG	engineered mutation	UNP E9AN40
B	285	GLU	-	cloning artifact	UNP E9AN40
B	286	PHE	-	cloning artifact	UNP E9AN40
B	287	GLU	-	cloning artifact	UNP E9AN40
B	288	LEU	-	cloning artifact	UNP E9AN40
B	289	GLY	-	cloning artifact	UNP E9AN40
B	290	ALA	-	cloning artifact	UNP E9AN40
B	291	PRO	-	cloning artifact	UNP E9AN40
B	292	ALA	-	cloning artifact	UNP E9AN40
B	293	GLY	-	cloning artifact	UNP E9AN40
B	294	ARG	-	cloning artifact	UNP E9AN40
B	295	GLN	-	cloning artifact	UNP E9AN40
B	296	ALA	-	cloning artifact	UNP E9AN40
B	297	CYS	-	cloning artifact	UNP E9AN40
B	298	GLY	-	cloning artifact	UNP E9AN40
B	299	ARG	-	cloning artifact	UNP E9AN40
B	300	ILE	-	cloning artifact	UNP E9AN40
B	301	MET	-	cloning artifact	UNP E9AN40
B	302	LEU	-	cloning artifact	UNP E9AN40
B	303	LYS	-	cloning artifact	UNP E9AN40
B	304	SER	-	cloning artifact	UNP E9AN40
B	305	ASN	-	cloning artifact	UNP E9AN40
B	306	ARG	-	cloning artifact	UNP E9AN40
B	307	LYS	-	cloning artifact	UNP E9AN40

- Molecule 2 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			22	6	14	2		
2	B	1	Total	C	H	O	0	0
			22	6	14	2		

- Molecule 3 is IMIDAZOLE (CCD ID: IMD) (formula: $C_3H_5N_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	N	0	0
			10	3	5	2		
3	A	1	Total	C	H	N	0	0
			10	3	5	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	136	Total 136	O 136	0	0
4	B	128	Total 128	O 128	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: *Leishmania mexicana* KKT4



- Molecule 1: *Leishmania mexicana* KKT4



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	31.31Å 37.71Å 122.39Å 90.00° 92.17° 90.00°	Depositor
Resolution (Å)	61.15 – 1.90 61.15 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.3 (61.15-1.90) 98.4 (61.15-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.84 (at 1.90Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.199 , 0.232 0.199 , 0.231	Depositor DCC
R_{free} test set	1069 reflections (4.67%)	wwPDB-VP
Wilson B-factor (Å ²)	17.6	Xtriage
Anisotropy	0.091	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 59.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	0.010 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	1816	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IMD, MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.95	6/881 (0.7%)	0.87	2/1187 (0.2%)
1	B	1.20	7/617 (1.1%)	0.96	0/831
All	All	1.06	13/1498 (0.9%)	0.91	2/2018 (0.1%)

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	199	GLU	C-O	-8.12	1.14	1.24
1	A	289	GLY	C-O	-7.27	1.16	1.24
1	B	208	ARG	C-O	-6.69	1.16	1.24
1	A	288	LEU	C-O	-5.97	1.16	1.24
1	A	187	LYS	C-O	-5.81	1.16	1.24
1	B	201	GLU	C-O	-5.71	1.17	1.24
1	B	209	VAL	C-O	-5.49	1.18	1.24
1	B	206	THR	C-O	-5.45	1.17	1.24
1	B	203	GLN	C-O	-5.36	1.17	1.24
1	A	188	LEU	C-O	-5.24	1.18	1.24
1	A	286	PHE	C-O	-5.22	1.18	1.24
1	B	202	LEU	C-O	-5.22	1.17	1.24
1	A	186	ASN	C-O	-5.12	1.18	1.24

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	286	PHE	CA-CB-CG	-5.29	108.51	113.80
1	A	285	GLU	O-C-N	5.16	128.03	122.15

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	874	0	836	10	0
1	B	614	0	585	7	0
2	A	8	14	14	4	0
2	B	8	14	14	5	0
3	A	10	10	10	1	0
4	A	136	0	0	4	1
4	B	128	0	0	2	1
All	All	1778	38	1459	19	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:193[A]:ARG:NH1	4:A:901:HOH:O	2.16	0.77
1:A:223:LEU:HD11	1:A:227:LYS:HZ2	1.54	0.71
1:B:213:SER:O	2:B:701:MPD:O2	2.10	0.69
1:A:224:GLN:HG2	2:A:801:MPD:H52	1.74	0.69
1:A:193[B]:ARG:HG3	4:A:971:HOH:O	1.97	0.65
1:A:205:THR:HG22	1:B:205[B]:THR:HG22	1.81	0.63
2:A:801:MPD:HM1	2:A:801:MPD:C5	2.30	0.62
1:A:224:GLN:HG2	2:A:801:MPD:C5	2.33	0.58
2:A:801:MPD:HM1	2:A:801:MPD:H53	1.83	0.58
1:A:223:LEU:HD11	1:A:227:LYS:NZ	2.21	0.55
2:B:701:MPD:H53	4:B:907:HOH:O	2.09	0.52
1:B:222:GLU:HG3	4:B:807:HOH:O	2.11	0.49
1:B:217:GLN:HG2	2:B:701:MPD:H32	1.97	0.47
1:A:223:LEU:HD21	1:B:222:GLU:OE2	2.16	0.45
1:A:221:ASP:OD2	4:A:902:HOH:O	2.21	0.45
3:A:802:IMD:H2	4:A:987:HOH:O	2.17	0.45
1:A:203:GLN:O	1:A:206:THR:HG22	2.20	0.41
1:B:217:GLN:CB	2:B:701:MPD:H32	2.50	0.41
1:B:217:GLN:CG	2:B:701:MPD:H32	2.52	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1008:HOH:O	4:A:1015:HOH:O[1_655]	1.96	0.24
4:B:903:HOH:O	4:B:914:HOH:O[1_655]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	111/126 (88%)	111 (100%)	0	0	100	100
1	B	77/126 (61%)	77 (100%)	0	0	100	100
All	All	188/252 (75%)	188 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	86/104 (83%)	83 (96%)	3 (4%)	32	24
1	B	62/104 (60%)	62 (100%)	0	100	100
All	All	148/208 (71%)	145 (98%)	3 (2%)	59	46

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	207[A]	GLN
1	A	207[B]	GLN
1	A	288	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MPD	B	701	-	7,7,7	0.35	0	9,10,10	1.10	1 (11%)
2	MPD	A	801	-	7,7,7	0.29	0	9,10,10	0.84	0
3	IMD	A	803	-	5,5,5	0.59	0	5,5,5	0.43	0
3	IMD	A	802	-	5,5,5	0.63	0	5,5,5	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MPD	B	701	-	-	0/5/5/5	-
2	MPD	A	801	-	-	2/5/5/5	-
3	IMD	A	803	-	-	-	0/1/1/1
3	IMD	A	802	-	-	-	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	701	MPD	CM-C2-C1	-2.08	105.98	110.63

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	801	MPD	C2-C3-C4-O4
2	A	801	MPD	C2-C3-C4-C5

There are no ring outliers.

3 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	701	MPD	5	0
2	A	801	MPD	4	0
3	A	802	IMD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	110/126 (87%)	0.66	12 (10%) 10 11	7, 29, 56, 60	3 (2%)
1	B	77/126 (61%)	0.21	3 (3%) 43 46	4, 21, 55, 60	2 (2%)
All	All	187/252 (74%)	0.47	15 (8%) 18 19	4, 25, 56, 60	5 (2%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	290	ALA	4.1
1	A	288	LEU	4.1
1	A	272	TYR	3.8
1	A	291	PRO	3.8
1	A	289	GLY	3.5
1	B	257	ARG	3.1
1	A	259	LEU	2.7
1	B	258	GLN	2.7
1	B	222	GLU	2.6
1	A	266	MET	2.5
1	A	286	PHE	2.3
1	A	284	GLN	2.3
1	A	260	LEU	2.2
1	A	222	GLU	2.1
1	A	250	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MPD	B	701	8/8	0.74	0.22	20,51,62,70	0
2	MPD	A	801	8/8	0.76	0.22	59,75,80,90	0
3	IMD	A	803	5/5	0.78	0.24	69,69,83,83	0
3	IMD	A	802	5/5	0.85	0.15	55,58,70,70	0

6.5 Other polymers [i](#)

There are no such residues in this entry.