



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 11:09 AM UTC

PDB ID : 6ZPX / pdb_00006zpx
Title : Structure of Unliganded MgGH51 α -L-Arabinofuranosidase Crystal Type 1
Authors : McGregor, N.G.S.; Davies, G.J.
Deposited on : 2020-07-09
Resolution : 1.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1553 (1.30-1.30)
Clashscore	190562	1595 (1.30-1.30)
Ramachandran outliers	187476	1551 (1.30-1.30)
Sidechain outliers	187428	1551 (1.30-1.30)
RSRZ outliers	180081	1549 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	627	 91% 8%
2	A	2	 100%
2	B	2	 100%
2	D	2	 100%
3	C	3	 100%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 10353 atoms, of which 4754 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MgGH51.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	AAA	627	Total	C	H	N	O	S	310	12	0
			9468	3093	4636	798	934	7			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	A	2	Total	C	H	N	O		5	0	0
			55	16	27	2	10				
2	B	2	Total	C	H	N	O		5	0	0
			55	16	27	2	10				
2	D	2	Total	C	H	N	O		5	0	0
			55	16	27	2	10				

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	3	Total	C	H	N	O		8	0	0
			76	22	37	2	15				

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	1	Total 1	Cl 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	AAA	643	Total 643	O 643	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MgGH51




- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



MAG1
MAG2
BMA3

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.98Å 65.80Å 193.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	62.36 – 1.30 62.36 – 1.30	Depositor EDS
% Data completeness (in resolution range)	99.9 (62.36-1.30) 99.9 (62.36-1.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.69 (at 1.30Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.178 , 0.217 0.178 , 0.214	Depositor DCC
R_{free} test set	9384 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	14.8	Xtriage
Anisotropy	0.441	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 34.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10353	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, CL, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.26	23/4994 (0.5%)	1.23	8/6842 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	AAA	0	3

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	AAA	139	GLY	C-O	6.99	1.33	1.24
1	AAA	337	HIS	CE1-NE2	6.74	1.39	1.32
1	AAA	442	GLY	C-O	6.23	1.29	1.23
1	AAA	485	HIS	CE1-NE2	6.12	1.38	1.32
1	AAA	29	GLY	N-CA	6.05	1.49	1.45
1	AAA	365	ASP	C-O	5.91	1.31	1.24
1	AAA	546	GLY	C-O	5.90	1.32	1.24
1	AAA	260	ASP	C-O	5.84	1.31	1.24
1	AAA	500	ALA	C-O	5.78	1.31	1.24
1	AAA	190	ILE	C-O	5.77	1.29	1.24
1	AAA	77	ALA	C-O	5.56	1.31	1.24
1	AAA	479	TYR	C-O	5.50	1.30	1.23
1	AAA	115	SER	C-O	5.29	1.30	1.23
1	AAA	85[A]	SER	C-O	5.28	1.30	1.23
1	AAA	85[B]	SER	C-O	5.28	1.30	1.23
1	AAA	320	ILE	N-CA	5.20	1.51	1.46
1	AAA	65	LEU	C-O	5.17	1.30	1.24
1	AAA	171	ASP	C-O	5.16	1.29	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	AAA	590	THR	N-CA	5.15	1.49	1.46
1	AAA	84	PHE	C-O	5.11	1.30	1.24
1	AAA	615	PRO	C-O	5.08	1.29	1.23
1	AAA	213	ILE	C-O	-5.06	1.18	1.24
1	AAA	473	ILE	N-CA	5.04	1.51	1.46

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	93	ARG	CB-CA-C	7.66	122.59	110.19
1	AAA	139	GLY	O-C-N	7.07	130.60	122.33
1	AAA	139	GLY	CA-C-O	-6.83	111.39	118.97
1	AAA	6	ASN	CA-CB-CG	6.66	119.26	112.60
1	AAA	426	PHE	CA-CB-CG	6.43	120.23	113.80
1	AAA	314	ASP	CA-CB-CG	6.09	118.69	112.60
1	AAA	39	GLN	CB-CA-C	5.35	118.57	109.53
1	AAA	260	ASP	CA-CB-CG	-5.12	107.48	112.60

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	AAA	220[A]	MET	Mainchain
1	AAA	359	TYR	Sidechain
1	AAA	65	LEU	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	4832	4636	4615	20	0
2	A	28	27	25	0	0
2	B	28	27	25	0	0
2	D	28	27	25	0	0
3	C	39	37	34	0	0
4	AAA	1	0	0	0	0
5	AAA	643	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5599	4754	4724	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:64:LYS:HB2	1:AAA:85[B]:SER:OG	2.01	0.60
1:AAA:119:ARG:NH1	5:AAA:805:HOH:O	2.39	0.54
1:AAA:364:HIS:HE1	5:AAA:1334:HOH:O	1.95	0.49
1:AAA:401:VAL:O	1:AAA:430:TYR:HA	2.13	0.48
1:AAA:515:PHE:CD2	1:AAA:551:LYS:HE3	2.49	0.47
1:AAA:221:GLY:N	1:AAA:222:PRO:CD	2.80	0.45
1:AAA:241:ARG:HD2	1:AAA:290:TRP:CD2	2.52	0.45
1:AAA:569:PHE:CD1	1:AAA:626:THR:HB	2.50	0.45
1:AAA:220[B]:MET:HE2	1:AAA:509:PHE:CE1	2.53	0.44
1:AAA:347:GLU:OE2	1:AAA:398:ASP:OD2	2.37	0.43
1:AAA:28:SER:HA	1:AAA:498:TYR:OH	2.18	0.43
1:AAA:291:ALA:O	1:AAA:362:ARG:HD2	2.20	0.42
1:AAA:159[B]:ILE:HD13	5:AAA:989:HOH:O	2.18	0.42
1:AAA:220[A]:MET:HG2	1:AAA:509:PHE:CE1	2.55	0.42
1:AAA:220[B]:MET:HE2	1:AAA:509:PHE:HE1	1.85	0.42
1:AAA:116:LEU:HD12	1:AAA:116:LEU:C	2.45	0.42
1:AAA:290:TRP:CH2	1:AAA:292:GLY:HA2	2.54	0.41
1:AAA:396:SER:HA	1:AAA:424:THR:O	2.21	0.41
1:AAA:426:PHE:CE1	1:AAA:428:GLY:HA2	2.55	0.41
1:AAA:226:ARG:HA	1:AAA:286:ILE:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	637/627 (102%)	621 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	514/510 (101%)	512 (100%)	2 (0%)	84	65

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	93	ARG
1	AAA	376	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

9 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	1	1,2	14,14,15	1.01	1 (7%)	17,19,21	1.06	1 (5%)
2	NAG	A	2	2	14,14,15	1.25	1 (7%)	17,19,21	1.18	2 (11%)
2	NAG	B	1	1,2	14,14,15	1.19	2 (14%)	17,19,21	1.46	4 (23%)
2	NAG	B	2	2	14,14,15	1.05	0	17,19,21	1.56	3 (17%)
3	NAG	C	1	1,3	14,14,15	0.95	0	17,19,21	1.16	1 (5%)
3	NAG	C	2	3	14,14,15	0.95	1 (7%)	17,19,21	1.42	3 (17%)
3	BMA	C	3	3	11,11,12	1.20	1 (9%)	15,15,17	2.66	6 (40%)
2	NAG	D	1	1,2	14,14,15	1.24	2 (14%)	17,19,21	1.43	4 (23%)
2	NAG	D	2	2	14,14,15	0.80	0	17,19,21	1.49	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	2	2	-	1/6/23/26	0/1/1/1
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
3	NAG	C	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	BMA	C	3	3	-	0/2/19/22	0/1/1/1
2	NAG	D	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	0/6/23/26	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	NAG	C1-C2	3.22	1.56	1.52
2	A	2	NAG	O4-C4	3.16	1.50	1.43
3	C	3	BMA	C2-C3	2.84	1.56	1.52
2	D	1	NAG	C1-C2	2.77	1.56	1.52
3	C	2	NAG	O4-C4	2.43	1.49	1.43
2	A	1	NAG	C1-C2	2.30	1.55	1.52
2	D	1	NAG	O5-C5	2.15	1.47	1.43
2	B	1	NAG	O7-C7	-2.04	1.18	1.23

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	3	BMA	O3-C3-C4	6.61	125.96	110.38
3	C	3	BMA	C1-C2-C3	4.60	116.34	109.64
2	D	2	NAG	O5-C1-C2	-3.47	105.92	111.29
3	C	3	BMA	O2-C2-C1	-3.28	101.71	109.22
3	C	3	BMA	O4-C4-C5	-3.09	101.72	109.32
2	B	1	NAG	C6-C5-C4	3.04	120.48	113.02
2	D	1	NAG	O7-C7-N2	3.01	127.30	121.98
2	D	2	NAG	O6-C6-C5	-2.98	101.17	111.33
2	D	1	NAG	O5-C5-C4	-2.94	103.68	110.83
2	B	2	NAG	O5-C5-C6	2.94	113.38	107.66
2	B	2	NAG	C4-C3-C2	2.85	115.19	111.02
3	C	2	NAG	O5-C5-C6	2.54	112.60	107.66
3	C	3	BMA	C2-C3-C4	-2.50	106.47	110.86
2	D	2	NAG	C3-C4-C5	-2.46	105.78	110.23
2	D	1	NAG	C8-C7-N2	-2.36	112.21	116.12
3	C	1	NAG	C1-O5-C5	2.34	115.32	112.19
3	C	2	NAG	O6-C6-C5	-2.31	103.48	111.33
3	C	2	NAG	O5-C5-C4	-2.24	105.39	110.83
2	B	2	NAG	C3-C4-C5	-2.20	106.24	110.23
2	A	1	NAG	C1-C2-N2	-2.19	106.98	110.43
2	D	1	NAG	O5-C1-C2	-2.18	107.92	111.29
2	A	2	NAG	O4-C4-C5	2.12	114.54	109.32
2	A	2	NAG	C3-C4-C5	-2.12	106.39	110.23
3	C	3	BMA	C3-C4-C5	2.09	114.02	110.23
2	B	1	NAG	C8-C7-N2	2.06	119.54	116.12
2	B	1	NAG	O5-C5-C4	-2.06	105.81	110.83
2	B	1	NAG	O7-C7-N2	-2.00	118.44	121.98

There are no chirality outliers.


All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	2	NAG	O5-C5-C6-O6


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	627/627 (100%)	0.07	3 (0%) 87 90	8, 18, 28, 46	13 (2%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	138	ALA	3.3
1	AAA	139	GLY	2.3
1	AAA	6	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	A	1	14/15	-	-	21,26,30,30	2
2	NAG	A	2	14/15	-	-	28,34,38,44	3
3	NAG	C	2	14/15	0.89	0.10	19,25,31,32	2
2	NAG	B	2	14/15	0.93	0.08	30,41,48,55	3
2	NAG	B	1	14/15	0.94	0.07	23,26,31,31	2
3	NAG	C	1	14/15	0.95	0.07	17,19,23,33	2
2	NAG	D	2	14/15	0.96	0.07	25,42,47,58	3
2	NAG	D	1	14/15	0.98	0.05	18,21,23,25	2
3	BMA	C	3	11/12	-	-	38,48,53,54	4

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around Chain B:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	CL	AAA	710	1/1	0.90	0.18	33,33,33,33	1

6.5 Other polymers [i](#)

There are no such residues in this entry.