



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:26 AM UTC

PDB ID : 2ZRJ / pdb_00002zrj
Title : MsRecA Q196A ATPgS form IV
Authors : Prabu, J.R.; Manjunath, G.P.; Chandra, N.R.; Muniyappa, K.; Vijayan, M.
Deposited on : 2008-08-27
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	349	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	502	-	X	-	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	503	-	X	-	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 2425 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Protein recA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	314	Total	C	N	O	S	0	0	0
			2294	1437	405	446	6			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	196	ALA	GLN	engineered mutation	UNP Q59560

- Molecule 2 is PHOSPHOTHIOPHOSPHORIC ACID-ADENYLATE ESTER (CCD ID: AGS) (formula: $C_{10}H_{16}N_5O_{12}P_3S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P S	0	0
			31	10	5	12	3 1		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is 2-{2-[2-(2-{2-[2-(2-ETHOXY-ETHOXY)-ETHOXY]-ETHOXY}-ETHOXY)-ETHOXY]-ETHOXY}-ETHANOL (CCD ID: PE4) (formula: $C_{16}H_{34}O_8$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			24	16	8		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	63	Total	O	0	0
			63	63		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Protein recA



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	101.90Å 101.90Å 75.33Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	24.97 – 2.60 24.97 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.9 (24.97-2.60) 99.8 (24.97-2.60)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.83 (at 2.60Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.218 , 0.252 0.218 , 0.252	Depositor DCC
R_{free} test set	1378 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	49.8	Xtriage
Anisotropy	0.760	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 44.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.044 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2425	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PE4, AGS, CL, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/2320	1.09	21/3134 (0.7%)

There are no bond length outliers.

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	147	SER	N-CA-C	9.87	131.83	110.80
1	A	322	GLU	N-CA-C	-8.69	101.75	111.14
1	A	289	SER	N-CA-C	-8.01	98.63	110.48
1	A	40	SER	N-CA-C	-7.46	97.20	108.46
1	A	168	LEU	N-CA-C	7.31	119.04	111.14
1	A	180	MET	N-CA-C	7.11	119.11	111.36
1	A	146	ASP	CB-CA-C	6.33	120.17	109.84
1	A	45	GLY	N-CA-C	-6.30	106.73	114.92
1	A	236	GLY	N-CA-C	-6.16	98.57	113.18
1	A	6	PRO	N-CA-C	6.12	120.10	110.50
1	A	222	SER	N-CA-C	-6.11	105.99	113.50
1	A	96	ASP	N-CA-C	5.78	118.14	108.73
1	A	161	MET	N-CA-C	5.77	127.14	111.00
1	A	238	ASP	N-CA-C	5.72	118.27	110.55
1	A	160	GLU	N-CA-C	5.59	119.86	111.81
1	A	214	GLY	N-CA-C	-5.52	108.27	114.40
1	A	167	GLY	N-CA-C	-5.49	108.06	115.36
1	A	147	SER	CB-CA-C	-5.40	99.68	110.42
1	A	120	GLN	CA-C-N	5.29	125.58	119.92
1	A	120	GLN	C-N-CA	5.29	125.58	119.92
1	A	284	GLY	N-CA-C	5.26	125.64	113.18

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2294	0	2260	96	0
2	A	31	0	12	2	0
3	A	1	0	0	1	0
4	A	12	0	8	2	0
5	A	24	0	34	1	0
6	A	63	0	0	2	0
All	All	2425	0	2314	98	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

All (98) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:12:LEU:HG	1:A:16:MET:HE2	1.31	1.13
1:A:151:LEU:HD12	1:A:173:MET:HE1	1.30	1.12
1:A:151:LEU:CD1	1:A:173:MET:HE1	1.81	1.09
1:A:155:ALA:HB1	1:A:160:GLU:HG3	1.42	1.02
1:A:44:THR:HG23	1:A:46:SER:H	1.30	0.95
1:A:146:ASP:OD1	4:A:503:GOL:H11	1.78	0.84
1:A:229:ARG:HD2	1:A:242:ASN:HB3	1.61	0.83
1:A:44:THR:HG22	1:A:50:ASP:OD2	1.84	0.78
1:A:80:HIS:CE1	1:A:270:ILE:H	2.03	0.77
1:A:291:SER:O	1:A:302:GLN:HG3	1.85	0.76
1:A:80:HIS:HE1	1:A:270:ILE:H	1.35	0.74
1:A:44:THR:HG21	1:A:49:LEU:HB3	1.68	0.74
1:A:155:ALA:O	1:A:160:GLU:HG2	1.89	0.72
1:A:98:GLU:OE2	1:A:150:ALA:HB2	1.89	0.71
1:A:151:LEU:HD13	1:A:173:MET:HE1	1.70	0.69
1:A:151:LEU:HD12	1:A:173:MET:CE	2.18	0.68
1:A:317:VAL:O	1:A:318:ALA:HB3	1.94	0.68
1:A:63:VAL:C	1:A:64:ILE:HD12	2.18	0.67
1:A:278:ASP:O	1:A:281:VAL:HG12	1.95	0.67
1:A:80:HIS:CE1	1:A:270:ILE:HG12	2.31	0.66
1:A:291:SER:O	1:A:302:GLN:HA	1.98	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:292:TRP:CE2	1:A:302:GLN:HB2	2.34	0.63
1:A:280:GLY:C	1:A:286:ILE:HG23	2.24	0.62
1:A:323:LYS:O	1:A:327:GLU:HG3	1.98	0.62
1:A:328:LYS:C	1:A:329:LEU:HD12	2.26	0.61
1:A:219:PHE:O	5:A:600:PE4:H22	2.01	0.61
1:A:155:ALA:CB	1:A:160:GLU:HG3	2.26	0.61
1:A:325:ILE:O	1:A:329:LEU:HD13	2.02	0.59
1:A:109:LEU:HD22	1:A:269:GLY:HA3	1.85	0.59
1:A:144:VAL:HG22	1:A:192:ILE:HB	1.85	0.59
1:A:44:THR:HG23	1:A:46:SER:N	2.10	0.58
1:A:9:GLU:O	1:A:13:GLU:HG2	2.04	0.58
1:A:44:THR:OG1	1:A:49:LEU:HD23	2.03	0.58
1:A:233:LEU:O	1:A:240:VAL:HG22	2.04	0.57
1:A:145:ILE:N	1:A:145:ILE:HD12	2.20	0.57
1:A:285:PHE:CE2	1:A:324:LYS:HB3	2.40	0.57
1:A:294:THR:HG22	1:A:299:GLN:NE2	2.21	0.56
1:A:60:ARG:HA	1:A:190:THR:OG1	2.06	0.55
1:A:290:GLY:O	1:A:291:SER:HB2	2.06	0.55
1:A:123:THR:HG22	1:A:157:ILE:HD12	1.89	0.55
1:A:44:THR:CG2	1:A:46:SER:H	2.13	0.55
1:A:291:SER:O	1:A:302:GLN:CG	2.53	0.55
1:A:147:SER:HB3	1:A:150:ALA:HB3	1.89	0.54
1:A:64:ILE:HD12	1:A:64:ILE:N	2.23	0.53
1:A:98:GLU:CG	1:A:147:SER:HB2	2.38	0.53
1:A:321:ILE:O	1:A:325:ILE:HG13	2.10	0.52
2:A:500:AGS:O2G	2:A:500:AGS:O2B	2.29	0.51
1:A:64:ILE:HD13	1:A:190:THR:HG23	1.91	0.51
1:A:329:LEU:HD12	1:A:329:LEU:N	2.26	0.51
1:A:272:ARG:HH21	1:A:322:GLU:CD	2.19	0.51
1:A:5:ALA:HB3	1:A:6:PRO:HD3	1.93	0.51
1:A:171:ARG:HG3	1:A:175:GLN:NE2	2.26	0.50
1:A:80:HIS:HE1	1:A:270:ILE:N	2.08	0.50
1:A:62:ARG:HD2	1:A:222:SER:OG	2.11	0.49
1:A:102:ASP:OD1	1:A:102:ASP:C	2.56	0.49
1:A:43:PRO:HB3	3:A:501:CL:CL	2.49	0.49
1:A:16:MET:HE1	1:A:30:ARG:NH1	2.28	0.49
1:A:328:LYS:HB3	1:A:329:LEU:HD12	1.95	0.48
1:A:232:THR:HG23	1:A:232:THR:O	2.13	0.48
1:A:126:GLN:O	1:A:130:ILE:HG13	2.14	0.48
1:A:318:ALA:O	1:A:322:GLU:N	2.45	0.48
1:A:320:GLU:O	1:A:323:LYS:HB3	2.14	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:286:ILE:HD11	1:A:293:PHE:HB3	1.96	0.47
1:A:317:VAL:HG12	1:A:318:ALA:H	1.80	0.47
1:A:262:PHE:CE1	1:A:270:ILE:HD12	2.50	0.47
1:A:233:LEU:HD11	1:A:263:ASP:HB3	1.97	0.47
1:A:272:ARG:NE	1:A:322:GLU:OE2	2.42	0.46
1:A:152:VAL:HG23	1:A:157:ILE:HD11	1.98	0.46
1:A:152:VAL:HG23	1:A:157:ILE:CG1	2.46	0.45
1:A:210:THR:HA	1:A:214:GLY:HA3	1.98	0.45
1:A:33:GLU:HG2	6:A:625:HOH:O	2.16	0.45
1:A:36:ARG:O	1:A:38:PRO:HD3	2.17	0.44
1:A:108:LYS:HD3	1:A:267:GLY:O	2.17	0.44
1:A:30:ARG:HD2	6:A:616:HOH:O	2.18	0.44
1:A:98:GLU:HG2	1:A:147:SER:HB2	2.00	0.43
1:A:121:PRO:HG2	1:A:127:ALA:HA	1.99	0.43
1:A:35:VAL:HG11	1:A:254:SER:OG	2.18	0.43
1:A:128:LEU:HD13	1:A:176:ALA:CB	2.47	0.43
1:A:13:GLU:OE1	1:A:16:MET:HE3	2.18	0.42
1:A:100:ALA:O	1:A:101:LEU:C	2.62	0.42
1:A:243:ARG:HH22	1:A:278:ASP:CG	2.27	0.42
1:A:5:ALA:N	1:A:6:PRO:CD	2.82	0.42
1:A:213:GLY:C	1:A:215:LYS:N	2.76	0.42
1:A:67:TYR:HB3	1:A:195:ASN:HB3	2.02	0.42
1:A:281:VAL:HG13	1:A:282:GLU:N	2.35	0.41
1:A:121:PRO:HG2	1:A:127:ALA:CB	2.50	0.41
1:A:279:MET:HA	1:A:282:GLU:HG2	2.02	0.41
2:A:500:AGS:O2G	4:A:503:GOL:H12	2.20	0.41
1:A:317:VAL:O	1:A:318:ALA:CB	2.60	0.41
1:A:316:ASP:C	1:A:317:VAL:O	2.61	0.41
1:A:147:SER:O	1:A:148:VAL:C	2.64	0.41
1:A:273:GLU:CD	1:A:273:GLU:H	2.29	0.41
1:A:121:PRO:CG	1:A:127:ALA:HA	2.51	0.41
1:A:213:GLY:C	1:A:215:LYS:H	2.27	0.41
1:A:326:LYS:O	1:A:330:GLY:N	2.53	0.40
1:A:20:ASP:O	1:A:24:GLY:N	2.43	0.40
1:A:119:SER:O	1:A:121:PRO:HD3	2.22	0.40
1:A:293:PHE:O	1:A:300:LEU:HB2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	308/349 (88%)	282 (92%)	24 (8%)	2 (1%)	21	42

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	39	ILE
1	A	166	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/274 (82%)	221 (98%)	5 (2%)	45	72

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	43	PRO
1	A	141	ASP
1	A	171	ARG
1	A	234	LYS
1	A	286	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
1	A	80	HIS
1	A	126	GLN
1	A	169	GLN
1	A	175	GLN
1	A	299	GLN
1	A	319	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	AGS	A	500	-	32,33,33	1.27	4 (12%)	45,52,52	1.88	10 (22%)
5	PE4	A	600	-	23,23,23	0.54	0	22,22,22	0.53	0
4	GOL	A	503	-	5,5,5	1.43	1 (20%)	5,5,5	2.36	5 (100%)
4	GOL	A	502	-	5,5,5	1.39	1 (20%)	5,5,5	2.37	5 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AGS	A	500	-	-	4/21/38/38	0/3/3/3
5	PE4	A	600	-	-	10/21/21/21	-
4	GOL	A	503	-	-	2/4/4/4	-
4	GOL	A	502	-	-	3/4/4/4	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	AGS	C5-N7	-3.23	1.33	1.39
4	A	503	GOL	O2-C2	-2.41	1.36	1.43
2	A	500	AGS	C8-N9	-2.36	1.33	1.37
4	A	502	GOL	O2-C2	-2.30	1.36	1.43
2	A	500	AGS	PA-O3A	2.20	1.61	1.59
2	A	500	AGS	PB-O3A	2.12	1.61	1.59

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	AGS	C5-C4-N3	-5.00	119.83	126.72
2	A	500	AGS	N3-C2-N1	-4.65	121.54	128.58
2	A	500	AGS	PB-O3B-PG	-4.23	117.69	133.17
2	A	500	AGS	O3G-PG-O3B	3.84	117.46	104.64
2	A	500	AGS	N3-C4-N9	3.54	133.19	127.17
2	A	500	AGS	C2-N3-C4	3.50	120.39	111.83
2	A	500	AGS	N9-C8-N7	-2.90	109.83	113.94
2	A	500	AGS	O2G-PG-O3B	2.72	113.71	104.64
2	A	500	AGS	C5-N7-C8	2.65	107.61	103.45
4	A	502	GOL	O2-C2-C3	2.58	119.84	109.18
4	A	502	GOL	O2-C2-C1	2.56	119.80	109.18
4	A	503	GOL	O2-C2-C3	2.56	119.78	109.18
4	A	503	GOL	O2-C2-C1	2.52	119.61	109.18
2	A	500	AGS	C4-C5-N7	-2.47	107.75	110.58
4	A	503	GOL	C3-C2-C1	2.40	120.61	111.80
4	A	502	GOL	C3-C2-C1	2.33	120.33	111.80
4	A	503	GOL	O3-C3-C2	2.24	120.45	110.38
4	A	502	GOL	O3-C3-C2	2.22	120.36	110.38
4	A	502	GOL	O1-C1-C2	2.15	120.05	110.38
4	A	503	GOL	O1-C1-C2	2.07	119.69	110.38

There are no chirality outliers.

All (19) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	500	AGS	PB-O3B-PG-O2G
4	A	502	GOL	C1-C2-C3-O3
4	A	503	GOL	C1-C2-C3-O3
5	A	600	PE4	C10-C9-O5-C8
5	A	600	PE4	C11-C12-O7-C13
5	A	600	PE4	O3-C5-C6-O4
5	A	600	PE4	O4-C7-C8-O5
4	A	502	GOL	O1-C1-C2-O2
4	A	503	GOL	O1-C1-C2-O2
5	A	600	PE4	O6-C11-C12-O7
5	A	600	PE4	O6-C10-C9-O5
2	A	500	AGS	PA-O3A-PB-O2B
5	A	600	PE4	C13-C14-O8-C15
5	A	600	PE4	C9-C10-O6-C11
5	A	600	PE4	C6-C5-O3-C4
4	A	502	GOL	O2-C2-C3-O3
2	A	500	AGS	PG-O3B-PB-O2B
2	A	500	AGS	PA-O3A-PB-O1B
5	A	600	PE4	C12-C11-O6-C10

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	AGS	2	0
5	A	600	PE4	1	0
4	A	503	GOL	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	314/349 (89%)	0.15	27 (8%) 16 12	33, 52, 95, 114	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	331	ILE	5.1
1	A	39	ILE	4.6
1	A	289	SER	3.9
1	A	297	GLY	3.8
1	A	236	GLY	3.6
1	A	210	THR	3.6
1	A	235	ASP	3.4
1	A	5	ALA	3.3
1	A	166	VAL	2.9
1	A	161	MET	2.8
1	A	299	GLN	2.8
1	A	211	THR	2.8
1	A	330	GLY	2.7
1	A	209	GLU	2.7
1	A	147	SER	2.6
1	A	292	TRP	2.6
1	A	198	ARG	2.5
1	A	35	VAL	2.5
1	A	237	THR	2.4
1	A	160	GLU	2.4
1	A	37	GLN	2.4
1	A	38	PRO	2.3
1	A	159	GLY	2.3
1	A	212	THR	2.3
1	A	284	GLY	2.1
1	A	6	PRO	2.1
1	A	168	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	A	502	6/6	0.64	0.14	78,86,87,87	0
5	PE4	A	600	24/24	0.70	0.22	45,72,90,93	0
2	AGS	A	500	31/31	0.84	0.16	43,64,71,87	31
3	CL	A	501	1/1	0.88	0.12	71,71,71,71	0
4	GOL	A	503	6/6	0.92	0.14	56,60,63,64	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around PE4 A 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.