



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:40 PM UTC

PDB ID : 3ZRB / pdb_00003zrb
Title : Structural basis for agonism and antagonism for a set of chemically related progesterone receptor modulators
Authors : Lusher, S.J.; Raaijmakers, H.C.A.; Vu-Pham, D.; Dechering, K.; Wai Lam, T.; Brown, A.R.; Hamilton, N.M.; Nimz, O.; Azevedo, R.; McGuire, R.; Oubrie, A.; de Vlieg, J.
Deposited on : 2011-06-15
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	260	3% 88% 8% .
1	B	260	8% 91% 5% .

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4384 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PROGESTERONE RECEPTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	250	Total	C	N	O	S	0	4	0
			2048	1326	340	367	15			
1	B	250	Total	C	N	O	S	0	2	0
			2029	1315	335	364	15			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	674	GLY	-	expression tag	UNP P06401
A	675	SER	-	expression tag	UNP P06401
A	676	HIS	-	expression tag	UNP P06401
A	677	MET	-	expression tag	UNP P06401
B	674	GLY	-	expression tag	UNP P06401
B	675	SER	-	expression tag	UNP P06401
B	676	HIS	-	expression tag	UNP P06401
B	677	MET	-	expression tag	UNP P06401

- Molecule 2 is 2-CHLORO-N-[[4-(3,5-DIMETHYLISOXAZOL-4-YL)PHENYL]METHYL]-1,4-DIMETHYL-1H-PYRAZOLE-4-SULFONAMIDE (CCD ID: OR8) (formula: C₁₇H₁₉ClN₄O₃S).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	Cl	N	O	S	0	0
			26	17	1	4	3	1		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	O S	0	0
			5	4 1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is (R)-N-[1-[4-(3,5-DIMETHYLISOXAZOL-4-YL)PHENYL]ETHYL]-3,5-DIMETHYLISOXAZOLE-4-SULFONAMIDE (CCD ID: ORC) (formula: $C_{18}H_{21}N_3O_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	S	0	0
			26	18	3	4	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	138	Total	O	0	0
			138	138		
6	B	88	Total	O	0	0
			88	88		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: PROGESTERONE RECEPTOR

Chain A: 



• Molecule 1: PROGESTERONE RECEPTOR

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.25Å 64.06Å 69.69Å 90.00° 96.62° 90.00°	Depositor
Resolution (Å)	69.17 – 1.80 69.17 – 1.80	Depositor EDS
% Data completeness (in resolution range)	96.1 (69.17-1.80) 96.1 (69.17-1.80)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.95 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.184 , 0.221 0.205 , 0.242	Depositor DCC
R_{free} test set	2244 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	22.3	Xtriage
Anisotropy	0.130	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 36.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4384	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4, OR8, ORC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/2098	0.79	0/2837
1	B	0.41	0/2076	0.77	0/2808
All	All	0.46	0/4174	0.78	0/5645

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2048	0	2103	23	0
1	B	2029	0	2080	15	0
2	A	26	0	19	0	0
3	A	5	0	0	1	0
4	A	18	0	24	2	0
4	B	6	0	8	0	0
5	B	26	0	21	0	0
6	A	138	0	0	2	0
6	B	88	0	0	0	0
All	All	4384	0	4255	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:892:LEU:HD21	1:B:892:LEU:HD21	1.57	0.87
1:A:692[B]:MET:SD	1:A:815:GLN:OE1	2.42	0.78
1:B:789:MET:HE3	1:B:798:CYS:HB3	1.69	0.74
1:B:876:LEU:C	1:B:876:LEU:HD23	2.18	0.69
1:B:692[B]:MET:O	1:B:692[B]:MET:HE3	1.97	0.64
1:B:789:MET:HE3	1:B:798:CYS:CB	2.32	0.59
1:A:692[B]:MET:HE1	6:A:2131:HOH:O	2.02	0.58
1:A:896:ILE:HD11	1:B:892:LEU:HD11	1.88	0.56
1:A:705[A]:ASN:CG	1:A:705[A]:ASN:O	2.50	0.54
1:A:860:GLN:CG	1:A:869:ARG:HD3	2.39	0.53
1:A:725[A]:GLN:NE2	6:A:2025:HOH:O	2.41	0.53
1:A:705[A]:ASN:O	1:A:705[A]:ASN:ND2	2.42	0.53
1:A:860:GLN:HG3	1:A:869:ARG:HD3	1.92	0.52
1:A:704:ASP:OD1	1:A:706:THR:HG23	2.10	0.52
1:A:829:THR:HG21	4:A:1935:GOL:H31	1.95	0.49
1:A:737:PRO:HA	3:A:1933:SO4:O3	2.16	0.46
1:B:876:LEU:HD23	1:B:876:LEU:O	2.16	0.45
1:B:730:VAL:HG21	1:B:908:MET:HE1	1.98	0.45
1:A:723:GLU:CD	1:A:906:PRO:HB2	2.42	0.45
1:B:876:LEU:C	1:B:876:LEU:CD2	2.88	0.44
1:A:922:ALA:HB2	1:B:921:LEU:HD22	1.99	0.44
1:A:883:LEU:C	1:A:883:LEU:HD23	2.43	0.44
1:A:892:LEU:HG	1:A:896:ILE:HD12	1.99	0.44
1:B:931:HIS:O	1:B:932:LYS:C	2.60	0.44
1:A:690:LEU:C	1:A:690:LEU:HD23	2.43	0.43
1:A:896:ILE:CD1	1:B:892:LEU:HD11	2.49	0.43
1:A:829:THR:HG21	4:A:1935:GOL:C3	2.48	0.43
1:B:760:VAL:HG22	1:B:804:ILE:HD12	2.01	0.43
1:A:892:LEU:HD21	1:B:892:LEU:CD2	2.39	0.43
1:A:885:LYS:NZ	1:B:922:ALA:O	2.49	0.42
1:A:888:HIS:CD2	1:A:917:LEU:HD11	2.55	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	252/260 (97%)	243 (96%)	9 (4%)	0	100	100
1	B	250/260 (96%)	242 (97%)	8 (3%)	0	100	100
All	All	502/520 (96%)	485 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/241 (98%)	234 (99%)	2 (1%)	73	70
1	B	233/241 (97%)	231 (99%)	2 (1%)	70	67
All	All	469/482 (97%)	465 (99%)	4 (1%)	70	67

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	714	LEU
1	A	802	TRP
1	B	833	GLU
1	B	908	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	752	GLN
1	A	806	GLN
1	A	868	GLN
1	A	879	ASN
1	A	893	ASN
1	B	803	GLN
1	B	868	GLN
1	B	879	ASN
1	B	886	GLN
1	B	893	ASN
1	B	916	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	1933	-	4,4,4	0.34	0	6,6,6	0.36	0
4	GOL	A	1934	-	5,5,5	0.45	0	5,5,5	0.29	0
4	GOL	A	1935	-	5,5,5	0.46	0	5,5,5	0.69	0
5	ORC	B	1000	-	25,28,28	3.25	2 (8%)	31,42,42	2.75	13 (41%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	OR8	A	1000	-	25,28,28	2.41	3 (12%)	29,42,42	3.10	12 (41%)
4	GOL	A	1936	-	5,5,5	0.35	0	5,5,5	0.23	0
4	GOL	B	1933	-	5,5,5	0.36	0	5,5,5	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1934	-	-	2/4/4/4	-
4	GOL	A	1935	-	-	2/4/4/4	-
5	ORC	B	1000	-	-	0/17/19/19	0/3/3/3
2	OR8	A	1000	-	-	3/14/16/16	0/3/3/3
4	GOL	A	1936	-	-	1/4/4/4	-
4	GOL	B	1933	-	-	4/4/4/4	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	1000	ORC	O23-N22	-11.55	1.21	1.42
5	B	1000	ORC	O15-N16	-11.15	1.22	1.42
2	A	1000	OR8	O10-N9	-10.96	1.22	1.42
2	A	1000	OR8	S14-N13	3.09	1.66	1.61
2	A	1000	OR8	C18-CL	2.16	1.74	1.70

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1000	OR8	O10-C11-C22	7.62	128.37	116.59
2	A	1000	OR8	O17-S14-N13	-6.36	101.46	107.41
2	A	1000	OR8	O10-N9-C8	6.22	112.66	106.20
5	B	1000	ORC	O15-C14-C18	5.83	125.60	116.59
5	B	1000	ORC	O23-N22-C21	5.75	112.18	106.20
5	B	1000	ORC	O23-C24-C26	5.69	125.38	116.59
2	A	1000	OR8	C22-C11-C7	-5.19	126.78	133.91
5	B	1000	ORC	O15-N16-C17	4.93	111.33	106.20
5	B	1000	ORC	C19-C17-N16	4.91	128.40	119.86
2	A	1000	OR8	O17-S14-O16	4.17	127.46	119.82
5	B	1000	ORC	C26-C24-C20	-4.15	128.21	133.91
2	A	1000	OR8	C23-C8-N9	4.07	126.93	119.86

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1000	OR8	C26-N19-N20	3.74	124.89	119.29
2	A	1000	OR8	O16-S14-N13	3.45	110.63	107.41
2	A	1000	OR8	C11-O10-N9	3.39	112.28	108.51
5	B	1000	ORC	C17-C12-S10	-3.38	124.71	127.23
5	B	1000	ORC	C19-C17-C12	-3.35	124.84	129.58
2	A	1000	OR8	C21-N20-N19	3.21	109.12	105.55
2	A	1000	OR8	C6-C1-C2	-3.01	117.58	120.80
5	B	1000	ORC	C25-C21-N22	3.00	125.07	119.86
5	B	1000	ORC	C24-O23-N22	2.92	111.75	108.51
5	B	1000	ORC	C14-O15-N16	2.68	111.49	108.51
2	A	1000	OR8	C3-C2-C7	-2.30	117.16	120.79
5	B	1000	ORC	C9-C7-C5	-2.25	106.94	112.22
5	B	1000	ORC	C24-C20-C21	-2.22	103.09	104.38

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1934	GOL	C1-C2-C3-O3
4	A	1935	GOL	O1-C1-C2-C3
4	B	1933	GOL	O2-C2-C3-O3
4	B	1933	GOL	O1-C1-C2-C3
4	B	1933	GOL	C1-C2-C3-O3
4	A	1934	GOL	O2-C2-C3-O3
4	A	1935	GOL	O1-C1-C2-O2
4	B	1933	GOL	O1-C1-C2-O2
4	A	1936	GOL	O2-C2-C3-O3
2	A	1000	OR8	C12-N13-S14-O17
2	A	1000	OR8	C21-C15-S14-O16
2	A	1000	OR8	C18-C15-S14-O16


There are no ring outliers.


2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1933	SO4	1	0
4	A	1935	GOL	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	250/260 (96%)	0.09	8 (3%)	50 50	7, 20, 41, 61	4 (1%)
1	B	250/260 (96%)	0.69	21 (8%)	17 15	16, 33, 58, 78	2 (0%)
All	All	500/520 (96%)	0.39	29 (5%)	29 27	7, 26, 53, 78	6 (1%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	706	THR	5.6
1	B	864	VAL	5.3
1	B	863	VAL	3.8
1	A	708	PRO	3.7
1	B	708	PRO	3.5
1	A	705[A]	ASN	3.4
1	B	698	VAL	3.3
1	A	683	LEU	3.3
1	B	711	SER	3.2
1	B	701	ALA	3.2
1	A	863	VAL	3.2
1	A	864	VAL	3.2
1	B	683	LEU	3.2
1	B	706	THR	3.1
1	B	871	TYR	2.9
1	B	707	LYS	2.7
1	B	812	GLN	2.6
1	B	860	GLN	2.5
1	A	707	LYS	2.5
1	B	799	LEU	2.5
1	A	861	LYS	2.4
1	B	703	HIS	2.4
1	B	783	ILE	2.3
1	B	702	GLY	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	861	LYS	2.1
1	B	862	GLY	2.1
1	B	795	TYR	2.0
1	B	787	GLN	2.0
1	B	689	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	A	1934	6/6	0.80	0.18	45,47,47,48	0
5	ORC	B	1000	26/26	0.86	0.14	35,39,41,42	0
4	GOL	B	1933	6/6	0.87	0.13	46,46,47,48	0
4	GOL	A	1936	6/6	0.88	0.14	43,43,45,47	0
2	OR8	A	1000	26/26	0.89	0.12	19,23,29,36	0
4	GOL	A	1935	6/6	0.90	0.14	30,32,33,33	0
3	SO4	A	1933	5/5	0.97	0.08	26,26,32,32	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ORC B 1000:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around OR8 A 1000:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.