



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 04:12 AM UTC

PDB ID : 3ZRY / pdb_00003zry
Title : Rotor architecture in the F(1)-c(10)-ring complex of the yeast F-ATP synthase
Authors : Giraud, M.-F.; Dautant, A.
Deposited on : 2011-06-21
Resolution : 6.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 6.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1162 (9.00-4.00)
Clashscore	190562	1020 (8.98-4.02)
Ramachandran outliers	187476	1050 (9.00-4.00)
Sidechain outliers	187428	1014 (9.00-4.00)
RSRZ outliers	180081	1155 (9.00-4.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	510	4% 62% 30% 5%
1	B	510	7% 56% 35% 5% 5%
1	C	510	5% 61% 32% 5%
2	D	478	4% 67% 29% •
2	E	478	7% 63% 33% •

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	478	
3	G	278	
4	H	138	
5	I	61	
6	J	76	
6	K	76	
6	L	76	
6	M	76	
6	N	76	
6	O	76	
6	P	76	
6	Q	76	
6	R	76	
6	S	76	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	ANP	A	600	X	-	-	-
7	ANP	B	600	X	-	-	-
7	ANP	C	600	X	-	-	-
7	ANP	D	600	X	-	-	-
7	ANP	F	600	X	-	X	-
8	MG	A	700	-	-	-	X
8	MG	B	700	-	-	-	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 30100 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP SYNTHASE SUBUNIT ALPHA, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	483	Total	C	N	O	S	0	0	1
			3665	2314	649	699	3			
1	B	484	Total	C	N	O	S	0	0	1
			3670	2317	650	700	3			
1	C	485	Total	C	N	O	S	0	0	1
			3674	2319	651	701	3			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	305	SER	PRO	variant	UNP P07251
B	305	SER	PRO	variant	UNP P07251
C	305	SER	PRO	variant	UNP P07251

- Molecule 2 is a protein called ATP SYNTHASE SUBUNIT BETA, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	471	Total	C	N	O	S	0	0	1
			3550	2250	605	689	6			
2	E	469	Total	C	N	O	S	0	0	1
			3537	2243	603	685	6			
2	F	470	Total	C	N	O	S	0	0	1
			3544	2247	604	687	6			

- Molecule 3 is a protein called ATP SYNTHASE SUBUNIT GAMMA, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	266	Total	C	N	O	S	0	0	1
			2022	1270	355	387	10			

- Molecule 4 is a protein called ATP SYNTHASE SUBUNIT DELTA, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	H	119	Total	C	N	O	S	0	0	0
			747	468	132	145	2			


- Molecule 5 is a protein called ATP SYNTHASE CATALYTIC SECTOR F1 EPSILON SUB-UNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	I	49	Total	C	N	O		0	0	1
			325	201	57	67				

- Molecule 6 is a protein called ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	J	73	Total	C	N	O	S	0	0	1
			515	343	81	88	3			
6	K	73	Total	C	N	O	S	0	0	1
			517	347	80	87	3			
6	L	72	Total	C	N	O	S	0	0	1
			507	337	80	87	3			
6	M	73	Total	C	N	O	S	0	0	1
			515	342	81	88	4			
6	N	73	Total	C	N	O	S	0	0	1
			515	342	81	88	4			
6	O	74	Total	C	N	O	S	0	0	1
			523	348	82	89	4			
6	P	75	Total	C	N	O	S	0	0	1
			534	357	83	90	4			
6	Q	75	Total	C	N	O	S	0	0	1
			534	357	83	90	4			
6	R	74	Total	C	N	O	S	0	0	1
			523	348	82	89	4			
6	S	74	Total	C	N	O	S	0	0	1
			523	348	82	89	4			

- Molecule 7 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (CCD ID: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	A	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
7	B	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
7	C	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
7	D	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
7	F	1	Total	C	N	O	P	0	0
			31	10	6	12	3		


- Molecule 8 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	Mg	0	0
			1	1		
8	B	1	Total	Mg	0	0
			1	1		
8	C	1	Total	Mg	0	0
			1	1		
8	D	1	Total	Mg	0	0
			1	1		
8	F	1	Total	Mg	0	0
			1	1		


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ATP SYNTHASE SUBUNIT ALPHA, MITOCHONDRIAL



• Molecule 1: ATP SYNTHASE SUBUNIT ALPHA, MITOCHONDRIAL






• Molecule 1: ATP SYNTHASE SUBUNIT ALPHA, MITOCHONDRIAL




• Molecule 2: ATP SYNTHASE SUBUNIT BETA, MITOCHONDRIAL







• Molecule 2: ATP SYNTHASE SUBUNIT BETA, MITOCHONDRIAL




• Molecule 2: ATP SYNTHASE SUBUNIT BETA, MITOCHONDRIAL







• Molecule 3: ATP SYNTHASE SUBUNIT GAMMA, MITOCHONDRIAL



• Molecule 4: ATP SYNTHASE SUBUNIT DELTA, MITOCHONDRIAL




• Molecule 5: ATP SYNTHASE CATALYTIC SECTOR F1 EPSILON SUBUNIT




• Molecule 6: ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL







• Molecule 6: ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL




• Molecule 6: ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL



• Molecule 6: ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL




• Molecule 6: ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL




• Molecule 6: ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL




• Molecule 6: ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL




- Molecule 6: ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL



- Molecule 6: ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL



- Molecule 6: ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	107.69Å 174.91Å 164.26Å 90.00° 93.92° 90.00°	Depositor
Resolution (Å)	54.62 – 6.50 54.62 – 6.50	Depositor EDS
% Data completeness (in resolution range)	96.6 (54.62-6.50) 96.5 (54.62-6.50)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.33 (at 6.69Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.317 , 0.339 0.306 , 0.346	Depositor DCC
R_{free} test set	560 reflections (4.72%)	wwPDB-VP
Wilson B-factor (Å ²)	274.6	Xtriage
Anisotropy	0.372	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 235.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.22$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.72	EDS
Total number of atoms	30100	wwPDB-VP
Average B, all atoms (Å ²)	309.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.28	0/3719	0.79	0/5034
1	B	0.29	0/3724	0.80	5/5041 (0.1%)
1	C	0.28	0/3729	0.82	3/5049 (0.1%)
2	D	0.30	0/3606	0.84	12/4891 (0.2%)
2	E	0.28	0/3593	0.77	1/4872 (0.0%)
2	F	0.28	0/3600	0.79	2/4883 (0.0%)
3	G	0.32	0/2045	0.83	4/2752 (0.1%)
4	H	0.30	0/755	0.87	1/1035 (0.1%)
5	I	0.38	0/327	1.00	4/447 (0.9%)
6	J	0.32	0/522	0.84	1/708 (0.1%)
6	K	0.33	0/525	0.99	3/712 (0.4%)
6	L	0.32	0/514	0.91	2/697 (0.3%)
6	M	0.33	0/522	0.91	1/707 (0.1%)
6	N	0.32	0/522	0.89	1/707 (0.1%)
6	O	0.35	0/530	0.94	1/718 (0.1%)
6	P	0.34	0/542	0.90	2/734 (0.3%)
6	Q	0.33	0/542	0.90	2/734 (0.3%)
6	R	0.34	0/530	0.94	3/718 (0.4%)
6	S	0.36	0/530	0.95	3/718 (0.4%)
All	All	0.30	0/30377	0.83	51/41157 (0.1%)

There are no bond length outliers.

The worst 5 of 51 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	K	39	ARG	N-CA-C	-7.65	104.85	114.56
2	D	224	GLU	CA-C-N	7.37	125.04	119.66
2	D	224	GLU	C-N-CA	7.37	125.04	119.66
1	C	289	ARG	CA-C-N	-7.09	113.08	120.38
1	C	289	ARG	C-N-CA	-7.09	113.08	120.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3665	0	3748	126	0
1	B	3670	0	3753	159	0
1	C	3674	0	3754	130	0
2	D	3550	0	3621	122	0
2	E	3537	0	3610	134	0
2	F	3544	0	3616	148	0
3	G	2022	0	2069	214	0
4	H	747	0	592	69	0
5	I	325	0	249	48	0
6	J	515	0	558	64	0
6	K	517	0	559	38	0
6	L	507	0	547	39	0
6	M	515	0	559	33	0
6	N	515	0	559	38	0
6	O	523	0	570	46	0
6	P	534	0	579	35	0
6	Q	534	0	579	43	0
6	R	523	0	570	36	0
6	S	523	0	570	52	0
7	A	31	0	13	1	0
7	B	31	0	13	2	0
7	C	31	0	13	5	0
7	D	31	0	13	6	0
7	F	31	0	13	10	0
8	A	1	0	0	0	0
8	B	1	0	0	0	0
8	C	1	0	0	0	0
8	D	1	0	0	0	0
8	F	1	0	0	0	0
All	All	30100	0	30727	1282	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 1282 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:M:40:ASN:HB3	6:M:41:PRO:HA	1.20	1.15
1:C:291:PRO:HD2	3:G:273:GLY:CA	1.76	1.15
1:C:509:THR:HG23	1:C:510:PHE:N	1.54	1.14
2:E:390:ILE:HG21	3:G:28:SER:CB	1.76	1.14
3:G:191:SER:HB2	3:G:194:PHE:CB	1.77	1.12

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	479/510 (94%)	430 (90%)	46 (10%)	3 (1%)	21	59
1	B	480/510 (94%)	433 (90%)	45 (9%)	2 (0%)	30	67
1	C	483/510 (95%)	445 (92%)	36 (8%)	2 (0%)	30	67
2	D	469/478 (98%)	429 (92%)	37 (8%)	3 (1%)	21	59
2	E	467/478 (98%)	425 (91%)	36 (8%)	6 (1%)	9	42
2	F	468/478 (98%)	419 (90%)	41 (9%)	8 (2%)	7	36
3	G	260/278 (94%)	225 (86%)	28 (11%)	7 (3%)	4	25
4	H	109/138 (79%)	88 (81%)	16 (15%)	5 (5%)	2	16
5	I	43/61 (70%)	27 (63%)	8 (19%)	8 (19%)	0	2
6	J	71/76 (93%)	64 (90%)	6 (8%)	1 (1%)	9	40
6	K	71/76 (93%)	65 (92%)	5 (7%)	1 (1%)	9	40
6	L	70/76 (92%)	65 (93%)	4 (6%)	1 (1%)	9	40
6	M	71/76 (93%)	66 (93%)	3 (4%)	2 (3%)	4	24
6	N	71/76 (93%)	66 (93%)	3 (4%)	2 (3%)	4	24

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
6	O	72/76 (95%)	67 (93%)	4 (6%)	1 (1%)	9	40
6	P	73/76 (96%)	67 (92%)	4 (6%)	2 (3%)	4	25
6	Q	73/76 (96%)	63 (86%)	9 (12%)	1 (1%)	9	40
6	R	72/76 (95%)	66 (92%)	5 (7%)	1 (1%)	9	40
6	S	72/76 (95%)	66 (92%)	4 (6%)	2 (3%)	4	24
All	All	3974/4201 (95%)	3576 (90%)	340 (9%)	58 (2%)	8	40

5 of 58 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	29	GLU
2	F	28	SER
3	G	152	SER
3	G	185	ALA
3	G	186	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	388/412 (94%)	364 (94%)	24 (6%)	16	38
1	B	388/412 (94%)	350 (90%)	38 (10%)	7	24
1	C	388/412 (94%)	353 (91%)	35 (9%)	9	27
2	D	380/384 (99%)	353 (93%)	27 (7%)	13	35
2	E	378/384 (98%)	356 (94%)	22 (6%)	18	40
2	F	379/384 (99%)	353 (93%)	26 (7%)	14	36
3	G	216/236 (92%)	185 (86%)	31 (14%)	3	13
4	H	53/112 (47%)	41 (77%)	12 (23%)	1	6
5	I	23/48 (48%)	19 (83%)	4 (17%)	2	9
6	J	53/56 (95%)	41 (77%)	12 (23%)	1	6
6	K	53/56 (95%)	47 (89%)	6 (11%)	5	19

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
6	L	52/56 (93%)	48 (92%)	4 (8%)	12	32
6	M	53/56 (95%)	47 (89%)	6 (11%)	5	19
6	N	53/56 (95%)	48 (91%)	5 (9%)	8	26
6	O	54/56 (96%)	48 (89%)	6 (11%)	6	20
6	P	55/56 (98%)	50 (91%)	5 (9%)	9	27
6	Q	55/56 (98%)	47 (86%)	8 (14%)	3	13
6	R	54/56 (96%)	44 (82%)	10 (18%)	1	8
6	S	54/56 (96%)	49 (91%)	5 (9%)	8	26
All	All	3129/3344 (94%)	2843 (91%)	286 (9%)	9	27

5 of 286 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
6	K	16	THR
6	L	33	LEU
6	P	59	GLU
2	D	9	ILE
1	C	498	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 36 such sidechains are listed below:

Mol	Chain	Res	Type
3	G	224	GLN
6	S	40	ASN
5	I	15	ASN
6	O	2	GLN
1	B	454	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 5 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	ANP	D	600	8	33,33,33	3.49	10 (30%)	45,52,52	3.03	21 (46%)
7	ANP	C	600	8	33,33,33	3.51	9 (27%)	45,52,52	2.95	21 (46%)
7	ANP	A	600	8	33,33,33	3.48	9 (27%)	45,52,52	3.07	22 (48%)
7	ANP	F	600	8	33,33,33	3.55	8 (24%)	45,52,52	2.97	22 (48%)
7	ANP	B	600	8	33,33,33	3.47	9 (27%)	45,52,52	3.03	20 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	ANP	D	600	8	2/2/7/8	9/18/38/38	0/3/3/3
7	ANP	C	600	8	2/2/7/8	10/18/38/38	0/3/3/3
7	ANP	A	600	8	2/2/7/8	9/18/38/38	0/3/3/3
7	ANP	F	600	8	2/2/7/8	9/18/38/38	0/3/3/3
7	ANP	B	600	8	2/2/7/8	4/18/38/38	0/3/3/3

The worst 5 of 45 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	F	600	ANP	PA-O3A	-13.18	1.45	1.59
7	C	600	ANP	PA-O3A	-12.99	1.45	1.59
7	D	600	ANP	PA-O3A	-12.93	1.45	1.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	600	ANP	PA-O3A	-12.80	1.45	1.59
7	B	600	ANP	PA-O3A	-12.76	1.45	1.59

The worst 5 of 106 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	D	600	ANP	O1B-PB-N3B	-9.49	97.80	111.77
7	A	600	ANP	O1B-PB-N3B	-9.41	97.91	111.77
7	C	600	ANP	O1B-PB-N3B	-8.90	98.67	111.77
7	F	600	ANP	O1B-PB-N3B	-8.62	99.08	111.77
7	B	600	ANP	O1B-PB-N3B	-8.11	99.83	111.77

5 of 10 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
7	A	600	ANP	C3'
7	A	600	ANP	C1'
7	B	600	ANP	C3'
7	B	600	ANP	C1'
7	C	600	ANP	C3'

5 of 41 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	600	ANP	PG-N3B-PB-O1B
7	A	600	ANP	PG-N3B-PB-O3A
7	A	600	ANP	PA-O3A-PB-O1B
7	A	600	ANP	C5'-O5'-PA-O1A
7	B	600	ANP	PB-N3B-PG-O1G


There are no ring outliers.


5 monomers are involved in 24 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	D	600	ANP	6	0
7	C	600	ANP	5	0
7	A	600	ANP	1	0
7	F	600	ANP	10	0
7	B	600	ANP	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In


addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

The following chains have linkage breaks:

Mol	Chain	Number of breaks
3	G	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	G	200:ASP	C	201:THR	N	3.99

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	483/510 (94%)	0.08	19 (3%)	43	42	190, 219, 279, 304	0
1	B	484/510 (94%)	0.14	35 (7%)	21	27	184, 237, 325, 351	0
1	C	485/510 (95%)	0.11	27 (5%)	30	34	255, 279, 425, 448	0
2	D	471/478 (98%)	0.02	21 (4%)	38	38	217, 250, 325, 361	0
2	E	469/478 (98%)	0.21	35 (7%)	20	26	215, 255, 301, 336	0
2	F	470/478 (98%)	0.02	14 (2%)	52	46	242, 287, 363, 404	0
3	G	266/278 (95%)	0.26	16 (6%)	27	32	315, 367, 397, 413	0
4	H	119/138 (86%)	-0.18	4 (3%)	48	44	339, 382, 438, 476	0
5	I	49/61 (80%)	-0.39	0	100	100	298, 368, 423, 442	0
6	J	73/76 (96%)	0.38	8 (10%)	10	18	407, 446, 474, 482	0
6	K	73/76 (96%)	0.94	13 (17%)	4	11	414, 437, 478, 491	0
6	L	72/76 (94%)	0.50	8 (11%)	10	18	399, 482, 524, 524	0
6	M	73/76 (96%)	-0.30	1 (1%)	73	62	424, 488, 520, 520	0
6	N	73/76 (96%)	-0.11	4 (5%)	30	34	430, 526, 526, 526	0
6	O	74/76 (97%)	0.16	4 (5%)	31	34	420, 455, 486, 493	0
6	P	75/76 (98%)	0.61	10 (13%)	7	15	419, 452, 486, 506	0
6	Q	75/76 (98%)	0.52	6 (8%)	18	25	445, 458, 488, 517	0
6	R	74/76 (97%)	0.17	5 (6%)	23	28	414, 436, 466, 485	0
6	S	74/76 (97%)	0.02	3 (4%)	41	41	378, 432, 464, 480	0
All	All	4032/4201 (95%)	0.13	233 (5%)	29	33	184, 280, 466, 526	0

The worst 5 of 233 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	42	PRO	10.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
6	P	67	MET	9.9
6	K	56	ALA	9.7
6	Q	3	LEU	8.7
6	K	53	LEU	8.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
8	MG	A	700	1/1	0.24	0.46	394,394,394,394	0
8	MG	B	700	1/1	0.28	0.42	410,410,410,410	0
8	MG	F	700	1/1	0.37	0.22	376,376,376,376	0
8	MG	C	700	1/1	0.60	0.23	405,405,405,405	0
7	ANP	C	600	31/31	0.69	0.09	309,327,335,341	0
7	ANP	F	600	31/31	0.87	0.10	216,249,272,272	0
8	MG	D	700	1/1	0.88	0.19	395,395,395,395	0
7	ANP	B	600	31/31	0.89	0.09	190,214,226,227	0
7	ANP	A	600	31/31	0.92	0.14	199,210,219,227	0
7	ANP	D	600	31/31	0.92	0.13	231,238,281,287	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ANP C 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ANP F 600:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around ANP B 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ANP A 600:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.