



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 07:29 PM UTC

PDB ID : 7ZR4 / pdb_00007zr4
Title : Molybdenum storage protein loaded with polyoxotungstates in the in vivo-like state
Authors : Ermler, U.; Aziz, I.
Deposited on : 2022-05-03
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	
2	B	269	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	UNL	A	304	-	-	X	-
5	UNL	B	302	-	-	X	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 4403 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Molybdenum storage protein subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	246	Total	C	N	O	S	0	5	0
			1855	1171	351	330	3			


- Molecule 2 is a protein called Molybdenum storage protein subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	268	Total	C	N	O	S	0	6	0
			2002	1267	354	373	8			

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
4	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 5 is UNKNOWN LIGAND (CCD ID: UNL) (formula:) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	3	Total	O	P	W	44	0
			90	70	1	19		
5	B	3	Total	O	W		45	0
			89	69	20			

- Molecule 6 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Cl	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	151	Total	O	0	2
			153	153		
7	B	149	Total	O	0	1
			150	150		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Molybdenum storage protein subunit alpha



- Molecule 2: Molybdenum storage protein subunit beta



4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	115.67Å 115.67Å 234.49Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.49 – 1.70 46.49 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.0 (46.49-1.70) 99.4 (46.49-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.13 (at 1.70Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.218 , 0.240 0.218 , 0.238	Depositor DCC
R_{free} test set	5048 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	29.6	Xtriage
Anisotropy	0.451	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 39.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4403	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, UNL, ATP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/1921	0.51	0/2615
2	B	0.80	4/2063 (0.2%)	0.79	8/2802 (0.3%)
All	All	0.62	4/3984 (0.1%)	0.67	8/5417 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	126	VAL	C-O	-23.53	0.93	1.23
2	B	125	VAL	C-N	-11.87	1.19	1.33
2	B	131	LEU	C-O	-11.86	1.07	1.23
2	B	130	GLY	C-O	-7.64	1.13	1.23

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	131	LEU	CB-CA-C	15.16	134.04	111.76
2	B	132	SER	N-CA-C	-11.72	93.49	109.54
2	B	129	ALA	N-CA-C	6.91	125.52	110.80
2	B	131	LEU	N-CA-C	-6.51	95.34	107.71
2	B	4	SER	N-CA-C	-6.35	104.41	111.71
2	B	130	GLY	CA-C-N	-5.98	113.57	123.37
2	B	130	GLY	C-N-CA	-5.98	113.57	123.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	131	LEU	N-CA-CB	-5.95	95.81	110.32

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	131	LEU	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1855	0	1902	18	0
2	B	2002	0	2052	35	0
3	A	1	0	0	0	0
4	A	31	0	12	1	0
4	B	31	0	12	0	0
5	A	90	0	0	3	0
5	B	89	0	0	5	0
6	A	1	0	0	0	0
7	A	153	0	0	1	0
7	B	150	0	0	4	0
All	All	4403	0	3978	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:126:VAL:CG2	2:B:129:ALA:HB2	1.57	1.33
2:B:126:VAL:CG2	2:B:129:ALA:CB	2.20	1.20
2:B:126:VAL:HG21	2:B:129:ALA:CB	1.76	1.12
2:B:126:VAL:HG23	2:B:129:ALA:HB2	1.12	1.08
2:B:126:VAL:HG23	2:B:129:ALA:CB	1.89	0.93

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:126:VAL:HG21	2:B:129:ALA:HB1	1.53	0.91
2:B:3:ASN:HB2	2:B:6:ALA:H	1.46	0.80
1:A:274:ARG:HD2	1:A:275:PRO:HD2	1.68	0.76
2:B:127:GLY:O	7:B:401:HOH:O	2.11	0.67
5:A:304:UNL:W5	5:A:304:UNL:O25	1.43	0.62
2:B:46:GLN:NE2	7:B:403:HOH:O	2.32	0.62
2:B:104:SER:HB3	5:B:302:UNL:O2	2.01	0.60
2:B:170:ASP:HB2	2:B:227:GLU:CD	2.30	0.56
2:B:35:LEU:HD11	2:B:130:GLY:O	2.05	0.55
5:A:304:UNL:O25	5:A:304:UNL:O14	2.25	0.55
1:A:110:GLY:HA3	5:A:304:UNL:O24	2.08	0.54
1:A:157:HIS:O	2:B:179[B]:GLN:HG2	2.09	0.53
2:B:192:ASP:O	2:B:264:GLY:HA2	2.09	0.52
2:B:244:GLN:NE2	2:B:263:VAL:O	2.36	0.52
1:A:235:VAL:HG12	2:B:157:ARG:HG3	1.91	0.50
2:B:222:HIS:HD2	2:B:231:LEU:HD12	1.76	0.50
1:A:193:ASP:HA	1:A:249:ASN:HB2	1.94	0.50
2:B:3:ASN:HB2	2:B:6:ALA:CB	2.43	0.49
2:B:131:LEU:HB3	5:B:303:UNL:O91	2.13	0.49
1:A:133:VAL:HG11	1:A:177[B]:LEU:HD21	1.95	0.48
2:B:228:PHE:CG	2:B:229:PRO:HD3	2.47	0.48
2:B:73:LEU:HD21	2:B:136:LEU:HD11	1.95	0.48
1:A:227:PRO:HD3	4:A:302:ATP:C6	2.49	0.48
2:B:120:LYS:NZ	7:B:406:HOH:O	2.46	0.48
2:B:104:SER:HA	2:B:149:MET:HG3	1.97	0.46
1:A:132:THR:HG23	1:A:140:HIS:CE1	2.50	0.46
1:A:159:GLU:CD	1:A:167:PRO:HD2	2.41	0.46
2:B:149:MET:HB2	5:B:302:UNL:O9	2.15	0.46
1:A:177[A]:LEU:HG	2:B:155:TRP:NE1	2.31	0.46
2:B:252:GLY:O	2:B:256[B]:ARG:HG3	2.16	0.46
1:A:133:VAL:CG1	1:A:177[B]:LEU:HD21	2.46	0.45
2:B:108:ASP:OD2	5:B:302:UNL:O1	2.34	0.45
2:B:131:LEU:HD23	7:B:402:HOH:O	2.16	0.45
2:B:75:GLY:HA2	2:B:146:PHE:O	2.17	0.44
1:A:213:GLU:HG3	1:A:268:LEU:HB2	2.00	0.44
2:B:147:SER:HB2	5:B:302:UNL:O9	2.17	0.44
1:A:99:GLY:O	1:A:157:HIS:ND1	2.43	0.44
2:B:130:GLY:HA3	2:B:180:PHE:CZ	2.54	0.43
1:A:49:ARG:NE	7:A:408:HOH:O	2.53	0.42
1:A:103:PRO:HD3	1:A:156:HIS:HA	2.01	0.42
2:B:203:LYS:H	2:B:203:LYS:HG3	1.62	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:195:TYR:HB3	2:B:205:ALA:HB1	2.02	0.41
1:A:141:LEU:HD23	1:A:141:LEU:HA	1.82	0.40
1:A:99:GLY:HA3	2:B:179[B]:GLN:CG	2.51	0.40
2:B:101:GLN:HG2	2:B:153:LYS:HB3	2.02	0.40
1:A:88:PHE:CE1	1:A:105:ALA:HB2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	249/275 (90%)	244 (98%)	5 (2%)	0	100	100
2	B	272/269 (101%)	260 (96%)	10 (4%)	2 (1%)	18	7
All	All	521/544 (96%)	504 (97%)	15 (3%)	2 (0%)	30	16

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	127	GLY
2	B	129	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/215 (90%)	192 (100%)	1 (0%)	81	76
2	B	211/205 (103%)	208 (99%)	3 (1%)	59	46
All	All	404/420 (96%)	400 (99%)	4 (1%)	68	58

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	129	GLU
2	B	3	ASN
2	B	132	SER
2	B	203	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	3	ASN
2	B	70	HIS
2	B	222	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic and 6 are unknown - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ATP	A	302	3	32,33,33	0.96	3 (9%)	48,52,52	0.74	0
4	ATP	B	301	-	32,33,33	0.78	1 (3%)	48,52,52	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ATP	A	302	3	-	3/22/38/38	0/3/3/3
4	ATP	B	301	-	-	5/22/38/38	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	302	ATP	PG-O3G	-2.42	1.45	1.54
4	A	302	ATP	PB-O2B	-2.20	1.45	1.55
4	A	302	ATP	PG-O2G	-2.18	1.46	1.54
4	B	301	ATP	PG-O3G	-2.00	1.47	1.54

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	301	ATP	C4'-C5'-O5'-PA
4	B	301	ATP	O4'-C1'-N9-C8
4	B	301	ATP	C3'-C4'-C5'-O5'
4	B	301	ATP	C2'-C1'-N9-C8
4	A	302	ATP	C2'-C1'-N9-C4
4	B	301	ATP	C2'-C1'-N9-C4
4	A	302	ATP	O4'-C1'-N9-C8
4	A	302	ATP	C2'-C1'-N9-C8


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	302	ATP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	125:VAL	C	126:VAL	N	1.19

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	246/275 (89%)	0.96	31 (12%) 8 7	22, 34, 64, 88	5 (2%)
2	B	268/269 (99%)	1.22	45 (16%) 4 3	22, 37, 70, 96	6 (2%)
All	All	514/544 (94%)	1.10	76 (14%) 5 5	22, 35, 70, 96	11 (2%)

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	128	GLY	18.7
2	B	127	GLY	13.7
2	B	131	LEU	13.5
2	B	129	ALA	12.7
1	A	131	PRO	11.4
2	B	130	GLY	10.9
2	B	108	ASP	7.3
1	A	129	GLU	7.1
1	A	276	ALA	7.1
1	A	132	THR	6.7
1	A	130	HIS	6.5
1	A	31	GLY	6.5
1	A	223	GLU	6.5
1	A	152	PHE	6.2
2	B	132	SER	5.7
2	B	133	ALA	5.4
2	B	221	LEU	5.2
1	A	135	ASP	5.0
2	B	3	ASN	4.8
2	B	223	ASP	4.6
2	B	222	HIS	4.6
2	B	151	PRO	4.5
1	A	32	LYS	4.4
2	B	46	GLN	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	126	VAL	4.1
2	B	228	PHE	4.0
2	B	101	GLN	4.0
1	A	208	ALA	3.9
1	A	213	GLU	3.9
2	B	201	THR	3.6
2	B	203	LYS	3.5
1	A	275	PRO	3.4
2	B	202	SER	3.4
1	A	221	LYS	3.3
2	B	83	ARG	3.3
1	A	92	LEU	3.2
1	A	94	LEU	3.2
1	A	274	ARG	3.2
1	A	272	GLY	3.1
2	B	149	MET	3.1
2	B	5	THR	3.1
2	B	153	LYS	3.1
2	B	147	SER	3.1
2	B	270	SER	3.0
2	B	262	HIS	2.9
2	B	251	PRO	2.9
1	A	33	ARG	2.9
2	B	136	LEU	2.8
2	B	119	ALA	2.8
2	B	134	VAL	2.8
2	B	71	LYS	2.7
2	B	218	ALA	2.6
2	B	220	GLY	2.6
1	A	203	PRO	2.5
1	A	156	HIS	2.5
2	B	6	ALA	2.5
1	A	207	GLN	2.4
1	A	219	LEU	2.4
2	B	73	LEU	2.4
2	B	7	GLU	2.4
1	A	222	SER	2.3
2	B	4	SER	2.3
1	A	205	ARG	2.3
2	B	125	VAL	2.3
2	B	219	LYS	2.3
1	A	134	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	104	SER	2.3
1	A	262	GLY	2.2
1	A	128	VAL	2.2
1	A	209	ARG	2.1
2	B	200	LYS	2.1
2	B	87	SER	2.1
2	B	180	PHE	2.1
1	A	95	GLY	2.0
2	B	179[A]	GLN	2.0
1	A	35	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	UNL	B	302	15/-	0.56	0.42	117,172,269,323	0
5	UNL	A	305	33/-	0.59	0.23	113,177,277,295	32
5	UNL	A	303	14/-	0.87	0.21	73,139,144,157	13
5	UNL	A	304	43/-	0.90	0.29	43,73,197,248	24
5	UNL	B	304	23/-	0.92	0.21	58,82,132,209	23
5	UNL	B	303	51/-	0.94	0.28	41,80,279,305	35
4	ATP	B	301	31/31	0.95	0.10	35,47,55,56	17
3	MG	A	301	1/1	0.97	0.06	30,30,30,30	0
4	ATP	A	302	31/31	0.98	0.06	27,33,38,42	0
6	CL	A	306	1/1	0.99	0.14	30,30,30,30	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around UNL A 305:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UNL A 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UNL A 304:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UNL B 304:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around UNL B 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ATP B 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.