



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 07:30 PM EDT

PDB ID : 6ZSN / pdb_00006zsn
Title : Crystal structure of rsGCaMP double mutant Ile80His/Val116Ile in the OFF state (illuminated)
Authors : Janowski, R.; Fuenzalida-Werner, J.P.; Mishra, K.; Stiel, A.C.; Niessing, D.
Deposited on : 2020-07-16
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	418	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3300 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein, Green fluorescent protein, Calmodulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	398	Total	C	N	O	S	0	0	0
			3169	1990	536	629	14			

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	36	MET	-	initiating methionine	UNP P42212
A	37	VAL	-	expression tag	UNP P42212
A	38	ASP	-	expression tag	UNP P42212
A	39	SER	-	expression tag	UNP P42212
A	40	SER	-	expression tag	UNP P42212
A	41	ARG	-	expression tag	UNP P42212
A	42	ARG	-	expression tag	UNP P42212
A	43	LYS	-	expression tag	UNP P42212
A	44	TRP	-	expression tag	UNP P42212
A	45	ASN	-	expression tag	UNP P42212
A	46	LYS	-	expression tag	UNP P42212
A	47	THR	-	expression tag	UNP P42212
A	48	GLY	-	expression tag	UNP P42212
A	49	HIS	-	expression tag	UNP P42212
A	50	ALA	-	expression tag	UNP P42212
A	51	VAL	-	expression tag	UNP P42212
A	52	ARG	-	expression tag	UNP P42212
A	53	ALA	-	expression tag	UNP P42212
A	54	ILE	-	expression tag	UNP P42212
A	55	GLY	-	expression tag	UNP P42212
A	56	ARG	-	expression tag	UNP P42212
A	57	LEU	-	expression tag	UNP P42212
A	58	SER	-	expression tag	UNP P42212
A	59	SER	-	expression tag	UNP P42212
A	60	LEU	-	expression tag	UNP P42212
A	61	GLU	-	expression tag	UNP P42212
A	66	LYS	MET	conflict	UNP P42212

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	76	ALA	VAL	conflict	UNP P42212
A	80	HIS	ILE	conflict	UNP P42212
A	88	GLY	SER	conflict	UNP P42212
A	93	TYR	ASP	conflict	UNP P42212
A	116	ILE	THR	conflict	UNP P42212
A	119	LYS	ALA	conflict	UNP P42212
A	144	LEU	HIS	conflict	UNP P42212
A	152	GLY	-	linker	UNP P42212
A	153	GLY	-	linker	UNP P42212
A	154	THR	-	linker	UNP P42212
A	155	GLY	-	linker	UNP P42212
A	156	GLY	-	linker	UNP P42212
A	157	SER	-	linker	UNP P42212
A	158	MET	-	linker	UNP P42212
A	159	VAL	-	linker	UNP P42212
A	?	-	PHE	deletion	UNP P42212
A	?	-	SER	deletion	UNP P42212
A	222	LEU	TYR	conflict	UNP P42212
A	224	PIA	GLY	conflict	UNP P42212
A	227	LEU	GLN	conflict	UNP P42212
A	251	ILE	VAL	conflict	UNP P42212
A	373	THR	ILE	conflict	UNP K4DIE3
A	381	TYR	ASP	conflict	UNP K4DIE3
A	452	LEU	-	expression tag	UNP K4DIE3
A	453	GLU	-	expression tag	UNP K4DIE3
A	454	HIS	-	expression tag	UNP K4DIE3
A	455	HIS	-	expression tag	UNP K4DIE3

- Molecule 2 is FORMIC ACID (CCD ID: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			3	1	2		
2	A	1	Total	C	O	0	0
			3	1	2		
2	A	1	Total	C	O	0	0
			3	1	2		
2	A	1	Total	C	O	0	0
			3	1	2		
2	A	1	Total	C	O	0	0
			3	1	2		

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Ca	0	0
			4	4		

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Na	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	108	Total 108	O 108	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Green fluorescent protein,Green fluorescent protein,Calmodulin



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	118.97Å 118.97Å 96.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.61 – 2.60 46.61 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.61-2.60) 99.9 (46.61-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.77 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.191 , 0.259 0.193 , 0.256	Depositor DCC
R_{free} test set	1092 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	54.5	Xtriage
Anisotropy	0.019	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 28.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3300	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, PIA, FMT, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.03	0/3205	1.39	1/4315 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	103	ASP	CA-CB-CG	5.61	118.21	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3169	0	3067	26	0
2	A	18	0	6	0	0
3	A	4	0	0	0	0
4	A	1	0	0	0	0
5	A	108	0	0	0	0
All	All	3300	0	3073	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:419:LEU:HD12	1:A:419:LEU:O	1.90	0.72
1:A:408:LEU:O	1:A:412:MET:HG2	1.89	0.71
1:A:389:ARG:HD2	1:A:441:TYR:CE1	2.43	0.54
1:A:410:HIS:O	1:A:414:ASN:HB2	2.07	0.53
1:A:419:LEU:HD13	1:A:424:VAL:CG2	2.37	0.53
1:A:229:PHE:CE2	1:A:277:LEU:HD22	2.44	0.53
1:A:420:THR:C	1:A:422:GLU:H	2.18	0.52
1:A:388:ILE:HG21	1:A:445:VAL:HG22	1.93	0.51
1:A:176:LEU:C	1:A:176:LEU:HD23	2.36	0.50
1:A:412:MET:HB3	1:A:419:LEU:HD21	1.92	0.50
1:A:64:TYR:O	1:A:76:ALA:HA	2.12	0.50
1:A:419:LEU:HD13	1:A:424:VAL:HG22	1.94	0.49
1:A:229:PHE:HE2	1:A:277:LEU:HD22	1.78	0.49
1:A:350:GLU:O	1:A:354:MET:HG3	2.12	0.49
1:A:340:ARG:HA	1:A:344:GLN:O	2.14	0.48
1:A:131:MET:HG3	1:A:301:TYR:OH	2.16	0.45
1:A:191:GLY:HA3	1:A:201:THR:O	2.17	0.45
1:A:367:ASP:HB3	1:A:369:PRO:HD2	1.97	0.45
1:A:47:THR:O	1:A:51:VAL:HG23	2.17	0.44
1:A:64:TYR:OH	1:A:357:GLU:OE1	2.31	0.44
1:A:321:LEU:HD21	1:A:415:LEU:HD22	2.00	0.43
1:A:368:PHE:HB3	1:A:369:PRO:HD3	1.99	0.42
1:A:63:VAL:O	1:A:113:TYR:HA	2.21	0.41
1:A:98:ASN:HA	1:A:251:ILE:O	2.21	0.41
1:A:268:ALA:HA	1:A:280:ARG:O	2.21	0.41
1:A:403:ILE:HB	1:A:439:VAL:HB	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	391/418 (94%)	373 (95%)	18 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	342/358 (96%)	322 (94%)	20 (6%)	18	39

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	ARG
1	A	66	LYS
1	A	79	LYS
1	A	99	THR
1	A	103	ASP
1	A	164	GLU
1	A	173	LEU
1	A	201	THR
1	A	265	LYS
1	A	281	ILE
1	A	321	LEU
1	A	341	SER
1	A	377	ARG
1	A	385	GLU
1	A	390	GLU
1	A	409	ARG
1	A	414	ASN
1	A	422	GLU
1	A	427	MET
1	A	447	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	97	GLN
1	A	111	ASN
1	A	438	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	PIA	A	224	1	20,21,22	3.48	4 (20%)	28,29,31	5.13	15 (53%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PIA	A	224	1	-	3/8/27/28	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	224	PIA	CB2-CA2	14.13	1.48	1.35
1	A	224	PIA	CA2-C2	-3.52	1.44	1.48
1	A	224	PIA	CA2-N2	-3.01	1.32	1.38
1	A	224	PIA	C2-N3	-2.96	1.33	1.40

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	224	PIA	CB2-CA2-C2	16.48	142.34	122.36
1	A	224	PIA	CB2-CA2-N2	-11.73	112.83	128.76
1	A	224	PIA	CA2-C2-N3	6.88	109.28	103.50
1	A	224	PIA	C2-N3-C1	-6.84	104.90	108.07
1	A	224	PIA	O2-C2-CA2	-6.82	126.67	131.02
1	A	224	PIA	CA1-C1-N2	-5.99	116.54	124.04
1	A	224	PIA	C2-CA2-N2	-5.83	104.77	108.95
1	A	224	PIA	CA1-C1-N3	5.54	131.25	124.69
1	A	224	PIA	CA2-N2-C1	4.22	109.10	105.80
1	A	224	PIA	C3-CA3-N3	3.79	121.04	112.43
1	A	224	PIA	C1-CA1-N1	-3.53	98.74	109.13
1	A	224	PIA	CD2-CG2-CD1	3.04	122.16	117.65
1	A	224	PIA	O3-C3-CA3	-2.95	112.20	125.47
1	A	224	PIA	CE2-CD2-CG2	-2.84	117.55	121.22
1	A	224	PIA	CD1-CE1-CZ	-2.09	117.67	119.88

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	224	PIA	C2-CA2-CB2-CG2
1	A	224	PIA	N2-CA2-CB2-CG2
1	A	224	PIA	C3-CA3-N3-C1

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 5 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FMT	A	504	-	2,2,2	0.45	0	1,1,1	0.09	0
2	FMT	A	503	-	2,2,2	0.22	0	1,1,1	0.19	0
2	FMT	A	506	-	2,2,2	0.50	0	1,1,1	0.03	0
2	FMT	A	505	-	2,2,2	0.44	0	1,1,1	0.08	0
2	FMT	A	502	-	2,2,2	0.55	0	1,1,1	0.01	0
2	FMT	A	501	-	2,2,2	0.23	0	1,1,1	0.14	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	397/418 (94%)	-0.18	13 (3%)	49 43	40, 55, 114, 175	3 (0%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	42	ARG	6.4
1	A	450	ALA	4.7
1	A	145	GLY	4.6
1	A	418	LYS	4.0
1	A	419	LEU	3.6
1	A	307	LEU	3.1
1	A	377	ARG	2.7
1	A	43	LYS	2.6
1	A	44	TRP	2.5
1	A	431	ALA	2.5
1	A	46	LYS	2.2
1	A	413	THR	2.2
1	A	161	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PIA	A	224	20/21	0.95	0.08	39,46,61,62	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	FMT	A	505	3/3	0.83	0.18	66,66,76,79	0
2	FMT	A	502	3/3	0.88	0.14	73,73,92,92	0
2	FMT	A	504	3/3	0.94	0.11	56,56,64,66	0
2	FMT	A	501	3/3	0.95	0.11	56,56,66,68	0
2	FMT	A	506	3/3	0.95	0.16	58,58,62,71	0
2	FMT	A	503	3/3	0.96	0.10	63,63,67,69	0
4	NA	A	511	1/1	0.97	0.22	53,53,53,53	0
3	CA	A	508	1/1	0.98	0.03	47,47,47,47	0
3	CA	A	507	1/1	0.98	0.03	81,81,81,81	0
3	CA	A	510	1/1	0.99	0.04	62,62,62,62	0
3	CA	A	509	1/1	1.00	0.03	49,49,49,49	0

6.5 Other polymers [i](#)

There are no such residues in this entry.