



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 07:35 PM UTC

PDB ID : 1ZTM / pdb_00001ztm
Title : Structure of the Uncleaved Paramyxovirus (hPIV3) Fusion Protein
Authors : Yin, H.S.; Paterson, R.G.; Wen, X.; Lamb, R.A.; Jardetzky, T.S.
Deposited on : 2005-05-27
Resolution : 3.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2469 (3.10-3.02)
Clashscore	190562	2569 (3.10-3.02)
Ramachandran outliers	187476	2424 (3.10-3.02)
Sidechain outliers	187428	2423 (3.10-3.02)
RSRZ outliers	180081	2469 (3.10-3.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	490	% 46% 34% 6% 15%
1	B	490	2% 45% 36% 5% 15%
1	C	490	% 46% 36% 5% 13%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9432 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fusion glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	416	Total	C	N	O	S	0	0	0
			3137	1983	522	618	14			
1	B	418	Total	C	N	O	S	0	0	0
			3117	1970	516	617	14			
1	C	424	Total	C	N	O	S	0	0	0
			3149	1986	526	623	14			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	106	SER	ARG	engineered mutation	UNP P06828
B	106	SER	ARG	engineered mutation	UNP P06828
C	106	SER	ARG	engineered mutation	UNP P06828
A	494	GLY	-	cloning artifact	UNP P06828
B	494	GLY	-	cloning artifact	UNP P06828
C	494	GLY	-	cloning artifact	UNP P06828
A	495	GLY	-	cloning artifact	UNP P06828
B	495	GLY	-	cloning artifact	UNP P06828
C	495	GLY	-	cloning artifact	UNP P06828
A	496	PRO	-	cloning artifact	UNP P06828
B	496	PRO	-	cloning artifact	UNP P06828
C	496	PRO	-	cloning artifact	UNP P06828
A	497	LEU	-	cloning artifact	UNP P06828
B	497	LEU	-	cloning artifact	UNP P06828
C	497	LEU	-	cloning artifact	UNP P06828
A	498	VAL	-	cloning artifact	UNP P06828
B	498	VAL	-	cloning artifact	UNP P06828
C	498	VAL	-	cloning artifact	UNP P06828
A	499	PRO	-	cloning artifact	UNP P06828
B	499	PRO	-	cloning artifact	UNP P06828
C	499	PRO	-	cloning artifact	UNP P06828
A	500	ARG	-	cloning artifact	UNP P06828
B	500	ARG	-	cloning artifact	UNP P06828

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	500	ARG	-	cloning artifact	UNP P06828
A	501	GLY	-	cloning artifact	UNP P06828
B	501	GLY	-	cloning artifact	UNP P06828
C	501	GLY	-	cloning artifact	UNP P06828
A	502	SER	-	cloning artifact	UNP P06828
B	502	SER	-	cloning artifact	UNP P06828
C	502	SER	-	cloning artifact	UNP P06828
A	503	HIS	-	expression tag	UNP P06828
B	503	HIS	-	expression tag	UNP P06828
C	503	HIS	-	expression tag	UNP P06828
A	504	HIS	-	expression tag	UNP P06828
B	504	HIS	-	expression tag	UNP P06828
C	504	HIS	-	expression tag	UNP P06828
A	505	HIS	-	expression tag	UNP P06828
B	505	HIS	-	expression tag	UNP P06828
C	505	HIS	-	expression tag	UNP P06828
A	506	HIS	-	expression tag	UNP P06828
B	506	HIS	-	expression tag	UNP P06828
C	506	HIS	-	expression tag	UNP P06828
A	507	HIS	-	expression tag	UNP P06828
B	507	HIS	-	expression tag	UNP P06828
C	507	HIS	-	expression tag	UNP P06828
A	508	HIS	-	expression tag	UNP P06828
B	508	HIS	-	expression tag	UNP P06828
C	508	HIS	-	expression tag	UNP P06828

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	O	0	0
			1	1		


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Fusion glycoprotein



• Molecule 1: Fusion glycoprotein





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	105.56Å 122.17Å 195.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.02 – 3.05 30.02 – 3.05	Depositor EDS
% Data completeness (in resolution range)	99.5 (30.02-3.05) 99.4 (30.02-3.05)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.40 (at 3.06Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.217 , 0.264 0.213 , 0.258	Depositor DCC
R_{free} test set	2365 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	67.5	Xtriage
Anisotropy	0.772	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 64.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9432	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/3183	1.10	22/4337 (0.5%)
1	B	0.51	0/3163	1.03	19/4318 (0.4%)
1	C	0.49	0/3193	1.05	13/4356 (0.3%)
All	All	0.50	0/9539	1.06	54/13011 (0.4%)

There are no bond length outliers.

The worst 5 of 54 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	24	LYS	N-CA-C	-14.51	95.59	114.31
1	C	250	LYS	N-CA-C	-10.67	99.73	111.36
1	A	250	LYS	N-CA-C	-10.11	100.22	111.14
1	A	33	ASN	N-CA-C	9.10	117.48	108.75
1	B	250	LYS	N-CA-C	-8.41	102.11	111.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3137	0	3062	199	0
1	B	3117	0	3006	222	0
1	C	3149	0	3034	218	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	14	0	13	2	0
2	B	14	0	13	0	0
3	B	1	0	0	0	0
All	All	9432	0	9128	554	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 30.

The worst 5 of 554 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:413:THR:HG22	1:C:415:LYS:H	1.13	1.11
1:C:313:PRO:HD2	1:C:352:MET:HE1	1.35	1.03
1:C:259:THR:HG23	1:C:332:ILE:HG21	1.46	0.96
1:C:264:VAL:HG12	1:C:280:VAL:HA	1.49	0.94
1:B:168:LEU:HD21	1:C:169:ILE:HG22	1.50	0.94

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	412/490 (84%)	364 (88%)	41 (10%)	7 (2%)	7	25
1	B	414/490 (84%)	371 (90%)	36 (9%)	7 (2%)	7	25
1	C	420/490 (86%)	367 (87%)	46 (11%)	7 (2%)	7	25
All	All	1246/1470 (85%)	1102 (88%)	123 (10%)	21 (2%)	7	25

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	447	ASN
1	A	448	SER
1	B	359	ASN
1	B	380	ASN
1	B	447	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	346/433 (80%)	321 (93%)	25 (7%)	13	37
1	B	340/433 (78%)	319 (94%)	21 (6%)	16	42
1	C	339/433 (78%)	305 (90%)	34 (10%)	7	25
All	All	1025/1299 (79%)	945 (92%)	80 (8%)	11	34

5 of 80 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	170	VAL
1	C	366	THR
1	C	187	ILE
1	C	253	ILE
1	C	393	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 37 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	198	GLN
1	C	403	GLN
1	C	206	HIS
1	C	272	ASN
1	B	61	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1359	1	14,14,15	1.28	1 (7%)	17,19,21	0.90	0
2	NAG	A	1359	1	14,14,15	0.78	0	17,19,21	1.08	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1359	1	-	3/6/23/26	0/1/1/1
2	NAG	A	1359	1	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1359	NAG	C1-C2	3.93	1.57	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1359	NAG	C2-N2-C7	-2.31	119.81	122.90
2	A	1359	NAG	C4-C3-C2	-2.18	107.82	111.02

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1359	NAG	C8-C7-N2-C2
2	A	1359	NAG	O7-C7-N2-C2
2	B	1359	NAG	C8-C7-N2-C2
2	B	1359	NAG	O7-C7-N2-C2
2	B	1359	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1359	NAG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	416/490 (84%)	-0.35	6 (1%) 73 51	26, 53, 89, 123	0
1	B	418/490 (85%)	-0.33	8 (1%) 66 43	23, 52, 87, 116	0
1	C	424/490 (86%)	-0.23	5 (1%) 76 55	24, 56, 91, 117	0
All	All	1258/1470 (85%)	-0.30	19 (1%) 72 50	23, 54, 90, 123	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	449	VAL	3.9
1	B	24	LYS	3.7
1	A	484	ILE	3.3
1	A	24	LYS	2.9
1	B	446	ASN	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	B	1359	14/15	0.44	0.15	49,99,105,105	0
2	NAG	A	1359	14/15	0.76	0.13	65,84,97,98	0

6.5 Other polymers [i](#)

There are no such residues in this entry.