



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 01:53 AM UTC

PDB ID : 4ZT3 / pdb_00004zt3
Title : Trypanosoma brucei methionyl-tRNA synthetase in complex with inhibitor N-(3,5-dichlorobenzyl)-N'-(5-fluoro-1H-imidazo[4,5-b]pyridin-2-yl)propane-1,3-diamine (Chem 1614)
Authors : Koh, C.-Y.; Hol, W.G.J.
Deposited on : 2015-05-14
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	542	4% 88% 8% .
1	B	542	8% 89% 8% .

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 8654 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methionyl-tRNA synthetase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	518	Total	As	C	N	O	S	0	0	0
			4117	1	2654	692	759	11			
1	B	526	Total	As	C	N	O	S	0	2	0
			4176	1	2688	708	768	11			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP Q38C91
A	-3	PRO	-	expression tag	UNP Q38C91
A	-2	GLY	-	expression tag	UNP Q38C91
A	-1	SER	-	expression tag	UNP Q38C91
A	0	MET	-	expression tag	UNP Q38C91
A	309	THR	ALA	conflict	UNP Q38C91
A	452	ALA	LYS	engineered mutation	UNP Q38C91
A	453	ARG	LYS	engineered mutation	UNP Q38C91
A	454	ALA	GLU	engineered mutation	UNP Q38C91
A	499	VAL	ALA	conflict	UNP Q38C91
A	503	ASN	SER	conflict	UNP Q38C91
B	-4	GLY	-	expression tag	UNP Q38C91
B	-3	PRO	-	expression tag	UNP Q38C91
B	-2	GLY	-	expression tag	UNP Q38C91
B	-1	SER	-	expression tag	UNP Q38C91
B	0	MET	-	expression tag	UNP Q38C91
B	309	THR	ALA	conflict	UNP Q38C91
B	452	ALA	LYS	engineered mutation	UNP Q38C91
B	453	ARG	LYS	engineered mutation	UNP Q38C91
B	454	ALA	GLU	engineered mutation	UNP Q38C91
B	499	VAL	ALA	conflict	UNP Q38C91
B	503	ASN	SER	conflict	UNP Q38C91

- Molecule 2 is METHIONINE (CCD ID: MET) (formula: C₅H₁₁NO₂S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			9	5	1	2	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: $\text{C}_2\text{H}_6\text{OS}$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	B	1	Total	C	O	S	0	0
			4	2	1	1		
4	B	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 5 is SULFATE ION (CCD ID: SO_4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is N-(3,5-dichlorobenzyl)-N'-(5-fluoro-1H-imidazo[4,5-b]pyridin-2-yl)propane-1,3-diamine (CCD ID: 4RQ) (formula: C₁₆H₁₆Cl₂FN₅).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	B	1	Total	C	Cl	F	N	0	0
			24	16	2	1	5		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	142	Total 142	O 142	0	0
7	B	135	Total 135	O 135	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Methionyl-tRNA synthetase



• Molecule 1: Methionyl-tRNA synthetase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	86.70Å 106.05Å 207.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.64 – 2.80 38.64 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (38.64-2.80) 99.9 (38.64-2.80)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.03 (at 2.81Å)	Xtriage
Refinement program	REFMAC refmac_5.8.0073	Depositor
R, R_{free}	0.209 , 0.249 (Not available) , 0.207	Depositor DCC
R_{free} test set	2414 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	42.0	Xtriage
Anisotropy	0.609	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 47.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8654	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, 4RQ, DMS, GOL, CAS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	0/4215	0.76	2/5740 (0.0%)
1	B	0.54	0/4282	0.77	2/5828 (0.0%)
All	All	0.54	0/8497	0.77	4/11568 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	586	GLY	N-CA-C	-5.73	104.46	112.13
1	B	290	GLY	N-CA-C	5.61	117.74	112.08
1	A	285	GLY	N-CA-C	5.39	117.86	110.69
1	B	766	SER	N-CA-C	5.36	117.20	110.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4117	0	4018	21	0
1	B	4176	0	4072	26	0
2	A	9	0	8	0	0
3	A	6	0	8	0	0
3	B	12	0	16	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	20	0	30	3	0
4	B	8	0	12	3	0
5	B	5	0	0	0	0
6	B	24	0	0	0	0
7	A	142	0	0	0	0
7	B	135	0	0	6	0
All	All	8654	0	8164	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:284:THR:HG22	1:B:326:ILE:HG21	1.74	0.69
1:A:284:THR:HG22	1:A:326:ILE:HG21	1.77	0.67
1:A:445:PRO:HG2	4:A:805:DMS:H21	1.76	0.67
1:B:488:ARG:NH2	1:B:504:GLU:OE2	2.27	0.67
1:B:720:LYS:CE	4:B:804:DMS:H11	2.27	0.64
1:A:579:TYR:OH	1:A:605:GLU:OE2	2.16	0.63
1:B:720:LYS:HE2	4:B:804:DMS:H11	1.80	0.61
1:B:368:GLU:OE1	7:B:901:HOH:O	2.18	0.56
1:B:521:LYS:HE3	7:B:1022:HOH:O	2.04	0.56
1:A:585:SER:HB3	1:A:593:TYR:OH	2.06	0.55
1:A:451:ARG:NH2	1:A:455:THR:HG21	2.22	0.54
1:A:370:PHE:CZ	1:A:448:SER:HB3	2.42	0.54
1:A:658:ILE:N	1:A:659:PRO:HA	2.24	0.53
1:A:577:LEU:HD13	1:A:598:MET:HE1	1.89	0.53
1:B:668:VAL:HG11	1:B:713:LEU:HG	1.93	0.51
4:A:803:DMS:H21	4:A:804:DMS:H23	1.93	0.50
1:B:658:ILE:N	1:B:659:PRO:HA	2.25	0.50
1:A:271:ARG:HH11	4:A:807:DMS:H23	1.76	0.50
1:B:451:ARG:HB2	1:B:456:LEU:HD11	1.92	0.50
1:B:720:LYS:HE3	4:B:804:DMS:H11	1.93	0.50
1:A:668:VAL:HG11	1:A:713:LEU:HG	1.93	0.49
1:B:261:TYR:CZ	1:B:265:ILE:HD11	2.48	0.48
1:A:519:ILE:HD12	1:A:523:HIS:HE1	1.79	0.48
1:A:563:PHE:CE1	1:A:568:LYS:HG3	2.49	0.48
1:B:250:TYR:CZ	1:B:293:VAL:HG23	2.49	0.47
1:A:519:ILE:HD12	1:A:523:HIS:CE1	2.50	0.46
1:B:519:ILE:HD12	1:B:523:HIS:CE1	2.51	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:602:LEU:HD21	1:A:765:ARG:NH1	2.31	0.46
1:B:519:ILE:HD12	1:B:523:HIS:HE1	1.80	0.46
1:B:602:LEU:HD21	1:B:765:ARG:NH1	2.31	0.46
1:B:408:ARG:NH2	7:B:906:HOH:O	2.43	0.45
1:A:261:TYR:CZ	1:A:265:ILE:HD11	2.51	0.45
1:B:714:SER:N	1:B:715:PRO:CD	2.80	0.45
1:A:618:THR:HG22	1:A:752:LEU:HD13	1.99	0.44
1:B:690:THR:HG21	7:B:1028:HOH:O	2.17	0.43
1:B:375:ASN:C	1:B:376:ILE:HD12	2.43	0.43
1:B:687:LEU:HD21	7:B:943:HOH:O	2.18	0.43
1:B:508:PHE:CG	1:B:509:PRO:HA	2.54	0.43
1:B:704:GLU:HA	1:B:704:GLU:OE1	2.19	0.43
1:A:605:GLU:O	1:A:609:THR:HB	2.19	0.43
1:A:488:ARG:NH2	1:A:504:GLU:OE2	2.52	0.42
1:A:704:GLU:HA	1:A:704:GLU:OE1	2.18	0.42
1:A:714:SER:N	1:A:715:PRO:CD	2.83	0.42
1:B:605:GLU:O	1:B:609:THR:HB	2.20	0.41
1:B:487:LEU:HD22	1:B:495:GLU:HG3	2.02	0.41
1:B:556:SER:HB3	1:B:559:LEU:CB	2.51	0.41
1:A:260:VAL:HG13	1:A:324:TYR:OH	2.21	0.40
1:B:434:ARG:HD3	7:B:936:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	511/542 (94%)	500 (98%)	11 (2%)	0	100	100
1	B	523/542 (96%)	510 (98%)	13 (2%)	0	100	100
All	All	1034/1084 (95%)	1010 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	438/468 (94%)	434 (99%)	4 (1%)	70	89
1	B	441/468 (94%)	437 (99%)	4 (1%)	70	89
All	All	879/936 (94%)	871 (99%)	8 (1%)	70	89

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	292	LYS
1	A	506	GLU
1	A	612	ASN
1	A	690	THR
1	B	588	SER
1	B	612	ASN
1	B	688	VAL
1	B	690	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	643	GLN
1	A	662	GLN
1	B	597	ASN
1	B	643	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CAS	A	470	1	5,8,9	1.04	0	1,9,11	2.03	1 (100%)
1	CAS	B	470	1	5,8,9	1.01	0	1,9,11	1.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	A	470	1	-	0/0/7/9	-
1	CAS	B	470	1	-	0/0/7/9	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	470	CAS	CB-CA-C	-2.03	105.28	110.80

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MET	A	801	-	7,8,8	0.94	1 (14%)	5,9,9	1.67	1 (20%)
4	DMS	A	803	-	3,3,3	0.52	0	3,3,3	0.60	0
3	GOL	B	802	-	5,5,5	0.27	0	5,5,5	0.22	0
4	DMS	A	805	-	3,3,3	0.49	0	3,3,3	0.67	0
4	DMS	A	804	-	3,3,3	0.54	0	3,3,3	0.59	0
4	DMS	A	806	-	3,3,3	0.53	0	3,3,3	0.51	0
4	DMS	B	803	-	3,3,3	0.49	0	3,3,3	0.89	0
3	GOL	A	802	-	5,5,5	0.34	0	5,5,5	0.11	0
3	GOL	B	801	-	5,5,5	0.27	0	5,5,5	0.80	0
5	SO4	B	805	-	4,4,4	0.48	0	6,6,6	0.17	0
6	4RQ	B	806	-	26,26,26	1.67	2 (7%)	32,35,35	0.94	3 (9%)
4	DMS	B	804	-	3,3,3	0.58	0	3,3,3	0.66	0
4	DMS	A	807	-	3,3,3	0.64	0	3,3,3	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MET	A	801	-	-	0/8/8/8	-
3	GOL	B	802	-	-	2/4/4/4	-
3	GOL	A	802	-	-	2/4/4/4	-
3	GOL	B	801	-	-	2/4/4/4	-
6	4RQ	B	806	-	-	1/9/9/9	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	806	4RQ	C-N4	7.98	1.42	1.31

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	MET	OXT-C	-2.24	1.23	1.30
6	B	806	4RQ	C4-N3	2.02	1.35	1.32

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	MET	OXT-C-O	-3.62	115.87	124.08
6	B	806	4RQ	C3-N-C4	-2.67	103.25	107.60
6	B	806	4RQ	C15-C3-N	2.62	109.98	105.47
6	B	806	4RQ	C3-C15-N3	-2.43	105.63	109.96

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	801	GOL	C1-C2-C3-O3
3	B	802	GOL	C1-C2-C3-O3
3	A	802	GOL	C1-C2-C3-O3
3	B	801	GOL	O2-C2-C3-O3
3	B	802	GOL	O2-C2-C3-O3
6	B	806	4RQ	C5-C6-C7-N2
3	A	802	GOL	O2-C2-C3-O3


There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	803	DMS	1	0
4	A	805	DMS	1	0
4	A	804	DMS	1	0
4	B	804	DMS	3	0
4	A	807	DMS	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	517/542 (95%)	-0.11	22 (4%)	40 31	22, 40, 82, 150	0
1	B	525/542 (96%)	0.17	46 (8%)	15 11	26, 48, 101, 121	2 (0%)
All	All	1042/1084 (96%)	0.03	68 (6%)	25 18	22, 44, 94, 150	2 (0%)

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	548	TRP	6.4
1	B	377	THR	5.9
1	A	756	VAL	5.5
1	A	547	TRP	5.3
1	A	550	LYS	5.1
1	B	757	GLU	4.8
1	A	549	THR	4.7
1	A	759	GLU	4.6
1	B	767	THR	4.0
1	B	360	GLU	3.9
1	B	398	THR	3.7
1	B	547	TRP	3.6
1	A	767	THR	3.5
1	B	376	ILE	3.5
1	B	391	LEU	3.4
1	A	766	SER	3.3
1	B	373	PRO	3.2
1	B	365	ILE	3.1
1	B	352	GLY	3.1
1	A	561	ASN	3.0
1	A	238	VAL	3.0
1	B	388	LYS	2.9
1	B	421[A]	HIS	2.9
1	B	372	THR	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	616[A]	ARG	2.8
1	B	396	VAL	2.8
1	B	370	PHE	2.8
1	A	590	ASP	2.8
1	A	491	GLU	2.7
1	A	380	VAL	2.6
1	B	371	LEU	2.6
1	B	366	SER	2.6
1	B	397	VAL	2.6
1	B	369	SER	2.5
1	B	394	GLY	2.5
1	A	589	ASP	2.5
1	B	765	ARG	2.5
1	A	760	VAL	2.5
1	A	685	TRP	2.5
1	B	491	GLU	2.4
1	B	305	TYR	2.4
1	B	353	ASP	2.4
1	B	374	GLN	2.4
1	B	556	SER	2.4
1	B	760	VAL	2.3
1	A	562	VAL	2.3
1	B	357	GLY	2.3
1	B	354	ILE	2.3
1	B	756	VAL	2.2
1	A	755	ALA	2.2
1	B	452	ALA	2.2
1	B	553	LYS	2.2
1	B	557	LYS	2.2
1	B	454	ALA	2.2
1	B	401	SER	2.2
1	A	382	LYS	2.1
1	B	555	ILE	2.1
1	B	395	HIS	2.1
1	B	400	VAL	2.1
1	B	356	LEU	2.1
1	B	399	TRP	2.1
1	A	302	VAL	2.1
1	B	358	ARG	2.0
1	A	381	ASP	2.0
1	A	588	SER	2.0
1	B	393	SER	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	488	ARG	2.0
1	B	389	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CAS	B	470	9/10	0.91	0.18	57,61,77,78	3
1	CAS	A	470	9/10	0.95	0.13	40,44,64,64	3

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	DMS	B	804	4/4	0.80	0.24	81,86,87,90	0
4	DMS	A	805	4/4	0.84	0.25	71,74,75,76	0
4	DMS	A	803	4/4	0.87	0.27	83,89,90,91	0
3	GOL	B	801	6/6	0.89	0.12	39,42,44,46	0
3	GOL	A	802	6/6	0.91	0.12	42,48,51,51	0
4	DMS	A	804	4/4	0.91	0.18	75,76,77,77	0
4	DMS	A	806	4/4	0.92	0.19	70,71,72,73	0
5	SO4	B	805	5/5	0.92	0.23	68,69,73,76	0
3	GOL	B	802	6/6	0.93	0.12	59,60,60,60	0
6	4RQ	B	806	24/24	0.93	0.12	50,60,65,66	0
4	DMS	A	807	4/4	0.94	0.14	52,53,55,61	0
4	DMS	B	803	4/4	0.96	0.14	56,60,61,63	0
2	MET	A	801	9/9	0.98	0.07	25,26,27,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.