



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 10:01 PM UTC

PDB ID : 5ZT0 / pdb_00005zt0
Title : Crystal Structure of Protein Phosphate 1 Complexed with PP1 binding domain of GL
Authors : Yu, J.; Xiang, S.
Deposited on : 2018-05-01
Resolution : 3.32 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1153 (3.34-3.30)
Clashscore	190562	1193 (3.34-3.30)
Ramachandran outliers	187476	1172 (3.34-3.30)
Sidechain outliers	187428	1171 (3.34-3.30)
RSRZ outliers	180081	1153 (3.34-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	294	90% 10%
1	B	294	85% 14%
1	C	294	90% 10%
1	D	294	86% 14%
1	E	294	89% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	294	% 87% 12%
2	G	75	19% 7% 75%
2	H	75	17% 8% 75%
2	I	75	17% 8% 75%
2	J	75	20% 5% 75%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 14774 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein phosphatase PP1-alpha catalytic subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	293	Total	C	N	O	S	0	0	0
			2358	1510	397	433	18			
1	B	293	Total	C	N	O	S	0	0	0
			2358	1510	397	433	18			
1	C	293	Total	C	N	O	S	0	0	0
			2358	1510	397	433	18			
1	D	293	Total	C	N	O	S	0	0	0
			2358	1510	397	433	18			
1	E	293	Total	C	N	O	S	0	0	0
			2358	1510	397	433	18			
1	F	293	Total	C	N	O	S	0	0	0
			2358	1510	397	433	18			

- Molecule 2 is a protein called Protein phosphatase 1 regulatory subunit 3B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	G	19	Total	C	N	O	S	0	0	0
			146	93	27	25	1			
2	H	19	Total	C	N	O	S	0	0	0
			146	93	27	25	1			
2	I	19	Total	C	N	O	S	0	0	0
			146	93	27	25	1			
2	J	19	Total	C	N	O	S	0	0	0
			146	93	27	25	1			

- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Mn	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Mn	0	0
			2	2		
3	C	2	Total	Mn	0	0
			2	2		
3	D	2	Total	Mn	0	0
			2	2		
3	E	2	Total	Mn	0	0
			2	2		
3	F	2	Total	Mn	0	0
			2	2		

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		
4	C	1	Total	O	P	0	0
			5	4	1		
4	D	1	Total	O	P	0	0
			5	4	1		
4	E	1	Total	O	P	0	0
			5	4	1		
4	F	1	Total	O	P	0	0
			5	4	1		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serine/threonine-protein phosphatase PP1-alpha catalytic subunit

Chain A: 



- Molecule 1: Serine/threonine-protein phosphatase PP1-alpha catalytic subunit

Chain B: 




- Molecule 1: Serine/threonine-protein phosphatase PP1-alpha catalytic subunit

Chain C: 



- Molecule 1: Serine/threonine-protein phosphatase PP1-alpha catalytic subunit

Chain D: 




- Molecule 1: Serine/threonine-protein phosphatase PP1-alpha catalytic subunit

Chain E:  89% 11%




- Molecule 1: Serine/threonine-protein phosphatase PP1-alpha catalytic subunit

Chain F:  87% 12%




- Molecule 2: Protein phosphatase 1 regulatory subunit 3B

Chain G:  19% 7% 75%




- Molecule 2: Protein phosphatase 1 regulatory subunit 3B

Chain H:  17% 8% 75%



- Molecule 2: Protein phosphatase 1 regulatory subunit 3B

Chain I:  17% 8% 75%



- Molecule 2: Protein phosphatase 1 regulatory subunit 3B

Chain J:  20% 5% 75%

SER	LYS	PRO	LEU	LEU	ARG	PRO	CYS	ILE	GLN	LEU	SER	SER	LYS	ASN	GLU	ALA	SER	GLY	MET	VAL	ALA	PRO	ALA	VAL	GLN	GLU	LYS	LYS	V59	D67	L73	I74	K77	VAL	PHE	SER	GLU	PHE	ASP	ASP	PRO	LEU	ASP	MET	PRO	PHE	ASN	ILE	THR	GLU	LEU	LEU	ASP	ASN	ILE	VAL	SER
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

LEU	THR	THR	ALA
-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	106.53Å 106.53Å 187.51Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	41.79 – 3.32 41.79 – 3.32	Depositor EDS
% Data completeness (in resolution range)	99.8 (41.79-3.32) 99.8 (41.79-3.32)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 3.32Å)	Xtriage
Refinement program	REFMAC 5.8.0222, PHENIX 1.10.1_2155	Depositor
R, R_{free}	0.200 , 0.233 0.206 , 0.235	Depositor DCC
R_{free} test set	1748 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	97.4	Xtriage
Anisotropy	0.332	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 51.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.000 for -h,-k,l 0.013 for h,-h-k,-l 0.207 for -k,-h,-l	Xtriage
Reported twinning fraction	0.410 for H, K, L 0.103 for K, H, -L 0.097 for -h,-k,l 0.390 for -K, -H, -L	Depositor
Outliers	0 of 35074 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	14774	wwPDB-VP
Average B, all atoms (Å ²)	106.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MN, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.13	0/2412	0.37	0/3259
1	B	0.13	0/2412	0.36	0/3259
1	C	0.12	0/2412	0.36	0/3259
1	D	0.13	0/2412	0.36	0/3259
1	E	0.13	0/2412	0.36	0/3259
1	F	0.14	0/2412	0.36	0/3259
2	G	0.36	0/146	0.49	0/193
2	H	0.27	0/146	0.45	0/193
2	I	0.23	0/146	0.61	0/193
2	J	0.18	0/146	0.37	0/193
All	All	0.14	0/15056	0.37	0/20326

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2358	0	2318	17	0
1	B	2358	0	2318	25	0
1	C	2358	0	2318	17	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	2358	0	2318	24	0
1	E	2358	0	2318	17	0
1	F	2358	0	2318	20	0
2	G	146	0	161	4	0
2	H	146	0	161	4	0
2	I	146	0	161	5	0
2	J	146	0	161	2	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
3	E	2	0	0	0	0
3	F	2	0	0	0	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
4	C	5	0	0	0	0
4	D	5	0	0	0	0
4	E	5	0	0	0	0
4	F	5	0	0	1	0
All	All	14774	0	14552	121	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 121 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:294:GLN:HA	2:H:75:MET:HB2	1.60	0.82
1:A:133:ILE:HG23	1:E:133:ILE:HG23	1.64	0.79
1:C:133:ILE:HG23	1:F:133:ILE:HG23	1.65	0.77
1:B:78:TYR:CE1	2:I:77:LYS:HD2	2.23	0.74
1:B:133:ILE:HG23	1:D:133:ILE:HG23	1.71	0.72

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	291/294 (99%)	272 (94%)	19 (6%)	0	100	100
1	B	291/294 (99%)	273 (94%)	18 (6%)	0	100	100
1	C	291/294 (99%)	269 (92%)	22 (8%)	0	100	100
1	D	291/294 (99%)	270 (93%)	21 (7%)	0	100	100
1	E	291/294 (99%)	270 (93%)	21 (7%)	0	100	100
1	F	291/294 (99%)	273 (94%)	18 (6%)	0	100	100
2	G	17/75 (23%)	17 (100%)	0	0	100	100
2	H	17/75 (23%)	16 (94%)	1 (6%)	0	100	100
2	I	17/75 (23%)	16 (94%)	1 (6%)	0	100	100
2	J	17/75 (23%)	16 (94%)	1 (6%)	0	100	100
All	All	1814/2064 (88%)	1692 (93%)	122 (7%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	258/259 (100%)	255 (99%)	3 (1%)	63	74
1	B	258/259 (100%)	256 (99%)	2 (1%)	73	78
1	C	258/259 (100%)	256 (99%)	2 (1%)	73	78
1	D	258/259 (100%)	257 (100%)	1 (0%)	84	83

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	258/259 (100%)	256 (99%)	2 (1%)	73	78
1	F	258/259 (100%)	256 (99%)	2 (1%)	73	78
2	G	16/67 (24%)	16 (100%)	0	100	100
2	H	16/67 (24%)	16 (100%)	0	100	100
2	I	16/67 (24%)	15 (94%)	1 (6%)	16	44
2	J	16/67 (24%)	14 (88%)	2 (12%)	4	19
All	All	1612/1822 (88%)	1597 (99%)	15 (1%)	70	77

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	246	ARG
2	J	59	VAL
1	E	246	ARG
2	J	67	ASP
1	F	296	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
2	G	68	ASN
2	G	69	GLN
1	D	99	GLN
1	D	214	GLN
1	D	239	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 18 ligands modelled in this entry, 12 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PO4	F	403	3	4,4,4	0.93	0	6,6,6	0.50	0
4	PO4	B	403	3	4,4,4	0.96	0	6,6,6	0.49	0
4	PO4	C	403	3	4,4,4	0.87	0	6,6,6	0.50	0
4	PO4	D	403	3	4,4,4	0.87	0	6,6,6	0.47	0
4	PO4	E	403	3	4,4,4	0.96	0	6,6,6	0.46	0
4	PO4	A	403	3	4,4,4	0.94	0	6,6,6	0.41	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	F	403	PO4	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	293/294 (99%)	-0.48	0 100 100	69, 96, 124, 155	0
1	B	293/294 (99%)	-0.30	1 (0%) 90 82	69, 107, 142, 204	0
1	C	293/294 (99%)	-0.51	1 (0%) 90 82	65, 92, 124, 143	0
1	D	293/294 (99%)	-0.35	2 (0%) 84 70	72, 104, 132, 153	0
1	E	293/294 (99%)	-0.45	0 100 100	74, 111, 155, 194	0
1	F	293/294 (99%)	-0.36	3 (1%) 79 63	76, 116, 163, 214	0
2	G	19/75 (25%)	-0.68	0 100 100	90, 113, 135, 139	0
2	H	19/75 (25%)	-0.84	0 100 100	86, 108, 140, 141	0
2	I	19/75 (25%)	-0.48	0 100 100	104, 120, 143, 151	0
2	J	19/75 (25%)	-0.72	0 100 100	102, 114, 128, 132	0
All	All	1834/2064 (88%)	-0.42	7 (0%) 88 79	65, 104, 146, 214	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	176	LEU	3.9
1	D	175	GLY	3.4
1	D	171	CYS	2.9
1	F	163	ALA	2.6
1	B	133	ILE	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MN	B	402	1/1	0.92	0.07	54,54,54,54	1
4	PO4	B	403	5/5	0.95	0.07	18,61,71,83	5
4	PO4	A	403	5/5	0.96	0.06	29,34,58,66	5
3	MN	F	402	1/1	0.96	0.06	49,49,49,49	1
4	PO4	C	403	5/5	0.96	0.07	18,64,77,111	5
3	MN	B	401	1/1	0.97	0.15	143,143,143,143	0
3	MN	E	402	1/1	0.97	0.05	44,44,44,44	1
4	PO4	D	403	5/5	0.97	0.05	25,49,65,102	5
4	PO4	E	403	5/5	0.97	0.05	21,65,71,113	5
3	MN	D	402	1/1	0.98	0.05	56,56,56,56	1
4	PO4	F	403	5/5	0.98	0.05	20,41,82,92	5
3	MN	C	402	1/1	0.99	0.04	21,21,21,21	1
3	MN	D	401	1/1	1.00	0.03	47,47,47,47	0
3	MN	A	402	1/1	1.00	0.02	31,31,31,31	1
3	MN	E	401	1/1	1.00	0.04	68,68,68,68	0
3	MN	C	401	1/1	1.00	0.06	65,65,65,65	0
3	MN	F	401	1/1	1.00	0.04	55,55,55,55	0
3	MN	A	401	1/1	1.00	0.02	47,47,47,47	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around MN B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PO4 B 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PO4 A 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around MN F 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PO4 C 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MN B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MN E 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PO4 D 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around PO4 E 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MN D 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around PO4 F 403:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around MN C 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MN D 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around MN A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around MN E 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MN C 401:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MN F 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.