



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:37 AM UTC

PDB ID : 5ZUU / pdb_00005zuu
Title : Crystal structure of AtCPSF30 YTH domain in complex with 10mer m6A-modified RNA
Authors : Wu, B.; Nie, H.; Li, S.; Patel, D.J.
Deposited on : 2018-05-08
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)
RNA backbone	3983	1010 (2.32-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	180	9% 82% 8% 10%
1	B	180	6% 76% 7% • 17%
1	C	180	8% 92% • • •
1	D	180	18% 73% 8% • 18%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	E	10	30%10%10%50%
2	F	10	30%10%10%50%
2	G	10	10%10%20%10%60%
2	I	10	10%20%20%60%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5701 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 30-kDa cleavage and polyadenylation specificity factor 30.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	174	Total	C	N	O	S	0	1	0
			1405	874	258	268	5			
1	A	162	Total	C	N	O	S	0	0	0
			1294	811	235	243	5			
1	B	149	Total	C	N	O	S	0	0	0
			1209	760	219	225	5			
1	D	147	Total	C	N	O	S	0	0	0
			1181	739	213	224	5			

- Molecule 2 is a RNA chain called RNA (5'-R(*(6MZ)P*CP*UP*AP*G)-3').


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	5	Total	C	N	O	P	0	0	0
			105	49	20	32	4			
2	F	5	Total	C	N	O	P	0	0	0
			105	49	20	32	4			
2	G	4	Total	C	N	O	P	0	0	0
			82	39	15	25	3			
2	I	4	Total	C	N	O	P	0	0	0
			82	39	15	25	3			

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: $C_8H_{18}N_2O_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	78	Total 78	O 78	0	0
5	A	48	Total 48	O 48	0	0
5	B	48	Total 48	O 48	0	0
5	D	24	Total 24	O 24	0	0
5	E	7	Total 7	O 7	0	0
5	F	10	Total 10	O 10	0	0
5	I	2	Total 2	O 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: 30-kDa cleavage and polyadenylation specificity factor 30





- Molecule 1: 30-kDa cleavage and polyadenylation specificity factor 30



- Molecule 1: 30-kDa cleavage and polyadenylation specificity factor 30




- Molecule 1: 30-kDa cleavage and polyadenylation specificity factor 30




- Molecule 2: RNA (5'-R*((6MZ)P*CP*UP*AP*(G))-3')

Chain E: 



- Molecule 2: RNA (5'-R(*(6MZ)P*CP*UP*AP*G)-3')

Chain F: 



- Molecule 2: RNA (5'-R(*(6MZ)P*CP*UP*AP*G)-3')

Chain G: 



- Molecule 2: RNA (5'-R(*(6MZ)P*CP*UP*AP*G)-3')

Chain I: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.11Å 74.29Å 151.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.95 30.00 – 1.95	Depositor EDS
% Data completeness (in resolution range)	86.3 (30.00-1.95) 86.3 (30.00-1.95)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.52 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.199 , 0.256 0.208 , 0.263	Depositor DCC
R_{free} test set	2513 reflections (4.38%)	wwPDB-VP
Wilson B-factor (Å ²)	18.5	Xtriage
Anisotropy	0.109	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 40.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5701	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 6MD, EPE, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.05	1/1322 (0.1%)	1.10	1/1787 (0.1%)
1	B	1.15	2/1232 (0.2%)	1.07	3/1661 (0.2%)
1	C	1.24	2/1434 (0.1%)	1.14	0/1936
1	D	0.98	0/1202	1.05	2/1621 (0.1%)
2	E	0.65	0/94	1.35	3/144 (2.1%)
2	F	0.92	0/94	1.77	2/144 (1.4%)
2	G	0.62	0/68	1.08	1/103 (1.0%)
2	I	0.62	0/68	1.07	1/103 (1.0%)
All	All	1.10	5/5514 (0.1%)	1.11	13/7499 (0.2%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	366	GLU	CD-OE1	6.02	1.36	1.25
1	C	249	ASN	C-O	-5.94	1.17	1.24
1	C	329	LYS	N-CA	-5.64	1.39	1.46
1	A	366	GLU	C-O	-5.52	1.17	1.24
1	B	277	GLU	C-O	-5.52	1.17	1.24

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	5	G	C1'-C2'-O2'	9.83	123.14	108.40
2	F	5	G	C3'-C2'-O2'	-8.89	97.37	110.70
2	E	5	G	C2'-C3'-O3'	-8.22	101.37	113.70
1	D	234	GLN	N-CA-C	7.35	121.50	112.54
2	G	2	C	C3'-C2'-O2'	6.64	120.67	110.70
1	A	236	VAL	N-CA-C	-5.66	99.44	107.98
2	E	5	G	C3'-C2'-O2'	-5.59	102.31	110.70
1	D	349	ILE	CB-CA-C	-5.40	107.17	113.22

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	2	C	C3'-C2'-O2'	5.40	118.79	110.70
1	B	256	GLN	N-CA-C	5.20	119.62	113.28
1	B	232	LEU	CA-C-N	-5.13	114.95	120.03
1	B	232	LEU	C-N-CA	-5.13	114.95	120.03
2	E	4	A	C1'-C2'-O2'	5.06	119.39	111.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1294	0	1261	8	0
1	B	1209	0	1190	4	0
1	C	1405	0	1363	3	0
1	D	1181	0	1163	9	0
2	E	105	0	58	0	0
2	F	105	0	58	2	0
2	G	82	0	47	2	0
2	I	82	0	47	0	0
3	A	6	0	8	0	0
4	B	15	0	18	0	0
5	A	48	0	0	0	0
5	B	48	0	0	0	0
5	C	78	0	0	1	0
5	D	24	0	0	1	0
5	E	7	0	0	0	0
5	F	10	0	0	0	0
5	I	2	0	0	0	0
All	All	5701	0	5213	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:249:ASN:HD21	1:A:311:LYS:H	1.24	0.85
1:D:332:HIS:HD2	5:D:404:HOH:O	1.72	0.71
1:C:249:ASN:HD21	1:C:311:LYS:H	1.39	0.70
1:D:270:ASN:HD21	1:D:301:ILE:HB	1.57	0.69
1:B:239:TYR:OH	1:B:366:GLU:HG3	2.02	0.59
2:F:4:A:H8	2:F:4:A:H5"	1.67	0.58
1:D:249:ASN:HB3	2:G:1:6MD:H15	1.86	0.57
1:D:269:LEU:HB3	1:D:297:MET:HE1	1.89	0.54
2:F:4:A:H5"	2:F:4:A:C8	2.43	0.53
1:A:249:ASN:HD21	1:A:311:LYS:N	2.01	0.53
1:D:232:LEU:HD11	1:D:234:GLN:NE2	2.26	0.51
1:D:351:ARG:NH1	2:G:2:C:OP1	2.38	0.50
1:A:237:ASN:OD1	1:A:278:ASN:HB2	2.12	0.50
1:D:232:LEU:HD11	1:D:234:GLN:HE21	1.77	0.49
1:D:297:MET:HE3	1:D:319:PHE:CE2	2.48	0.47
1:A:266:GLU:OE1	1:A:320:GLY:HA3	2.15	0.47
1:A:307:GLY:HA3	1:A:322:ASN:ND2	2.29	0.47
1:A:256:GLN:HE22	1:A:308:GLY:CA	2.29	0.44
1:D:283:PHE:CD1	1:D:323:TRP:CH2	3.06	0.43
1:B:289:ARG:HA	1:B:335:PHE:CE2	2.53	0.43
1:A:256:GLN:HE22	1:A:308:GLY:HA3	1.84	0.43
1:C:247:ARG:HD3	5:C:507:HOH:O	2.17	0.43
1:B:253:SER:HB2	1:B:259:TRP:HE3	1.85	0.42
1:C:247:ARG:O	1:C:248:GLU:C	2.63	0.42
1:A:304:TYR:CD1	1:A:304:TYR:C	2.97	0.42
1:B:250:PHE:CZ	1:B:292:GLN:HB3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	160/180 (89%)	154 (96%)	5 (3%)	1 (1%)	21 12

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	144/180 (80%)	140 (97%)	4 (3%)	0	100	100
1	C	173/180 (96%)	166 (96%)	5 (3%)	2 (1%)	10	4
1	D	143/180 (79%)	137 (96%)	6 (4%)	0	100	100
All	All	620/720 (86%)	597 (96%)	20 (3%)	3 (0%)	24	16

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	316	THR
1	C	314	HIS
1	C	308	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	141/156 (90%)	141 (100%)	0	100	100
1	B	134/156 (86%)	132 (98%)	2 (2%)	57	54
1	C	153/156 (98%)	151 (99%)	2 (1%)	61	58
1	D	132/156 (85%)	130 (98%)	2 (2%)	57	54
All	All	560/624 (90%)	554 (99%)	6 (1%)	65	64

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	258	VAL
1	C	318	GLN
1	B	264	SER
1	B	321	ARG
1	D	258	VAL
1	D	318	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15)

such sidechains are listed below:

Mol	Chain	Res	Type
1	C	230	HIS
1	C	249	ASN
1	A	249	ASN
1	A	255	GLN
1	A	256	GLN
1	B	249	ASN
1	B	255	GLN
1	B	340	ASN
1	B	348	ASN
1	D	234	GLN
1	D	245	ASN
1	D	249	ASN
1	D	256	GLN
1	D	318	ASN
1	D	332	HIS

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	E	3/10 (30%)	1 (33%)	0
2	F	3/10 (30%)	1 (33%)	0
2	G	2/10 (20%)	1 (50%)	0
2	I	2/10 (20%)	1 (50%)	0
All	All	10/40 (25%)	4 (40%)	0

All (4) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	E	5	G
2	F	4	A
2	G	4	A
2	I	4	A

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	501	-	5,5,5	0.50	0	5,5,5	0.73	0
4	EPE	B	501	-	15,15,15	1.74	1 (6%)	19,20,20	1.45	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	501	-	-	4/4/4/4	-
4	EPE	B	501	-	-	4/9/19/19	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	501	EPE	C10-S	-6.24	1.68	1.77

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	501	EPE	O2S-S-O1S	-3.52	102.39	113.82
4	B	501	EPE	O2S-S-C10	2.92	111.14	106.73
4	B	501	EPE	O3S-S-C10	2.30	110.51	106.00

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	GOL	O1-C1-C2-C3
4	B	501	EPE	C9-C10-S-O1S
4	B	501	EPE	C9-C10-S-O3S
3	A	501	GOL	C1-C2-C3-O3
3	A	501	GOL	O1-C1-C2-O2
3	A	501	GOL	O2-C2-C3-O3
4	B	501	EPE	N4-C7-C8-O8
4	B	501	EPE	C9-C10-S-O2S

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	162/180 (90%)	0.15	16 (9%) 13 14	13, 22, 62, 78	0
1	B	149/180 (82%)	0.08	11 (7%) 20 23	12, 23, 62, 88	0
1	C	174/180 (96%)	-0.07	14 (8%) 18 21	10, 16, 67, 107	1 (0%)
1	D	147/180 (81%)	1.13	33 (22%) 2 2	18, 45, 85, 97	0
2	E	4/10 (40%)	-0.56	0 100 100	15, 17, 24, 29	0
2	F	4/10 (40%)	-0.55	0 100 100	14, 16, 26, 31	0
2	G	3/10 (30%)	1.60	1 (33%) 1 0	66, 66, 70, 101	0
2	I	3/10 (30%)	1.21	1 (33%) 1 0	49, 49, 51, 91	0
All	All	646/760 (85%)	0.30	76 (11%) 9 10	10, 25, 74, 107	1 (0%)

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	317	ALA	8.4
1	D	232	LEU	5.8
1	C	316	THR	5.3
1	D	385	ALA	5.3
1	A	389	ALA	4.7
1	C	314	HIS	4.6
1	C	315	GLY	4.4
1	D	381	ILE	4.4
1	D	383	ALA	4.3
1	D	233	PRO	4.2
1	B	319	TYR	4.1
1	D	231	PRO	3.8
1	A	386	ALA	3.8
1	D	379	ILE	3.7
1	C	225	THR	3.6
1	D	259	TRP	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	301	ILE	3.5
1	A	228	THR	3.4
1	D	264	SER	3.3
1	D	230	HIS	3.2
1	D	267	ALA	3.2
1	B	386	ALA	3.2
1	B	310	TRP	3.2
1	B	320	GLY	3.1
1	D	302	GLY	3.1
1	C	223	ASP	3.1
1	D	260	ALA	3.1
1	A	385	ILE	3.0
1	A	306	GLY	3.0
1	B	387	ALA	3.0
1	B	390	LYS	3.0
1	C	318	GLN	3.0
1	A	316	THR	3.0
1	D	263	ARG	3.0
1	C	308	GLY	2.9
1	D	317	ARG	2.9
1	A	307	GLY	2.9
1	B	302	GLY	2.9
1	A	382	ALA	2.8
1	B	389	ALA	2.8
1	A	383	ILE	2.8
1	C	226	ASN	2.7
1	C	224	GLN	2.7
1	D	382	ALA	2.6
1	D	257	GLY	2.6
1	B	391	ARG	2.6
1	D	269	LEU	2.6
1	D	252	LEU	2.5
1	D	386	LYS	2.5
2	G	4	A	2.5
1	C	394	GLU	2.5
1	A	379	GLU	2.4
1	D	250	PHE	2.4
1	A	317	ALA	2.4
1	C	312	HIS	2.4
2	I	4	A	2.4
1	D	319	PHE	2.3
1	D	375	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	387	ALA	2.3
1	B	385	ILE	2.3
1	D	376	LEU	2.3
1	A	305	ILE	2.3
1	D	258	VAL	2.3
1	D	389	GLU	2.3
1	A	314	HIS	2.3
1	D	229	SER	2.3
1	A	304	TYR	2.2
1	D	273	PHE	2.2
1	A	315	GLY	2.2
1	D	388	GLU	2.2
1	D	234	GLN	2.2
1	B	384	SER	2.2
1	C	227	ARG	2.2
1	D	265	ASN	2.1
1	D	378	ALA	2.1
1	C	302	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EPE	B	501	15/15	0.77	0.19	56,66,80,84	0
3	GOL	A	501	6/6	0.92	0.14	36,43,47,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.