



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 7, 2026 – 02:03 AM UTC

PDB ID : 2ZVF / pdb_00002zvf
Title : Crystal structure of Archaeoglobus fulgidus alanyl-tRNA synthetase C-terminal dimerization domain
Authors : Naganuma, M.; Sekine, S.; Fukunaga, R.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2008-11-06
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	171	38% 56% 6%
1	B	171	44% 48% 7%
1	C	171	35% 56% 6%
1	D	171	39% 56% 5%
1	E	171	35% 56% 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	171	36% 56% 6% ••
1	G	171	38% 57% ••
1	H	171	% 38% 53% 8% •

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10745 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alanyl-tRNA synthetase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	170	Total	C	N	O	S	Se	0	0	0
			1341	841	236	257	1	6			
1	B	170	Total	C	N	O	S	Se	0	0	0
			1341	841	236	257	1	6			
1	C	166	Total	C	N	O	S	Se	0	0	0
			1315	827	232	249	1	6			
1	D	170	Total	C	N	O	S	Se	0	0	0
			1345	844	236	257	1	7			
1	E	170	Total	C	N	O	S	Se	0	0	0
			1345	844	236	257	1	7			
1	F	168	Total	C	N	O	S	Se	0	0	0
			1326	833	232	253	1	7			
1	G	170	Total	C	N	O	S	Se	0	0	0
			1342	841	236	258	1	6			
1	H	171	Total	C	N	O	S	Se	0	0	0
			1350	846	237	259	1	7			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MSE	-	initiating methionine	UNP O28029
B	0	MSE	-	initiating methionine	UNP O28029
C	0	MSE	-	initiating methionine	UNP O28029
D	0	MSE	-	initiating methionine	UNP O28029
E	0	MSE	-	initiating methionine	UNP O28029
F	0	MSE	-	initiating methionine	UNP O28029
G	0	MSE	-	initiating methionine	UNP O28029
H	0	MSE	-	initiating methionine	UNP O28029

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	F	1	Total	O	S	0	0
			5	4	1		
2	G	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	O	0	0
			4	4		
3	D	2	Total	O	0	0
			2	2		
3	E	1	Total	O	0	0
			1	1		
3	G	2	Total	O	0	0
			2	2		
3	H	1	Total	O	0	0
			1	1		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Alanyl-tRNA synthetase



• Molecule 1: Alanyl-tRNA synthetase



• Molecule 1: Alanyl-tRNA synthetase







• Molecule 1: Alanyl-tRNA synthetase

Chain D: 39% 56% 5%





• Molecule 1: Alanyl-tRNA synthetase

Chain E: 35% 56% 9%




• Molecule 1: Alanyl-tRNA synthetase


Chain F: 36% 56% 6%



• Molecule 1: Alanyl-tRNA synthetase

Chain G: 38% 57% ..





• Molecule 1: Alanyl-tRNA synthetase



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	124.11Å 131.71Å 138.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.20 50.00 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.1 (50.00-3.20) 99.1 (50.00-3.20)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 3.19Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.205 , 0.276 0.204 , 0.276	Depositor DCC
R_{free} test set	1898 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	76.1	Xtriage
Anisotropy	0.048	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 98.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	10745	wwPDB-VP
Average B, all atoms (Å ²)	83.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	0/1349	0.98	3/1793 (0.2%)
1	B	0.65	0/1349	1.05	2/1793 (0.1%)
1	C	0.46	0/1323	0.96	5/1759 (0.3%)
1	D	0.54	0/1353	0.93	5/1798 (0.3%)
1	E	0.52	0/1353	1.03	4/1798 (0.2%)
1	F	0.47	0/1333	0.92	3/1771 (0.2%)
1	G	0.56	0/1350	1.00	6/1793 (0.3%)
1	H	0.67	0/1358	1.03	6/1803 (0.3%)
All	All	0.56	0/10768	0.99	34/14308 (0.2%)

There are no bond length outliers.

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	890	ARG	N-CA-C	-7.93	102.62	111.82
1	G	890	ARG	N-CA-C	-7.72	103.12	112.54
1	C	890	ARG	N-CA-C	-7.41	102.36	111.40
1	H	780	ARG	N-CA-C	-7.08	103.50	111.14
1	H	890	ARG	N-CA-C	-6.86	103.73	111.14

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1341	0	1359	106	0
1	B	1341	0	1359	103	0
1	C	1315	0	1341	94	0
1	D	1345	0	1368	104	0
1	E	1345	0	1368	141	0
1	F	1326	0	1355	120	0
1	G	1342	0	1359	111	0
1	H	1350	0	1371	104	1
2	A	5	0	0	1	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
2	F	5	0	0	0	0
2	G	5	0	0	0	0
3	A	4	0	0	1	0
3	D	2	0	0	0	0
3	E	1	0	0	0	0
3	G	2	0	0	0	0
3	H	1	0	0	0	0
All	All	10745	0	10880	793	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 37.

The worst 5 of 793 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:871:GLY:HA3	1:D:884:VAL:HG22	1.26	1.11
1:B:822:LEU:HB3	1:B:849:VAL:HG21	1.31	1.08
1:G:757:ILE:HD11	1:H:769:VAL:HG12	1.36	1.08
1:E:884:VAL:HG12	1:E:886:GLN:H	1.22	1.02
1:H:877:LYS:H	1:H:877:LYS:HD2	1.24	1.01

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:870:LYS:NZ	1:H:870:LYS:NZ[4_565]	1.98	0.22

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	168/171 (98%)	148 (88%)	20 (12%)	0	100	100
1	B	168/171 (98%)	151 (90%)	17 (10%)	0	100	100
1	C	164/171 (96%)	137 (84%)	26 (16%)	1 (1%)	21	56
1	D	168/171 (98%)	151 (90%)	17 (10%)	0	100	100
1	E	168/171 (98%)	148 (88%)	18 (11%)	2 (1%)	10	42
1	F	166/171 (97%)	142 (86%)	23 (14%)	1 (1%)	21	56
1	G	168/171 (98%)	150 (89%)	17 (10%)	1 (1%)	21	56
1	H	169/171 (99%)	145 (86%)	23 (14%)	1 (1%)	21	56
All	All	1339/1368 (98%)	1172 (88%)	161 (12%)	6 (0%)	30	62

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	903	GLU
1	G	870	LYS
1	E	749	ARG
1	E	870	LYS
1	F	870	LYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	138/132 (104%)	126 (91%)	12 (9%)	9	36

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	138/132 (104%)	124 (90%)	14 (10%)	7	30
1	C	136/132 (103%)	125 (92%)	11 (8%)	11	39
1	D	139/132 (105%)	131 (94%)	8 (6%)	18	51
1	E	139/132 (105%)	125 (90%)	14 (10%)	7	30
1	F	137/132 (104%)	123 (90%)	14 (10%)	7	29
1	G	138/132 (104%)	132 (96%)	6 (4%)	26	59
1	H	139/132 (105%)	123 (88%)	16 (12%)	5	24
All	All	1104/1056 (104%)	1009 (91%)	95 (9%)	10	36

5 of 95 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	895	ASP
1	F	877	LYS
1	F	746	GLU
1	F	813	GLU
1	G	861	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	779	GLN
1	G	779	GLN
1	H	885	GLN
1	F	881	GLN
1	G	820	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	202	-	4,4,4	0.41	0	6,6,6	0.08	0
2	SO4	A	206	-	4,4,4	0.36	0	6,6,6	0.20	0
2	SO4	F	205	-	4,4,4	0.42	0	6,6,6	0.10	0
2	SO4	G	203	-	4,4,4	0.35	0	6,6,6	0.08	0
2	SO4	D	201	-	4,4,4	1.71	2 (50%)	6,6,6	0.23	0
2	SO4	C	204	-	4,4,4	0.40	0	6,6,6	0.11	0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	201	SO4	O2-S	2.29	1.58	1.44
2	D	201	SO4	O1-S	2.05	1.57	1.44

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	206	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9	
1	A	164/171 (95%)	-0.53	0	100	100	29, 72, 113, 150	0
1	B	164/171 (95%)	-0.64	0	100	100	28, 60, 103, 140	0
1	C	160/171 (93%)	-0.30	0	100	100	52, 98, 144, 169	0
1	D	163/171 (95%)	-0.45	0	100	100	51, 79, 112, 142	0
1	E	163/171 (95%)	-0.33	0	100	100	44, 81, 115, 153	0
1	F	161/171 (94%)	-0.33	0	100	100	51, 95, 134, 167	0
1	G	164/171 (95%)	-0.49	0	100	100	46, 80, 116, 152	0
1	H	164/171 (95%)	-0.55	1 (0%)	85	73	29, 65, 110, 160	0
All	All	1303/1368 (95%)	-0.45	1 (0%)	92	89	28, 80, 127, 169	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	737	GLY	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	D	201	5/5	0.82	0.15	121,127,137,137	0
2	SO4	F	205	5/5	0.84	0.10	103,106,121,124	0
2	SO4	C	204	5/5	0.88	0.09	125,125,128,134	0
2	SO4	G	203	5/5	0.88	0.10	126,127,130,135	0
2	SO4	B	202	5/5	0.91	0.11	79,103,109,114	0
2	SO4	A	206	5/5	0.94	0.09	75,82,90,95	0

6.5 Other polymers [i](#)

There are no such residues in this entry.