



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:24 AM UTC

PDB ID : 3ZVN / pdb_00003zvn
Title : The structural basis for substrate recognition by mammalian polynucleotide kinase 3' phosphatase
Authors : Garces, F.; Pearl, L.H.; Oliver, A.W.
Deposited on : 2011-07-25
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	416	11% 71% 19% • 9%
2	E	5	40% 40% 20%
2	F	5	60% 40%
2	G	5	20% 60% 40%
2	H	5	60% 40%
2	I	5	20% 80%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3683 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BIFUNCTIONAL POLYNUCLEOTIDE PHOSPHATASE/KINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	377	Total	C	N	O	S	0	0	0
			2904	1858	507	527	12			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	107	GLY	-	expression tag	UNP Q9JLV6
A	108	PRO	-	expression tag	UNP Q9JLV6
A	109	HIS	-	expression tag	UNP Q9JLV6
A	110	MET	-	expression tag	UNP Q9JLV6
A	170	ASN	ASP	engineered mutation	UNP Q9JLV6


- Molecule 2 is a DNA chain called 5'-D(*GP*TP*CP*AP*CP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	4	Total	C	N	O	P	0	0	0
			82	39	15	24	4			
2	F	5	Total	C	N	O	P	0	0	0
			101	48	18	30	5			
2	G	5	Total	C	N	O	P	0	0	0
			101	48	18	30	5			
2	H	5	Total	C	N	O	P	0	0	0
			101	48	18	30	5			
2	I	5	Total	C	N	O	P	0	0	0
			101	48	18	30	5			

- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mn	0	0
			1	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	210	Total 210	O 210	0	0
6	E	8	Total 8	O 8	0	0
6	F	11	Total 11	O 11	0	0
6	G	7	Total 7	O 7	0	0
6	H	9	Total 9	O 9	0	0
6	I	8	Total 8	O 8	0	0

- Molecule 1: BIFUNCTIONAL POLYNUCLEOTIDE PHOSPHATASE/KINASE




- Molecule 2: 5'-D(*GP*TP*CP*AP*CP)-3'

Chain H:  60% 40%



- Molecule 2: 5'-D(*GP*TP*CP*AP*CP)-3'

Chain I:  20% 80%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.27Å 72.99Å 114.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.49 – 2.15 36.49 – 2.15	Depositor EDS
% Data completeness (in resolution range)	89.2 (36.49-2.15) 89.0 (36.49-2.15)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 2.16Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.205 , 0.269 (Not available) , 0.257	Depositor DCC
R_{free} test set	1484 reflections (5.16%)	wwPDB-VP
Wilson B-factor (Å ²)	32.6	Xtriage
Anisotropy	0.785	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 50.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.028 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3683	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, MN, GOL, CSX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/2974	0.83	2/4043 (0.0%)
2	E	0.91	0/91	1.55	4/138 (2.9%)
2	F	0.85	0/112	1.43	3/170 (1.8%)
2	G	0.82	0/112	1.28	3/170 (1.8%)
2	H	0.84	0/112	1.51	4/170 (2.4%)
2	I	0.88	0/112	1.50	3/170 (1.8%)
All	All	0.54	0/3513	0.96	19/4861 (0.4%)

There are no bond length outliers.

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	1	DG	N3-C4-C5	-6.29	118.97	128.40
2	G	1	DG	N3-C4-C5	-5.98	119.43	128.40
2	F	1	DG	C2-N3-C4	5.96	120.75	111.80
2	F	1	DG	N3-C4-C5	-5.95	119.47	128.40
2	G	1	DG	C2-N3-C4	5.88	120.63	111.80
2	H	2	DT	C4'-C3'-O3'	-5.86	101.22	110.00
2	H	1	DG	C2-N3-C4	5.85	120.58	111.80
2	E	1	DG	C2-N3-C4	5.74	120.40	111.80
2	G	1	DG	N3-C4-N9	5.66	134.48	126.00
1	A	301	LYS	N-CA-C	5.61	117.20	111.14
2	I	1	DG	C2-N3-C4	5.48	120.03	111.80
2	E	1	DG	C5-C6-O6	-5.46	120.10	128.30
2	H	1	DG	N3-C4-N9	5.46	134.19	126.00
2	E	1	DG	N3-C4-C5	-5.46	120.22	128.40
1	A	181	GLY	N-CA-C	-5.42	107.73	115.64
2	I	1	DG	N3-C4-C5	-5.41	120.28	128.40
2	I	1	DG	N3-C4-N9	5.34	134.01	126.00
2	E	1	DG	N3-C4-N9	5.28	133.91	126.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	1	DG	N3-C4-N9	5.00	133.51	126.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2904	0	2835	53	0
2	E	82	0	46	1	0
2	F	101	0	57	1	0
2	G	101	0	57	3	0
2	H	101	0	57	0	0
2	I	101	0	57	5	0
3	A	1	0	0	0	0
4	A	12	0	16	2	0
5	A	27	0	12	1	0
6	A	210	0	0	5	0
6	E	8	0	0	2	0
6	F	11	0	0	1	0
6	G	7	0	0	0	0
6	H	9	0	0	0	0
6	I	8	0	0	2	0
All	All	3683	0	3137	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:4:DA:N1	6:E:2002:HOH:O	2.04	0.90
2:I:2:DT:O3'	6:I:2003:HOH:O	2.02	0.77
1:A:476:MET:HG3	2:I:4:DA:C2	2.20	0.75
1:A:268:LEU:HA	1:A:272:ALA:HB3	1.70	0.74

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:180:SER:OG	1:A:190:ASP:OD2	2.06	0.73
1:A:463:ARG:NE	6:A:2176:HOH:O	2.20	0.72
1:A:380:PHE:HB2	4:A:1523:GOL:H31	1.77	0.65
1:A:153:LEU:HB2	1:A:250:VAL:HB	1.83	0.61
1:A:195:TYR:OH	1:A:300:ARG:NH2	2.33	0.59
1:A:362:PRO:HD2	6:A:2095:HOH:O	2.02	0.58
1:A:509:ASP:OD1	1:A:510:PRO:HD2	2.03	0.57
1:A:222:GLY:HA3	1:A:253:HIS:HD2	1.70	0.57
1:A:387:SER:OG	6:A:2108:HOH:O	2.17	0.56
1:A:222:GLY:HA3	1:A:253:HIS:CD2	2.42	0.55
1:A:427:VAL:HB	1:A:428:PRO:HD3	1.90	0.54
1:A:394:ASN:OD1	1:A:396:ASP:HB2	2.08	0.53
1:A:187:SER:HB2	1:A:189:SER:H	1.73	0.53
1:A:155:PHE:HE2	1:A:239:LEU:HD12	1.74	0.52
1:A:488:PRO:HA	1:A:492:GLU:OE2	2.10	0.51
6:E:2005:HOH:O	2:I:2:DT:O4	2.16	0.50
1:A:223:ARG:HG3	2:G:3:DC:O3'	2.13	0.49
1:A:229:GLU:HA	1:A:232:LYS:HB2	1.95	0.49
1:A:146:TRP:CD2	1:A:155:PHE:HB2	2.48	0.48
1:A:223:ARG:HD2	2:G:3:DC:H2''	1.96	0.48
1:A:146:TRP:CE3	1:A:155:PHE:HB2	2.48	0.48
1:A:165:LYS:HD2	1:A:210:TYR:CE2	2.49	0.47
1:A:157:ALA:HB3	1:A:160:VAL:HG23	1.96	0.47
1:A:223:ARG:CD	2:G:3:DC:H2''	2.45	0.47
1:A:227:PRO:HG2	1:A:230:VAL:CG2	2.44	0.47
2:I:3:DC:P	6:I:2003:HOH:O	2.72	0.47
1:A:170:ASN:O	1:A:174:THR:HB	2.15	0.47
1:A:276:ILE:HD12	1:A:276:ILE:H	1.80	0.47
1:A:450:CSX:HB3	1:A:501:PRO:HA	1.97	0.46
4:A:1523:GOL:O3	5:A:1524:ADP:N7	2.46	0.46
1:A:256:LEU:HD23	1:A:465:MET:HB3	1.97	0.46
1:A:196:PRO:O	1:A:199:PRO:HD2	2.15	0.46
1:A:454:GLU:HA	1:A:454:GLU:OE1	2.15	0.46
1:A:228:ALA:O	1:A:232:LYS:N	2.40	0.46
1:A:514:ARG:HB3	1:A:514:ARG:CZ	2.46	0.45
2:I:2:DT:H2''	2:I:3:DC:O5'	2.16	0.45
1:A:166:VAL:HG22	1:A:211:LYS:HB3	1.98	0.45
1:A:394:ASN:ND2	1:A:421:ASP:O	2.50	0.45
1:A:509:ASP:CG	1:A:510:PRO:HD2	2.42	0.45
1:A:442:VAL:HB	1:A:443:PRO:HD2	1.98	0.45
1:A:152:LEU:HD11	1:A:264:MET:HG2	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:314:LEU:HD13	6:A:2049:HOH:O	2.18	0.43
1:A:219:MET:C	1:A:221:ILE:N	2.76	0.43
1:A:235:VAL:HA	1:A:238:VAL:HB	2.00	0.43
1:A:146:TRP:CZ3	1:A:155:PHE:HB2	2.54	0.43
1:A:162:PRO:HG2	1:A:274:GLU:CD	2.45	0.42
2:F:5:DC:OP2	6:F:2010:HOH:O	2.21	0.42
1:A:425:PRO:HB3	1:A:481:TYR:CE2	2.54	0.42
1:A:146:TRP:CE2	1:A:155:PHE:HB2	2.55	0.42
1:A:279:SER:HB3	1:A:282:ASP:HB2	2.02	0.41
1:A:305:PHE:HD1	6:A:2047:HOH:O	2.01	0.41
1:A:427:VAL:N	1:A:428:PRO:CD	2.83	0.41
1:A:259:LYS:HG2	1:A:312:PHE:CD1	2.56	0.41
1:A:163:GLN:HB2	1:A:211:LYS:HB2	2.03	0.40
1:A:255:GLY:HA2	1:A:465:MET:HG2	2.03	0.40
1:A:286:VAL:HA	1:A:321:ALA:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	292/348 (84%)	279 (96%)	13 (4%)	24	22

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	180	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	187	SER
1	A	201	LYS
1	A	223	ARG
1	A	235	VAL
1	A	256	LEU
1	A	270	GLU
1	A	303	LYS
1	A	347	SER
1	A	418	VAL
1	A	463	ARG
1	A	505	GLN
1	A	506	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	363	ASN
1	A	382	GLN
1	A	384	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSX	A	450	1	3,6,7	0.69	0	1,6,8	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSX	A	450	1	-	0/2/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	450	CSX	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	1525	-	5,5,5	0.34	0	5,5,5	0.35	0
5	ADP	A	1524	3	28,29,29	1.50	5 (17%)	43,45,45	1.85	11 (25%)
4	GOL	A	1523	-	5,5,5	0.36	0	5,5,5	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1525	-	-	0/4/4/4	-
5	ADP	A	1524	3	-	1/16/32/32	0/3/3/3
4	GOL	A	1523	-	-	2/4/4/4	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1524	ADP	C5-C4	4.46	1.47	1.39
5	A	1524	ADP	PA-O3A	3.75	1.63	1.59
5	A	1524	ADP	C5-C6	2.65	1.48	1.41
5	A	1524	ADP	C5-N7	-2.26	1.35	1.39
5	A	1524	ADP	C8-N7	2.23	1.36	1.31

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1524	ADP	C5-C4-N3	-5.60	119.00	126.72
5	A	1524	ADP	N3-C4-N9	4.68	135.12	127.17
5	A	1524	ADP	C2-N3-C4	3.56	120.51	111.83
5	A	1524	ADP	C4-C5-N7	-3.42	106.67	110.58
5	A	1524	ADP	N3-C2-N1	-3.32	123.55	128.58
5	A	1524	ADP	C4-N9-C8	3.30	109.21	105.74
5	A	1524	ADP	C5-N7-C8	2.81	107.87	103.45
5	A	1524	ADP	N9-C8-N7	-2.51	110.38	113.94
5	A	1524	ADP	O2A-PA-O1A	2.27	123.00	112.44
5	A	1524	ADP	C2-N1-C6	2.11	122.19	118.73
5	A	1524	ADP	O3'-C3'-C4'	-2.05	105.18	111.08

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	1523	GOL	C1-C2-C3-O3
5	A	1524	ADP	PA-O3A-PB-O3B
4	A	1523	GOL	O2-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1524	ADP	1	0
4	A	1523	GOL	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	376/416 (90%)	0.73	46 (12%) 8 9	22, 44, 76, 87	0
2	E	4/5 (80%)	-0.05	0 100 100	29, 29, 30, 36	1 (25%)
2	F	5/5 (100%)	-0.01	0 100 100	40, 42, 44, 49	0
2	G	5/5 (100%)	0.98	1 (20%) 3 3	48, 49, 51, 54	2 (40%)
2	H	5/5 (100%)	-0.23	0 100 100	36, 37, 44, 46	0
2	I	5/5 (100%)	0.09	0 100 100	40, 43, 44, 49	0
All	All	400/441 (90%)	0.70	47 (11%) 9 10	22, 44, 76, 87	3 (0%)

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	237	ALA	4.6
1	A	233	GLY	3.8
1	A	224	GLY	3.6
1	A	158	SER	3.6
1	A	149	LEU	3.6
1	A	277	PRO	3.6
1	A	209	GLY	3.5
1	A	154	VAL	3.5
1	A	302	LYS	3.5
1	A	160	VAL	3.5
1	A	507	HIS	3.1
1	A	207	ALA	3.1
1	A	471	ALA	3.1
1	A	240	GLU	3.1
1	A	146	TRP	3.0
1	A	272	ALA	2.8
1	A	222	GLY	2.8
1	A	155	PHE	2.7
1	A	235	VAL	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	228	ALA	2.5
1	A	299	GLY	2.5
1	A	441	GLY	2.5
1	A	226	LEU	2.5
1	A	317	GLY	2.5
1	A	161	LYS	2.4
1	A	183	VAL	2.4
1	A	221	ILE	2.4
1	A	274	GLU	2.3
1	A	251	ALA	2.3
1	A	232	LYS	2.3
1	A	468	PRO	2.3
1	A	469	SER	2.2
1	A	231	PHE	2.2
1	A	276	ILE	2.2
1	A	157	ALA	2.2
1	A	206	ALA	2.2
1	A	220	GLY	2.2
1	A	470	HIS	2.2
1	A	256	LEU	2.2
2	G	3	DC	2.1
1	A	150	LYS	2.1
1	A	511	ALA	2.1
1	A	268	LEU	2.0
1	A	246	PHE	2.0
1	A	152	LEU	2.0
1	A	159	GLY	2.0
1	A	164	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSX	A	450	7/8	0.92	0.11	28,32,42,48	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	A	1525	6/6	0.76	0.15	65,69,71,74	0
4	GOL	A	1523	6/6	0.93	0.09	29,30,33,33	6
5	ADP	A	1524	27/27	0.98	0.07	24,33,38,39	0
3	MN	A	901	1/1	0.99	0.05	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.