



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 07:35 PM EDT

PDB ID : 6ZVM / pdb_00006zvm
Title : Botulinum neurotoxin B2 binding domain in complex with GD1a
Authors : Davies, J.R.; Masuyer, G.; Stenmark, P.
Deposited on : 2020-07-25
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	438	8% 89% 7%
2	A	6	17% 67% 17%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4075 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Neurotoxin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	AAA	425	3677	2366	604	697	10	0	9	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	854	SER	-	expression tag	UNP Q8GR96
AAA	855	HIS	-	expression tag	UNP Q8GR96
AAA	856	MET	-	expression tag	UNP Q8GR96

- Molecule 2 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-galactopyranose-(1-4)-[N-acetyl-alpha-neuraminic acid-(2-3)]beta-D-galactopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	A	6	88	48	3	37	0	0	0

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	AAA	1	Total	C	O	0	0
			6	3	3		
3	AAA	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	296	Total	O	0	2
			298	298		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Neurotoxin



- Molecule 2: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-galactopyranose-(1-4)-[N-acetyl-alpha-neuraminic acid-(2-3)]beta-D-galactopyranose-(1-4)-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	139.64Å 56.16Å 76.79Å 90.00° 122.00° 90.00°	Depositor
Resolution (Å)	63.73 – 1.80 63.73 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.5 (63.73-1.80) 99.5 (63.73-1.80)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.06 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.164 , 0.203 0.176 , 0.207	Depositor DCC
R_{free} test set	2083 reflections (4.44%)	wwPDB-VP
Wilson B-factor (Å ²)	18.4	Xtriage
Anisotropy	0.514	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.015 for -h-2*k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4075	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SIA, NGA, BGC, GAL, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	0.92	0/3761	1.10	2/5069 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	AAA	860	ILE	CA-C-N	5.16	127.51	120.54
1	AAA	860	ILE	C-N-CA	5.16	127.51	120.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	3677	0	3549	22	1
2	A	88	0	74	2	0
3	AAA	12	0	16	0	0
4	AAA	298	0	0	5	0
All	All	4075	0	3639	23	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:1093[B]:MET:SD	4:AAA:1581:HOH:O	2.45	0.75
1:AAA:1099:TYR:CZ	1:AAA:1144:LYS:HD3	2.30	0.67
1:AAA:881[A]:TYR:CD2	1:AAA:918:ILE:HD11	2.33	0.64
1:AAA:1144:LYS:NZ	4:AAA:1405:HOH:O	2.33	0.61
1:AAA:1121:ILE:HB	1:AAA:1255:ASN:HB3	1.93	0.51
1:AAA:1265:LYS:HG2	4:AAA:1452:HOH:O	2.11	0.51
1:AAA:1149:ARG:NH2	1:AAA:1158:ASP:O	2.41	0.50
1:AAA:983:LYS:HG3	1:AAA:1026:ILE:HD12	1.94	0.50
1:AAA:857:ASN:HD22	1:AAA:859:GLU:H	1.60	0.49
1:AAA:983:LYS:CG	1:AAA:1026:ILE:HD12	2.42	0.49
1:AAA:1240[B]:ILE:HG22	4:AAA:1437:HOH:O	2.12	0.48
1:AAA:908:GLU:OE2	1:AAA:910:ARG:NE	2.48	0.47
1:AAA:1099:TYR:CE1	1:AAA:1144:LYS:HD3	2.50	0.47
1:AAA:1056[A]:MET:HE3	1:AAA:1056[A]:MET:HB2	1.85	0.45
1:AAA:1262:TRP:CE3	2:A:4:GAL:H5	2.52	0.45
1:AAA:953:ILE:HD11	1:AAA:1056[B]:MET:HE1	2.01	0.43
1:AAA:1098:TYR:CG	1:AAA:1282:PHE:HB3	2.54	0.43
1:AAA:1260:SER:HB3	1:AAA:1263:TYR:CD2	2.54	0.43
1:AAA:983:LYS:HG3	1:AAA:1026:ILE:CD1	2.49	0.42
1:AAA:867:ASN:OD1	1:AAA:869:ARG:CZ	2.68	0.41
2:A:6:SIA:H113	2:A:6:SIA:O7	2.20	0.41
1:AAA:973:TRP:CD2	1:AAA:1007:ILE:HG21	2.55	0.41
1:AAA:1164:GLU:OE2	4:AAA:1402:HOH:O	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:916:ASN:OD1	1:AAA:916:ASN:OD1[2_556]	2.13	0.07

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	428/438 (98%)	412 (96%)	15 (4%)	1 (0%)	43	31

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	1230	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	406/412 (98%)	404 (100%)	2 (0%)	81	80

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	1204	PHE
1	AAA	1265	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BGC	A	1	2	12,12,12	0.46	0	17,17,17	0.58	0
2	GAL	A	2	2	11,11,12	0.30	0	15,15,17	1.54	2 (13%)
2	NGA	A	3	2	14,14,15	0.68	0	17,19,21	1.17	2 (11%)
2	GAL	A	4	2	11,11,12	0.58	0	15,15,17	0.85	0
2	SIA	A	5	2	20,20,21	0.87	1 (5%)	21,28,31	0.98	2 (9%)
2	SIA	A	6	2	20,20,21	0.63	1 (5%)	21,28,31	1.06	2 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	A	1	2	-	0/2/22/22	0/1/1/1
2	GAL	A	2	2	-	0/2/19/22	0/1/1/1
2	NGA	A	3	2	-	1/6/23/26	0/1/1/1
2	GAL	A	4	2	-	0/2/19/22	0/1/1/1
2	SIA	A	5	2	-	2/18/34/38	0/1/1/1
2	SIA	A	6	2	-	2/18/34/38	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	5	SIA	O1B-C1	-3.22	1.20	1.30
2	A	6	SIA	O1B-C1	-2.32	1.23	1.30

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2	GAL	O3-C3-C2	-4.39	101.09	110.05
2	A	6	SIA	O1B-C1-C2	3.27	121.22	112.71
2	A	5	SIA	O1B-C1-C2	2.75	119.86	112.71
2	A	3	NGA	O3-C3-C4	2.74	116.83	110.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	3	NGA	O3-C3-C2	-2.57	104.07	109.40
2	A	6	SIA	O1A-C1-C2	-2.32	117.83	122.85
2	A	2	GAL	C1-C2-C3	2.10	112.70	109.64
2	A	5	SIA	O1B-C1-O1A	-2.00	119.53	124.08

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	5	SIA	C11-C10-N5-C5
2	A	5	SIA	O10-C10-N5-C5
2	A	6	SIA	C11-C10-N5-C5
2	A	6	SIA	O10-C10-N5-C5
2	A	3	NGA	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	4	GAL	1	0
2	A	6	SIA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	AAA	1301	-	5,5,5	0.18	0	5,5,5	0.63	0
3	GOL	AAA	1302	-	5,5,5	0.09	0	5,5,5	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	AAA	1301	-	-	2/4/4/4	-
3	GOL	AAA	1302	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	AAA	1301	GOL	C1-C2-C3-O3
3	AAA	1301	GOL	O2-C2-C3-O3
3	AAA	1302	GOL	C1-C2-C3-O3
3	AAA	1302	GOL	O1-C1-C2-O2
3	AAA	1302	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	425/438 (97%)	0.10	33 (7%) 19 17	6, 19, 51, 89	9 (2%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	1252	ASP	4.3
1	AAA	1244	TYR	3.8
1	AAA	856	MET	3.7
1	AAA	1132	ASN	3.4
1	AAA	1228	ASP	3.4
1	AAA	1032	GLY	3.4
1	AAA	1230	GLU	3.3
1	AAA	1151	SER	3.0
1	AAA	923	PHE	3.0
1	AAA	1133	TYR	2.9
1	AAA	1130	ASN	2.9
1	AAA	1049	ASP	2.7
1	AAA	1231	SER	2.6
1	AAA	924	LEU	2.6
1	AAA	917	ILE	2.6
1	AAA	1050	ARG	2.5
1	AAA	916	ASN	2.5
1	AAA	957	LYS	2.5
1	AAA	1229	GLU	2.5
1	AAA	1227	LYS	2.4
1	AAA	861	LEU	2.4
1	AAA	1270	LYS	2.4
1	AAA	1183	TYR	2.3
1	AAA	958	ASN	2.3
1	AAA	1047	ASP	2.3
1	AAA	947	HIS	2.2
1	AAA	919	PHE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	AAA	1033	GLU	2.2
1	AAA	978	ILE	2.1
1	AAA	1157	ASN	2.1
1	AAA	1204	PHE	2.0
1	AAA	1134	ILE	2.0
1	AAA	1232	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BGC	A	1	12/12	-	-	30,43,59,64	0
2	GAL	A	2	11/12	-	-	22,23,29,33	0
2	NGA	A	3	14/15	-	-	19,22,30,34	0
2	GAL	A	4	11/12	-	-	18,21,25,26	0
2	SIA	A	5	20/21	-	-	20,26,42,53	0
2	SIA	A	6	20/21	-	-	19,25,44,44	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	AAA	1302	6/6	0.87	0.18	23,56,58,66	0
3	GOL	AAA	1301	6/6	0.94	0.10	21,32,37,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.