



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 06:12 PM UTC

PDB ID : 2ZWL / pdb_00002zwl
Title : Human factor viia-tissue factor complexed with highly selective peptide inhibitor
Authors : Kadono, S.; Sakamoto, A.; Kikuchi, Y.; Oh-Eda, M.; Yabuta, N.; Koga, T.; Hattori, K.; Shiraishi, T.; Haramura, M.; Sato, H.; Ohta, M.; Kozono, T.
Deposited on : 2008-12-16
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	152	
2	H	254	
3	T	218	

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 5170 atoms, of which 30 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Factor VII light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	142	1134	683	189	247	15	0	0	0


- Molecule 2 is a protein called Factor VII heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	254	1974	1253	351	357	13	0	0	0

- Molecule 3 is a protein called Tissue factor.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	T	191	1557	993	251	308	5	0	0	0

- Molecule 4 is beta-D-glucopyranose (CCD ID: BGC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	L	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is alpha-L-fucopyranose (CCD ID: FUC) (formula: C₆H₁₂O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	L	1	Total	C	O	0	0
			10	6	4		

- Molecule 6 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	L	8	Total	Ca	0	0
			8	8		
6	H	1	Total	Ca	0	0
			1	1		

- Molecule 7 is 2-[[[(2R)-1-[[[(2S)-5-amino-1-[(4-carbamimidoylphenyl)methylamino]-1,5-dioxo-pentan-2-yl]amino]-3-(1H-indol-3-yl)-1-oxo-propan-2-yl]sulfamoyl]ethanoic acid (CCD ID: 567) (formula: C₂₆H₃₁N₇O₇S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
7	H	1	Total	C	H	N	O	S	41	0
			71	26	30	7	7	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	L	84	Total	O	0	0
			84	84		
8	H	207	Total	O	0	0
			207	207		
8	T	113	Total	O	0	0
			113	113		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Factor VII light chain



- Molecule 2: Factor VII heavy chain



- Molecule 3: Tissue factor



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	71.40Å 82.22Å 123.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.20 30.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.4 (30.00-2.20) 99.4 (30.00-2.20)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.03 (at 2.20Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.226 , 0.267 0.213 , 0.250	Depositor DCC
R_{free} test set	2631 reflections (7.03%)	wwPDB-VP
Wilson B-factor (Å ²)	22.6	Xtriage
Anisotropy	0.424	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 52.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5170	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 567, BGC, CA, FUC, CGU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	L	0.38	0/1028	0.82	2/1374 (0.1%)
2	H	0.45	0/2024	0.89	11/2755 (0.4%)
3	T	0.39	0/1591	0.83	1/2163 (0.0%)
All	All	0.42	0/4643	0.85	14/6292 (0.2%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	199	HIS	N-CA-C	-7.10	92.87	107.70
2	H	123	LEU	N-CA-C	-6.98	99.53	109.24
3	T	20	LYS	N-CA-C	-6.25	100.15	109.15
2	H	129(G)	VAL	N-CA-C	-6.12	99.49	108.54
1	L	91	CYS	N-CA-C	5.98	117.80	111.28
2	H	164	MET	N-CA-C	-5.89	102.19	110.50
1	L	41	TRP	N-CA-C	5.41	117.92	111.71
2	H	190	SER	N-CA-C	-5.35	102.07	110.36
2	H	28	PRO	N-CA-C	5.33	120.52	113.86
2	H	126	ARG	N-CA-C	5.28	116.72	111.07
2	H	51	TRP	N-CA-C	5.14	117.86	109.59
2	H	125	GLU	N-CA-C	-5.14	103.75	110.53
2	H	197	GLY	CA-C-N	5.09	124.99	119.85
2	H	197	GLY	C-N-CA	5.09	124.99	119.85

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1134	0	988	16	0
2	H	1974	0	1950	25	0
3	T	1557	0	1518	33	0
4	L	11	0	10	0	0
5	L	10	0	10	0	0
6	H	1	0	0	0	0
6	L	8	0	0	0	0
7	H	41	30	29	2	0
8	H	207	0	0	1	0
8	L	84	0	0	0	0
8	T	113	0	0	2	0
All	All	5140	30	4505	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:T:200:ARG:HG3	3:T:201:LYS:HG3	1.60	0.82
2:H:170(E):VAL:HG23	2:H:170(H):SER:HB2	1.65	0.76
3:T:20:LYS:HG2	3:T:133:LEU:HG	1.71	0.72
2:H:127:THR:O	2:H:129(B):ARG:HG2	1.91	0.70
1:L:42:ILE:HD11	8:T:247:HOH:O	1.91	0.69
1:L:69:ILE:HG23	3:T:20:LYS:HG3	1.74	0.68
2:H:170(E):VAL:CG2	2:H:170(H):SER:HB2	2.22	0.68
3:T:129:ASP:OD1	3:T:142:SER:HB2	1.94	0.67
3:T:116:PHE:HA	3:T:124:ASN:O	1.96	0.66
3:T:100:PHE:CE1	3:T:105:GLU:HG3	2.35	0.62
2:H:144:LEU:C	2:H:145:LEU:HD12	2.26	0.61
1:L:62:LYS:HE2	3:T:130:GLU:OE2	2.01	0.61
2:H:144:LEU:O	2:H:145:LEU:HD12	2.00	0.61
3:T:29:PRO:HG3	3:T:53:THR:O	2.00	0.61
2:H:45:THR:OG1	2:H:198:PRO:HG3	2.01	0.60
1:L:42:ILE:HD12	1:L:43:SER:N	2.18	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:166:GLN:O	2:H:170:GLN:HG3	2.03	0.58
1:L:105:HIS:HB2	1:L:108:THR:O	2.04	0.57
3:T:136:ARG:HG3	3:T:141:LEU:HD11	1.88	0.56
2:H:77:ASP:O	2:H:80:GLU:HG2	2.06	0.55
2:H:34:LEU:HD23	2:H:40:GLN:HA	1.87	0.55
3:T:76:PHE:CD1	3:T:94:TYR:HB3	2.42	0.55
2:H:89:ILE:HG21	2:H:241:LEU:HD13	1.89	0.54
1:L:101:TYR:CZ	2:H:125:GLU:HG3	2.43	0.54
1:L:13:LEU:HA	1:L:30:ILE:HD13	1.90	0.53
1:L:110:ARG:HD2	1:L:110:ARG:C	2.35	0.51
3:T:32:GLN:HA	3:T:78:TYR:O	2.12	0.50
3:T:16:SER:HA	3:T:20:LYS:O	2.10	0.50
3:T:20:LYS:HG2	3:T:133:LEU:CG	2.40	0.50
3:T:136:ARG:O	3:T:137:ASN:HB2	2.12	0.50
3:T:121:THR:HG22	3:T:122:LYS:HE2	1.95	0.49
2:H:144:LEU:HD21	2:H:152:ALA:HB2	1.94	0.49
1:L:101:TYR:CE2	2:H:125:GLU:HG3	2.47	0.48
3:T:29:PRO:HB2	3:T:32:GLN:O	2.13	0.48
3:T:136:ARG:NE	3:T:141:LEU:HD11	2.29	0.48
2:H:256:PHE:HA	2:H:257:PRO:C	2.39	0.47
1:L:16:CGU:O	1:L:22:CYS:HB3	2.14	0.47
2:H:87:GLN:HG3	2:H:250:VAL:O	2.15	0.47
3:T:48:LYS:HD2	3:T:62:GLU:HG3	1.97	0.47
1:L:4:PHE:CE2	1:L:5:LEU:HG	2.50	0.47
2:H:256:PHE:CD1	2:H:257:PRO:HA	2.50	0.47
3:T:72:LEU:HD13	3:T:73:ALA:N	2.29	0.46
3:T:167:THR:HG23	8:T:292:HOH:O	2.15	0.46
1:L:4:PHE:CD2	1:L:5:LEU:HG	2.52	0.45
2:H:147:ARG:O	2:H:147:ARG:HG2	2.17	0.45
3:T:186:CYS:HA	3:T:209:CYS:HA	1.98	0.45
3:T:158:TRP:CE2	3:T:186:CYS:HB2	2.52	0.45
3:T:185:TYR:H	3:T:210:MET:HG2	1.81	0.45
1:L:70:CYS:O	3:T:20:LYS:HE3	2.17	0.45
3:T:170:THR:HG22	3:T:175:PHE:CE1	2.52	0.45
1:L:62:LYS:HD3	1:L:62:LYS:C	2.42	0.44
1:L:118:TYR:OH	1:L:137:LYS:HD3	2.19	0.43
2:H:192:LYS:HD3	7:H:1:567:C8	2.49	0.43
3:T:121:THR:CG2	3:T:122:LYS:HE2	2.48	0.43
3:T:125:VAL:HB	3:T:175:PHE:HB2	2.01	0.42
3:T:18:ASN:HD21	3:T:109:GLY:HA3	1.84	0.42
3:T:101:THR:O	3:T:105:GLU:HG2	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:46:LEU:HD11	2:H:48:ASN:O	2.20	0.42
3:T:33:VAL:CG2	3:T:78:TYR:HB2	2.49	0.42
2:H:235:ILE:O	2:H:239:GLN:HG3	2.20	0.41
1:L:139:PRO:HG2	8:H:272:HOH:O	2.20	0.41
2:H:59:PHE:HA	2:H:60(B):ILE:HG12	2.01	0.41
2:H:187:GLY:HA2	2:H:221(A):ALA:O	2.21	0.41
3:T:33:VAL:HG22	3:T:78:TYR:HB2	2.03	0.41
3:T:110:GLN:HA	3:T:111:PRO:HD3	1.96	0.41
3:T:155:LEU:O	3:T:167:THR:HA	2.20	0.41
2:H:215:TRP:HB2	7:H:1:567:O1	2.21	0.40
2:H:240:LYS:O	2:H:240:LYS:HD3	2.21	0.40
2:H:32:LEU:HD11	2:H:40:GLN:HG3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	130/152 (86%)	120 (92%)	10 (8%)	0	100	100
2	H	252/254 (99%)	241 (96%)	10 (4%)	1 (0%)	30	34
3	T	185/218 (85%)	174 (94%)	9 (5%)	2 (1%)	11	10
All	All	567/624 (91%)	535 (94%)	29 (5%)	3 (0%)	24	27

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	170(F)	GLY
3	T	137	ASN
3	T	120	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	114/122 (93%)	112 (98%)	2 (2%)	51	68
2	H	216/216 (100%)	214 (99%)	2 (1%)	70	84
3	T	180/199 (90%)	177 (98%)	3 (2%)	53	69
All	All	510/537 (95%)	503 (99%)	7 (1%)	59	75

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	18	LYS
1	L	52	SER
2	H	100	ASN
2	H	159	ASN
3	T	20	LYS
3	T	27	PRO
3	T	165	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	49	GLN
1	L	66	GLN
2	H	30	GLN
2	H	37	ASN
2	H	60(D)	ASN
2	H	63	ASN
2	H	100	ASN
2	H	110	GLN
2	H	159	ASN
2	H	166	GLN
2	H	233	GLN
3	T	11	ASN
3	T	69	GLN
3	T	107	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	T	124	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

10 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CGU	L	7	1,6	9,11,12	1.52	2 (22%)	10,14,16	0.86	1 (10%)
1	CGU	L	35	1	9,11,12	1.18	2 (22%)	10,14,16	0.53	0
1	CGU	L	20	1,6	9,11,12	1.27	2 (22%)	10,14,16	1.23	2 (20%)
1	CGU	L	19	1,6	9,11,12	1.29	2 (22%)	10,14,16	0.94	0
1	CGU	L	26	1,6	9,11,12	1.54	2 (22%)	10,14,16	0.80	0
1	CGU	L	14	1,6	9,11,12	1.30	2 (22%)	10,14,16	0.91	0
1	CGU	L	6	1,6	9,11,12	1.44	2 (22%)	10,14,16	0.79	0
1	CGU	L	25	1,6	9,11,12	1.19	2 (22%)	10,14,16	0.97	1 (10%)
1	CGU	L	16	1,6	9,11,12	1.53	2 (22%)	10,14,16	1.25	1 (10%)
1	CGU	L	29	1,6	9,11,12	1.53	2 (22%)	10,14,16	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CGU	L	7	1,6	-	2/13/14/16	-
1	CGU	L	35	1	-	7/13/14/16	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CGU	L	20	1,6	-	0/13/14/16	-
1	CGU	L	19	1,6	-	3/13/14/16	-
1	CGU	L	26	1,6	-	2/13/14/16	-
1	CGU	L	14	1,6	-	0/13/14/16	-
1	CGU	L	6	1,6	-	4/13/14/16	-
1	CGU	L	25	1,6	-	5/13/14/16	-
1	CGU	L	16	1,6	-	4/13/14/16	-
1	CGU	L	29	1,6	-	0/13/14/16	-

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	L	7	CGU	OE12-CD1	-3.47	1.19	1.30
1	L	29	CGU	OE22-CD2	-3.39	1.19	1.30
1	L	16	CGU	OE12-CD1	-3.28	1.20	1.30
1	L	26	CGU	OE22-CD2	-3.07	1.20	1.30
1	L	6	CGU	OE22-CD2	-3.02	1.21	1.30
1	L	26	CGU	OE12-CD1	-2.90	1.21	1.30
1	L	16	CGU	OE22-CD2	-2.76	1.21	1.30
1	L	14	CGU	OE12-CD1	-2.73	1.22	1.30
1	L	20	CGU	OE12-CD1	-2.71	1.22	1.30
1	L	19	CGU	OE12-CD1	-2.68	1.22	1.30
1	L	7	CGU	OE22-CD2	-2.59	1.22	1.30
1	L	6	CGU	OE12-CD1	-2.59	1.22	1.30
1	L	29	CGU	OE12-CD1	-2.58	1.22	1.30
1	L	20	CGU	OE22-CD2	-2.30	1.23	1.30
1	L	14	CGU	OE22-CD2	-2.29	1.23	1.30
1	L	25	CGU	OE12-CD1	-2.27	1.23	1.30
1	L	19	CGU	OE22-CD2	-2.19	1.23	1.30
1	L	25	CGU	OE22-CD2	-2.13	1.23	1.30
1	L	35	CGU	OE12-CD1	-2.05	1.24	1.30
1	L	35	CGU	OE22-CD2	-2.02	1.24	1.30

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	16	CGU	CB-CG-CD1	-3.01	106.99	113.11
1	L	20	CGU	CB-CA-C	-2.48	107.17	110.99
1	L	20	CGU	CB-CG-CD2	-2.36	108.32	113.11
1	L	7	CGU	CB-CG-CD1	-2.18	108.68	113.11
1	L	25	CGU	CB-CG-CD1	-2.06	108.93	113.11

There are no chirality outliers.

All (27) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	L	6	CGU	CA-CB-CG-CD1
1	L	6	CGU	CA-CB-CG-CD2
1	L	25	CGU	CA-CB-CG-CD2
1	L	35	CGU	CA-CB-CG-CD1
1	L	6	CGU	OE21-CD2-CG-CB
1	L	6	CGU	OE22-CD2-CG-CB
1	L	7	CGU	OE21-CD2-CG-CB
1	L	7	CGU	OE22-CD2-CG-CB
1	L	16	CGU	OE21-CD2-CG-CB
1	L	16	CGU	OE22-CD2-CG-CB
1	L	25	CGU	OE11-CD1-CG-CB
1	L	25	CGU	OE12-CD1-CG-CB
1	L	26	CGU	OE21-CD2-CG-CB
1	L	26	CGU	OE22-CD2-CG-CB
1	L	35	CGU	OE21-CD2-CG-CB
1	L	35	CGU	OE22-CD2-CG-CB
1	L	16	CGU	OE21-CD2-CG-CD1
1	L	16	CGU	OE22-CD2-CG-CD1
1	L	19	CGU	OE11-CD1-CG-CD2
1	L	35	CGU	OE21-CD2-CG-CD1
1	L	35	CGU	OE22-CD2-CG-CD1
1	L	19	CGU	OE11-CD1-CG-CB
1	L	19	CGU	OE12-CD1-CG-CB
1	L	25	CGU	OE11-CD1-CG-CD2
1	L	25	CGU	OE12-CD1-CG-CD2
1	L	35	CGU	OE11-CD1-CG-CD2
1	L	35	CGU	OE12-CD1-CG-CD2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	L	16	CGU	1	0

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 12 ligands modelled in this entry, 9 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	567	H	1	-	42,43,43	1.91	10 (23%)	54,60,60	1.38	9 (16%)
5	FUC	L	1060	1	10,10,11	0.66	0	14,14,16	0.70	0
4	BGC	L	1052	1	11,11,12	0.56	0	15,15,17	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	567	H	1	-	-	4/39/40/40	0/3/3/3
5	FUC	L	1060	1	-	-	0/1/1/1
4	BGC	L	1052	1	-	1/2/19/22	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	H	1	567	S1-N4	6.75	1.72	1.61
7	H	1	567	C9-C8	6.43	1.58	1.51
7	H	1	567	C10-C7	2.95	1.43	1.38
7	H	1	567	C6-C11	2.89	1.43	1.38
7	H	1	567	C4-C3	2.63	1.44	1.41
7	H	1	567	C13-C12	2.46	1.40	1.36
7	H	1	567	C10-C6	2.29	1.43	1.38
7	H	1	567	O7-C8	-2.21	1.23	1.30
7	H	1	567	C23-C22	2.14	1.42	1.39
7	H	1	567	C5-C14	2.01	1.58	1.52

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	H	1	567	C21-N6-C20	-3.79	117.09	122.29
7	H	1	567	C7-C3-C4	-3.52	118.78	122.19
7	H	1	567	C22-C26-N1	2.98	122.54	118.01
7	H	1	567	O7-C8-C9	2.96	118.50	112.83
7	H	1	567	O3-S1-N4	-2.79	103.14	107.11
7	H	1	567	O4-S1-N4	-2.67	103.30	107.11
7	H	1	567	C7-C3-N7	2.33	134.47	130.74
7	H	1	567	O6-C8-C9	-2.25	118.67	122.86
7	H	1	567	O3-S1-C9	2.17	111.71	108.15

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	L	1052	BGC	O5-C5-C6-O6
7	H	1	567	C17-C22-C26-N1
7	H	1	567	C23-C22-C26-N1
7	H	1	567	C17-C22-C26-N2
7	H	1	567	C23-C22-C26-N2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	H	1	567	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	132/152 (86%)	0.38	5 (3%) 44 41	15, 29, 45, 51	0
2	H	254/254 (100%)	-0.24	6 (2%) 59 56	8, 16, 37, 52	0
3	T	191/218 (87%)	0.57	29 (15%) 5 4	13, 27, 56, 69	0
All	All	577/624 (92%)	0.17	40 (6%) 23 20	8, 23, 49, 69	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	170(G)	ASP	6.3
3	T	137	ASN	5.2
3	T	160	SER	4.9
1	L	107	GLY	4.1
3	T	156	TYR	3.9
1	L	4	PHE	3.8
3	T	119	VAL	3.6
2	H	170(E)	VAL	3.3
3	T	136	ARG	3.2
2	H	147	ARG	3.2
3	T	31	ASN	3.2
3	T	182	GLY	3.1
3	T	30	VAL	3.0
3	T	171	ASN	3.0
1	L	108	THR	2.9
3	T	116	PHE	2.9
3	T	164	GLY	2.8
3	T	198	VAL	2.8
3	T	130	GLU	2.8
1	L	106	THR	2.7
3	T	32	GLN	2.7
3	T	138	ASN	2.7
3	T	159	LYS	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	T	181	LYS	2.6
3	T	120	GLY	2.6
3	T	28	LYS	2.6
3	T	122	LYS	2.5
3	T	197	THR	2.4
3	T	184	ASN	2.4
2	H	247	ARG	2.4
3	T	180	ASP	2.3
3	T	41	LYS	2.3
3	T	29	PRO	2.2
2	H	170(H)	SER	2.1
1	L	88	GLN	2.1
2	H	170(D)	LYS	2.0
3	T	210	MET	2.0
3	T	135	ARG	2.0
3	T	131	ARG	2.0
3	T	165	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CGU	L	35	12/13	0.59	0.22	42,52,55,56	0
1	CGU	L	19	12/13	0.79	0.12	32,36,39,41	0
1	CGU	L	14	12/13	0.81	0.12	28,35,37,40	0
1	CGU	L	29	12/13	0.85	0.11	27,29,36,37	0
1	CGU	L	25	12/13	0.87	0.12	25,38,42,43	0
1	CGU	L	20	12/13	0.88	0.10	28,30,33,33	0
1	CGU	L	26	12/13	0.88	0.11	24,26,26,28	0
1	CGU	L	7	12/13	0.90	0.09	23,24,28,28	0
1	CGU	L	6	12/13	0.90	0.11	19,25,26,27	0
1	CGU	L	16	12/13	0.91	0.10	20,24,29,29	0

6.3 Carbohydrates ⓘ


There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	BGC	L	1052	11/12	0.81	0.18	45,46,48,51	0
6	CA	L	158	1/1	0.83	0.18	52,52,52,52	0
5	FUC	L	1060	10/11	0.86	0.10	35,36,39,39	0
6	CA	L	160	1/1	0.86	0.14	40,40,40,40	0
6	CA	H	258	1/1	0.91	0.07	33,33,33,33	0
6	CA	L	159	1/1	0.94	0.15	43,43,43,43	0
7	567	H	1	41/41	0.94	0.07	10,11,17,18	41
6	CA	L	157	1/1	0.95	0.08	31,31,31,31	0
6	CA	L	154	1/1	0.95	0.08	32,32,32,32	0
6	CA	L	153	1/1	0.96	0.07	30,30,30,30	0
6	CA	L	155	1/1	0.96	0.07	29,29,29,29	0
6	CA	L	156	1/1	0.98	0.06	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.