



wwPDB EM Validation Summary Report ⓘ

Mar 12, 2026 – 08:34 AM UTC

PDB ID : 6ZWM / pdb_00006zwm
EMDB ID : EMD-11488
Title : cryo-EM structure of human mTOR complex 2, overall refinement
Authors : Scaiola, A.; Mangia, F.; Inseng, S.; Boehringer, D.; Ban, N.; Maier, T.
Deposited on : 2020-07-28
Resolution : 3.20 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	15020 (2.70 - 3.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	2549	
1	B	2549	
2	C	326	
2	D	326	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	1708	15% 57% 8% 35%
3	F	1708	58% 7% 35%
4	G	522	10% 18% 81%
4	H	522	5% 17% 81%

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 56947 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase mTOR.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	2184	Total	C	N	O	S	0	0
			16337	10356	2906	2977	98		
1	B	2185	Total	C	N	O	S	0	0
			16304	10330	2904	2972	98		

- Molecule 2 is a protein called Target of rapamycin complex subunit LST8.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	C	319	Total	C	N	O	S	0	0
			2465	1533	437	477	18		
2	D	319	Total	C	N	O	S	0	0
			2465	1533	437	477	18		


- Molecule 3 is a protein called Rapamycin-insensitive companion of mTOR.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	E	1117	Total	C	N	O	S	0	0
			8931	5689	1584	1611	47		
3	F	1117	Total	C	N	O	S	0	0
			8931	5689	1584	1611	47		

- Molecule 4 is a protein called Target of rapamycin complex 2 subunit MAPKAP1.


Mol	Chain	Residues	Atoms					AltConf	Trace
4	G	98	Total	C	N	O	S	0	0
			655	399	125	127	4		
4	H	98	Total	C	N	O	S	0	0
			655	399	125	127	4		

- Molecule 5 is PHOSPHOTHIOPHOSPHORIC ACID-ADENYLATE ESTER (CCD ID: AGS) (formula: C₁₀H₁₆N₅O₁₂P₃S) (labeled as "Ligand of Interest" by depositor).



Id	Chain	Residues	Atoms						AltConf
5	A	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
5	B	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
5	E	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
5	F	1	Total 31	C 10	N 5	O 12	P 3	S 1	0

- Molecule 6 is INOSITOL HEXAKISPHOSPHATE (CCD ID: IHP) (formula: $\text{C}_6\text{H}_{18}\text{O}_{24}\text{P}_6$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				AltConf
6	A	1	Total	C	O	P	0
			36	6	24	6	
6	B	1	Total	C	O	P	0
			36	6	24	6	

- Molecule 7 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
7	E	1	Total	Zn	0
			1	1	
7	F	1	Total	Zn	0
			1	1	

- Molecule 8 is ACETYL GROUP (CCD ID: ACE) (formula: C₂H₄O).




Mol	Chain	Residues	Atoms			AltConf
8	G	1	Total	C	O	0
			3	2	1	
8	H	1	Total	C	O	0
			3	2	1	


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Serine/threonine-protein kinase mTOR









• Molecule 2: Target of rapamycin complex subunit LST8




• Molecule 2: Target of rapamycin complex subunit LST8





• Molecule 3: Rapamycin-insensitive companion of mTOR



PHE
LYS
LEU
THR
TYR
LEU
SER
ASN
HIS
ASP
TYR
LYS
HIS
LEU
TYR
PHE
GLU
SER
SER
ASP
ALA
ALA
THR
THR
VAL
ASN
GLU
ILE
VAL
LEU
LYS
VAL
ASN
TYR
ILE
GLU
GLY
SER
SER
ARG
ALA
SER
THR
ALA
GLN
LYS
GLN
GLN
ARG
LYS
LEU
ASN
ARG
THR
PHE
SER

PHE
GLN
GLU
GLY
LYS
SER
GLY
GLN

● Molecule 4: Target of rapamycin complex 2 subunit MAPKAP1



MET
A2
N6
Q16
D22
D23
T24
G25
N26
D35
V36
ASP
LEU
GLU
LYS
ILE
HIS
PRO
PRO
SER
MET
PRO
GLY
ASP
SER
GLY
SER
ASN
GLY
THR
GLN
GLY
TYR
VAL
TYR
ALA
GLN
SER
VAL
ASP
ILE
THR
SER
TRP
ASP
PHE
ILE
ARG

ARG
ARG
S84
N85
E91
A95
Q100
I101
N105
E110
R111
N112
Q115
S116
A117
Q118
E119
F124
L129
K130
E131
K132
P133
P134
I135
S136
G137
K138
Q139
S140
R145
L146
GLU
CYS
PRO
LEU
GLN
LEU
ASN
ASN
PHE
ASN
GLU
TYR
SER
LYS
PHE
ASP
GLY

LYS
GLY
HIS
VAL
GLY
THR
THR
ALA
THR
LYS
LYS
ILE
ASP
VAL
TYR
PRO
LEU
HIS
SER
SER
GLN
ASP
ARG
LEU
LEU
PRO
MET
THR
VAL
VAL
THR
VAL
PHE
THR
MET
THR

ASN
VAL
SER
ALA
TYR
CYS
LEU
HIS
ASN
ALA
GLU
ASP
LYS
GLY
GLU
VAL
ASP
THR
ASP
PHE
PRO
PRO
LEU
LYS
VAL
SER
ASN
PRO
ILE
HIS
LYS
PHE
GLY
PHE
SER
GLY
PRO
THR
LEU
ALA
VAL
LEU
VAL
GLY
LYS
TYR
SER
SER
PRO
GLY
LEU
THR
SER
LYS
GLU
THR
ASP
VAL
LEU
LEU
SER
PHE
VAL
THR
GLU
ILE
ASN
ALA

ALA
HIS
GLY
PHE
SER
LEU
ILE
GLN
VAL
ILE
ASN
ASN
THR
LYS
VAL
VAL
ASP
THR
MET
LYS
GLU
ILE
LEU
LEU
PHE
GLU
ASP
LEU
LYS
GLN
ALA
VAL
ILE
ASP
ILE
ARG
GLY
LYS
GLN
LYS
VAL
GLY
SER
GLN
LYS
VAL
MET
LEU
LEU
SER
SER
GLY
PRO
TYR
ARG
LEU
GLY
LYS
PHE
LYS
GLN
SER
GLU
GLU
THR
PHE
VAL
VAL
ASP
LEU
LEU
THR
ASP
SER
LEU
THR
VAL
THR
GLU
LEU
LEU
THR
VAL
THR
GLN
LEU
ASN
HIS
PHE
ARG
LEU
VAL
VAL
ASP
PHE
THR
THR
ASP
SER
SER
SER
SER
VAL
GLN
LEU
GLY
ILE
GLN
SER

ALA
TRP
GLU
PHE
CYS
LEU
VAL
ARG
GLU
VAL
THR
ASN
SER
SER
ARG
ALA
ASP
THR
GLY
VAL
PHE
TRP
ILE
ASP
PHE
ILE
ALA
ASP
LEU
LEU
MET
CYS
ALA
CYS
SER
HIS
HIS
TYR
LYS
SER
PHE
ILE
HIS
ARG
PHE
LYS
LEU
THR
PHE
THR
LEU
SER
SER
SER
SER
HIS
ALA
ILE
HIS
PHE
PHE
LYS
LEU
THR
THR
LYS
LYS
GLY
ASN
HIS
ASP
TYR
LYS
HIS

GLY
ASP
LYS
VAL
GLU
ILE
ASP
PRO
VAL
THR
ASN
GLN
LYS
ALA
ILE
SER
SER
ILE
ILE
ASP
LEU
LEU
CYS
ALA
GLN
LYS
ASN
ARG
THR
PHE
PHE
GLY
THR
LEU
LEU
ASN
HIS
ASP
TYR
LYS
HIS

LEU
TYR
PHE
GLU
SER
ASP
ALA
ALA
THR
VAL
GLU
ILE
VAL
LEU
LYS
VAL
ASN
TYR
ILE
LEU
GLU
SER
ARG
ALA
SER
THR
ALA
ASP
LEU
LEU
PHE
GLN
LYS
GLN
ARG
LYS
LEU
ASN
ARG
THR
SER
PHE
SER
PHE
PHE
GLN
LYS
GLU
LYS
LYS
SER
SER
GLY
GLN

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	293038	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	70	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	3.323	Depositor
Minimum map value	-1.756	Depositor
Average map value	-0.002	Depositor
Map value standard deviation	0.077	Depositor
Recommended contour level	0.375	Depositor
Map size ($\text{\AA}$)	430.08, 430.08, 430.08	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.344, 1.344, 1.344	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IHP, AGS, ZN, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.20	0/16632	0.46	0/22593
1	B	0.19	0/16598	0.44	0/22552
2	C	0.20	0/2523	0.54	0/3438
2	D	0.19	0/2523	0.53	0/3438
3	E	0.22	0/9092	0.51	0/12300
3	F	0.20	0/9092	0.46	1/12300 (0.0%)
4	G	0.22	0/660	0.63	1/900 (0.1%)
4	H	0.20	0/660	0.58	0/900
All	All	0.20	0/57780	0.48	2/78421 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	E	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	G	25	GLY	N-CA-C	6.26	120.21	112.64
3	F	141	VAL	N-CA-C	-5.36	107.60	112.96

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	E	367	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	16337	0	15369	117	0
1	B	16304	0	15298	107	0
2	C	2465	0	2351	33	0
2	D	2465	0	2351	31	0
3	E	8931	0	9080	81	0
3	F	8931	0	9080	72	0
4	G	655	0	519	2	0
4	H	655	0	519	7	0
5	A	31	0	12	1	0
5	B	31	0	12	0	0
5	E	31	0	12	1	0
5	F	31	0	12	1	0
6	A	36	0	6	2	0
6	B	36	0	6	2	0
7	E	1	0	0	0	0
7	F	1	0	0	0	0
8	G	3	0	3	0	0
8	H	3	0	3	0	0
All	All	56947	0	54633	432	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 432 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:552:GLY:O	3:E:556:LYS:HB2	1.75	0.87
3:E:920:ILE:HG23	3:E:964:ILE:HD11	1.75	0.69
1:A:1211:VAL:HA	3:E:553:THR:HG21	1.75	0.67
3:E:555:LEU:HD11	3:E:607:GLN:HG3	1.76	0.66
2:D:134:VAL:HG12	2:D:145:VAL:HG12	1.82	0.62

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	2143/2549 (84%)	2080 (97%)	62 (3%)	1 (0%)	100	100
1	B	2145/2549 (84%)	2071 (97%)	73 (3%)	1 (0%)	100	100
2	C	317/326 (97%)	299 (94%)	18 (6%)	0	100	100
2	D	317/326 (97%)	291 (92%)	26 (8%)	0	100	100
3	E	1103/1708 (65%)	1066 (97%)	37 (3%)	0	100	100
3	F	1103/1708 (65%)	1067 (97%)	36 (3%)	0	100	100
4	G	94/522 (18%)	86 (92%)	8 (8%)	0	100	100
4	H	94/522 (18%)	85 (90%)	9 (10%)	0	100	100
All	All	7316/10210 (72%)	7045 (96%)	269 (4%)	2 (0%)	100	100

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	121	VAL
1	A	121	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1563/2220 (70%)	1543 (99%)	20 (1%)	61	78

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	1552/2220 (70%)	1527 (98%)	25 (2%)	55	75
2	C	269/276 (98%)	260 (97%)	9 (3%)	33	65
2	D	269/276 (98%)	263 (98%)	6 (2%)	45	71
3	E	987/1539 (64%)	975 (99%)	12 (1%)	63	79
3	F	987/1539 (64%)	973 (99%)	14 (1%)	59	77
4	G	50/471 (11%)	48 (96%)	2 (4%)	28	61
4	H	50/471 (11%)	47 (94%)	3 (6%)	17	50
All	All	5727/9012 (64%)	5636 (98%)	91 (2%)	54	75

5 of 91 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	147	ASP
3	E	1667	LEU
2	D	308	TYR
3	E	691	LEU
3	F	441	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 105 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	118	ASN
3	E	748	ASN
3	F	881	HIS
3	E	49	GLN
3	E	264	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	AGS	E	4802	-	32,33,33	1.73	8 (25%)	45,52,52	2.16	16 (35%)
8	ACE	G	601	4	1,2,2	0.82	0	0,1,1	-	-
5	AGS	F	4802	-	32,33,33	1.72	8 (25%)	45,52,52	2.18	14 (31%)
6	IHP	B	2602	-	36,36,36	1.51	3 (8%)	60,60,60	1.08	5 (8%)
5	AGS	A	2601	-	32,33,33	1.71	8 (25%)	45,52,52	2.09	14 (31%)
6	IHP	A	2602	-	36,36,36	1.52	3 (8%)	60,60,60	1.09	6 (10%)
5	AGS	B	2601	-	32,33,33	1.73	8 (25%)	45,52,52	2.09	14 (31%)
8	ACE	H	601	4	1,2,2	0.73	0	0,1,1	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	AGS	E	4802	-	-	3/21/38/38	0/3/3/3
5	AGS	F	4802	-	-	3/21/38/38	0/3/3/3
6	IHP	B	2602	-	-	6/30/54/54	0/1/1/1
5	AGS	A	2601	-	-	6/21/38/38	0/3/3/3
6	IHP	A	2602	-	-	7/30/54/54	0/1/1/1
5	AGS	B	2601	-	-	6/21/38/38	0/3/3/3

The worst 5 of 38 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	2602	IHP	P4-O24	3.82	1.62	1.50
6	B	2602	IHP	P4-O24	3.82	1.62	1.50
5	E	4802	AGS	C2-N3	3.56	1.40	1.33
5	F	4802	AGS	C2-N3	3.55	1.40	1.33
5	A	2601	AGS	C2-N3	3.51	1.40	1.33

The worst 5 of 69 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	2601	AGS	N3-C2-N1	-7.09	117.85	128.58
5	F	4802	AGS	N3-C2-N1	-7.08	117.86	128.58
5	B	2601	AGS	N3-C2-N1	-7.08	117.87	128.58
5	E	4802	AGS	N3-C2-N1	-7.04	117.93	128.58
5	E	4802	AGS	C5-C4-N3	-4.67	120.29	126.72

There are no chirality outliers.

5 of 31 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	2601	AGS	PB-O3B-PG-O3G
5	B	2601	AGS	PB-O3B-PG-O3G
5	E	4802	AGS	C5'-O5'-PA-O1A
5	F	4802	AGS	C5'-O5'-PA-O1A
5	F	4802	AGS	O4'-C1'-N9-C8


There are no ring outliers.


5 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	E	4802	AGS	1	0
5	F	4802	AGS	1	0
6	B	2602	IHP	2	0
5	A	2601	AGS	1	0
6	A	2602	IHP	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring


in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-11488. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 160




Y Index: 160




Z Index: 160


6.2.2 Raw map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 179




Y Index: 182




Z Index: 157


6.3.2 Raw map



X Index: 176



Y Index: 163




Z Index: 159


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.375. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_11488_msk_2.map [i](#)



X




Y




Z


6.6.2 emd_11488_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 419 nm^3 ; this corresponds to an approximate mass of 379 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.312 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.312 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.20	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	4.22	7.53	4.34

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.22 differs from the reported value 3.2 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-11488 and PDB model 6ZWM. Per-residue inclusion information can be found in section [3](#) on page [7](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.375 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.375).

9.4 Atom inclusion [i](#)



At the recommended contour level, 80% of all backbone atoms, 72% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.375) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7250	0.3960
A	0.7370	0.3860
B	0.7520	0.4130
C	0.6640	0.3020
D	0.7610	0.3740
E	0.5830	0.3580
F	0.8290	0.4600
G	0.4070	0.3060
H	0.6470	0.3700

